

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Top10 Countability and Separation Axioms

KYB

Thrn, it's a Fact

mathrnfact@gmail.com

May 18, 2020

Overview

30.The Countability Axioms

Definition

Theorem

Exercise

31.The Separation Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

30.The Countability Axioms

Definition

Theorem

Exercise

31.The Separation Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Definition

Definition (countable basis at x)

if there is cble coll \mathcal{B} of nbd of x s.t. \forall nbd of x contains at least one of ele of \mathcal{B}

Definition (first cble axiom)

if X has cble basis at each x

Definition (second cble axiom)

if X has a cble basis

Definition (dense)

$A \subset X$ is dense if $\bar{A} = X$

Theorem

Theorem (30.1)

- ▶ (a) $A \subset X$. If $\exists \{x_n\}$ in A s.t. $x_n \rightarrow x$, then $x \in \bar{A}$.
The converse holds if X is first-cble.
- ▶ (b) $f: X \rightarrow Y$ fct. If f is conti, then for every convergent seq $x_n \rightarrow x$ in X , $f(x_n) \rightarrow f(x)$. The converse holds if X is first-cble.

Theorem (30.2)

- ▶ subspace, cble product of first(second)-cble is again first(secon)-cble.

30.The Countability Axioms

[Definition](#)[Theorem](#)[Exercise](#)

31.The Separation Axioms

[Definition](#)[Theorem](#)[Exercise](#)

32.Normal Spaces

[Theorem](#)[Exercise](#)

Theorem (30.3)

Suppose X has a cble basis. Then:

- ▶ *(a) Every open covering of X contains a cble subcoll covering X .*
- ▶ *(b) There exists a cble subset of X that is dense in X .*

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex30.2

Show that if X has a cble basis $\{B_n\}$, then every basis \mathcal{C} for X contains a cble basis for X .

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex30.3

Let X has a cble basis; let A be an ucble subset of X . Show that uncountably many points of A are limit points of A .

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex30.4

Show that every compact metrizable space X has a cble basis.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex30.5

- (a) Show that every metrizable space with a cble dense subset has a cble basis.
- (b) Show that every metrizable Lindelöf space has a cble basis.
- (Lindelöf space : every open covering contains a cble subcovering)

30.The Countability Axioms

Definition

Theorem

Exercise

31.The Separation Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex30.9

Let A be a closed subspace of X . Show that if X is Lindelöf, then A is Lindelöf.

Show by a example that if X has a cble dense subset, A need not have a cble dense subset.

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex30.14

Show that if X is Lindelöf and Y is compact, then $X \times Y$ is Lindelöf.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Definition

Definition

Suppose X is T_1 -space.

- ▶ Hausdorff
- ▶ Regular
- ▶ Normal

30.The Countability Axioms

Definition

Theorem

Exercise

31.The Separation Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Lemma (31.1)

Suppose X is T_1 -space.

- ▶ *(a) X is regular iff given pt x and nbd U of x , there is a nbd V of x s.t. $\bar{V} \subset U$.*
- ▶ *(b) X is normal iff given closed set A and open set U containing A , there is an open set V containing A s.t. $\bar{V} \subset U$*

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Theorem (31.2)

- ▶ (a) *subspace of Haus if Haus. product of Haus is Haus.*
- ▶ (b) *subspace of regular is regular. product of regular is regular.*

30.The Countability Axioms

Definition

Theorem

Exercise

31.The Separation Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex.31.1

Show that if X is regular, every pair of points of X have neighborhoods whose closures are disjoint.

30.The Countability Axioms

Definition

Theorem

Exercise

31.The Separation Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex.31.2

Show that if X is normal, every pair of disjoint closed sets have neighborhoods whose closures are disjoint.

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex.31.5

Let $f, g: X \rightarrow Y$ be conti; assume that Y is Haus. Show that $\{x \mid f(x) = g(x)\}$ is closed in X .

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex.31.6

Let $p : X \rightarrow Y$ be a closed conti surj map. Show that if X is normal, then so is Y .

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Theorem

Theorem (32.1)

Every regular space with a countable basis is normal.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Theorem

Theorem (32.2)

Every metrizable space is normal.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Theorem

Theorem (32.3)

Every compact Hausdorff is normal.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Theorem

Theorem (32.4)

Every well-ordered set X is normal in the order topology.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Exercise

Ex32.1

Show that a closed subspace of a normal space is normal.

30.The Countability Axioms

Definition

Theorem

Exercise

31.The Separation Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex32.2

Show that if $\prod X_\alpha$ is Haus (or regular, or normal), then so is X_α .

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex32.3

Show that every locally compact Hausdorff space is regular.

30. The
Countability
Axioms

Definition

Theorem

Exercise

31. The Separation
Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex32.4

Show that every regular Lindelöf space is normal.

30. The
Countability
Axioms

Definition

Theorem

Exercise

31. The Separation
Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex32.6

A space X is said to be completely normal if every subspace of X is normal. Show that X is completely normal iff for every pair A, B of separated sets in X (that is, $\bar{A} \cap B = \emptyset$ and $A \cap \bar{B} = \emptyset$), there exist disjoint open sets containing them.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex32.9(Challenging)

Theorem

If J is uncountable, then \mathbb{R}^J is not normal.

Proof) Let $X = (\mathbb{Z}_+)^J$. Since X is closed subspace of \mathbb{R}^J , it will suffice to show that X is not normal.

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex32.9

(a) If $\mathbf{x} \in X$ and if B is a finite subset of J , let $U(\mathbf{x}, B)$ denote the set consisting of all those elements \mathbf{y} of X such that $\mathbf{y}(\alpha) = \mathbf{x}(\alpha)$ for $\alpha \in B$. Show that the set $U(\mathbf{x}, B)$ are a basis for X .

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise

Ex32.9

(b) Define P_n to be the subset of X consisting of those \mathbf{x} such that on the set $J - \mathbf{x}^{-1}(n)$, the map \mathbf{x} is injective. Show that P_1 and P_2 are closed and disjoint.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex32.9

(c) Suppose U and V are open sets containing P_1 and P_2 , respectively. Given a sequence α_1, \dots of distinct elements of J , and a sequence $0 = n_0 < n_1 < \dots$ of integers, for $i \geq 1$ let us set $B_i = \{a_1, \dots, a_{n_i}\}$ and define $\mathbf{x}_i \in X$ by the equations $\mathbf{x}_i(\alpha_j) = j$ for $1 \leq j \leq n_{i-1}$, $\mathbf{x}_i(\alpha) = 1$ for all other values of α . Show that one can choose the sequences α_j and n_j so that for each i , one has the inclusion $U(\mathbf{x}_i, B_i) \subset U$.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

Ex32.9

(d) Let A be the set $\{\alpha_1, \dots\}$ constructed in (c). Define $\mathbf{y}: J \rightarrow \mathbb{Z}_+$ by the equations $\mathbf{y}(\alpha_j) = j$ for $\alpha_j \in A$, $\mathbf{y}(\alpha) = 2$ for all other values of α . Choose B so that $U(\mathbf{y}, B) \subset V$. Then choose i so that $B \cap A$ is contained in the set B_i . Show that $U(\mathbf{x}_{i+1}, B_{i+1}) \cap U(\mathbf{y}, B)$ is not empty.

30. The Countability Axioms

Definition

Theorem

Exercise

31. The Separation Axioms

Definition

Theorem

Exercise

32. Normal Spaces

Theorem

Exercise

The End

30.The
Countability
Axioms

Definition

Theorem

Exercise

31.The Separation
Axioms

Definition

Theorem

Exercise

32.Normal Spaces

Theorem

Exercise