EASI Report: (unnamed stream)

Scenario: Existing Conditions
Analysis point 43.6858, -72.2367 · Reach 1000.0 ft upstream
Scoring method: Alternative 2 with corrected cross-section quality checks (E5b) (method e91118cc821c), method package easi-screening v2

Complete screening coverage (includes proxy-derived ratings). Use the evidence profile and per-metric source tier to distinguish observations, models, and screening proxies.

Evidence profile: observed 0 · connected/nearby 4 · published model 2 · screening proxy 14 · manual 0 · unavailable 0

Shared evidence: BHR/ER incision geometry is reused by 4 metrics and should not be interpreted as independent evidence.

Basin characteristics

COMID	9327042
HUC12	010801040401
Watershed engine	STAF site engine v0.5.0
Drainage area	1.00 km²
Channel slope	0.0100 m/m (1.00%)
Stream order	2
Sinuosity	1.061
EPA ecoregion (Level III)	Northeastern Highlands (58)



Bars colored by condition band. Blue: Functioning · yellow: Functioning-at-Risk · red: Non-Functioning.

Watershed

Map background unavailable

Reach cross-sections



Bieger region: Appalachian Highlands · Bieger XS area: 0.8 m² · Bankfull width: 28.2 ft · Floodprone width: 40.7 ft · Section ER: 1.44 · Section BHR: 1.70 · Bankfull height: 0.43 ft · Low bank height: 0.73 ft
Metrics score on the reach medians of 9 sections: entrenchment ratio 1.54 (1.40 to 9.09), bank-height ratio 1.70 (0.72 to at least 2.00). The section at 300 ft, nearest both medians, is shown.

Function	Metric	Value	Rating	Idx	Fn	Conf
Catchment hydrology	Watershed Land-Cover Pressure	6.0% impervious cover (impervious governs)	Good	0.85	13	H
Reach inflow	Concentrated Runoff / Stormwater Inputs	1.71 km/km² road density	Fair	0.55	8	L
Streamflow regime	Flow Alteration (Regulation / Water Use)	degree of regulation 0.00% (storage 0 m³/km²; runoff 701 mm)	Good	0.85	13	M
Surface water storage	Percent Wetlands in Watershed	13.2% combined woody + herbaceous wetland cover	Good	0.85	13	H
Floodplain connectivity	Floodplain Access / Entrenchment	entrenchment ratio 1.54 (flood-prone width / bankfull width), reach median of 9 sections (1.40 to 9.09)	Good	0.85	13	M
High flow dynamics	Floodplain Engagement Frequency (bankfull recurrence)	bank-height ratio 1.70 (low-bank height / max bankfull depth), reach median of 9 sections (0.72 to at least 2.00)	Poor	0.20	3	M
Hyporheic connectivity	Hyporheic Exchange Indicators	channel slope 0.0100 m/m (slope pathway governs)	Good	0.85	13	L
Low flow and baseflow dynamics	Low-flow Wetted Connectivity†	EROM monthly flow variability (CV) 0.852	Good	0.85	13	L
Bed composition and large wood	Substrate Condition (grain size/embeddedness/fines/consolidation)	watershed agriculture 2.1%	Good	0.85	13	L

Function	Metric	Value	Rating	Idx	Fn	Conf
Channel and floodplain dynamics	Bank Erosion & Armoring Condition	bank-instability susceptibility from BHR 1.70, reach median of 9 sections (0.72 to at least 2.00)	Poor	0.20	3	L
Channel evolution	Channel Evolution Stage & Trends	channel-adjustment susceptibility (BHR 1.70, ER 1.54, reach medians of 9 sections), bhr governs	Poor	0.20	3	L
Sediment continuity	Sediment Supply Potential (watershed/banks)	soil erodibility governs (agriculture 2.1%, K 0.31, roads 1.71 km/km²)	Fair	0.55	8	M
Carbon processing	Detrital Processing (CPOM retention / shredders)	73.1% organic-matter supply potential (100 m corridor)	Fair	0.55	8	M
Light & thermal regime	Stream Temperature	thermal vulnerability: woody riparian 68.8%, impervious 6.0% (woodyRiparian governs)	Fair	0.55	8	L
Nutrient cycling	Nitrogen & Phosphorus Concentrations	TN 0.236 mg/L, TP 0.0091 mg/L (NAP region)	Good	0.85	13	L
Water & soil quality	Regulatory Impairment Status (305b/303d/TMDL)	nearest assessed unit 1 m away: IR category 4A (AU NHRIV801040401-03)	Fair	0.55	8	M/L
Community dynamics	Invasive / Non-native Species Presence	no established taxa recorded (HUC12)	Good	0.85	13	M
Habitat provision	In-stream Habitat Complexity & Cover	woody riparian cover 68.8% (sinuosity 1.06 shown as context)	Fair	0.55	8	L
Population support	Biological Integrity (IBI / community condition)†	predicted Good-BMMI probability 0.79	Good	0.85	13	M/L
Watershed connectivity	Fish Passage & Barrier Effects (longitudinal connectivity)	no mapped NID dams within 1 mile	Good	0.85	13	M

† Marked desktop evidence comes from the nearest StreamCat reach downstream.

Source details. This stream is outside the StreamCat network. Watershed metrics use the HR reach watershed (1.00 km²) from the STAF site engine. Scored from the nearest StreamCat reach, Mink Brook (COMID 9327042), 444 ft downstream, which drains 32 times this stream.

Generated from national datasets. A desktop screening estimate with per-metric confidence, not a field-validated assessment.