EPA's Concentration *versus* Time Database:

A Resource for Extrapolating Toxicokinetic Trends Across Chemicals

Christopher Cook^{1,2}, Lucas Albrecht¹, Grace Cary^{1,2}, Brenda Edwards¹, Derik Haggard¹, Nancy Hanley¹, Michael F. Hughes¹, Anna Jarnagin^{1,2}, Tirumula D. Kodavanti^{1,2}, Evgenia Koral-Bexell^{1,2}, Anna Kreutz^{1,2}, Mayla Ngo^{1,2}, Caroline Ring¹, Risa Sayre^{1,3}, Bhaskar Sharma¹, Jonathan Wall¹, Hiroshi Yamazaki⁴, John Wambaugh¹

Abstract No. 4656 Poster Board No. P702

¹Center for Computational Toxicology and Exposure, Office of Research and Development, United States Environmental Sciences and Engineering, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina; ⁴Showa Pharmaceutical University, Tokyo, Japan

1 Expansion of the Concentration versus Time Database

1a) What is the CvTdb?

• The Concentration *versus* Time Database (CvTdb) is a database of standardized time-course data of chemical compound concentrations measured *in vivo* from hundreds of diverse publications. It serves as a platform for assessing PK/TK models on a large set of data. (Sayre, Grulke, and Wambaugh, 2020).




Fig. 1. Plasma concentrations of DEHP (○) and MEHP (●) in rats after single oral administration of 2.8 g/kg DEHP (mean

Figure 1: Typical plot from which CvT data is extracted. From Sowa and Steibert (1985)


 The CvTdb contains 389 unique chemicals, including 70 pharmaceuticals (~30%). 101 analytes are present in EPA's ToxCast dataset (~53%).

1b) Integration of Data Shared by Other Institutions

- Showa Pharmaceutical University shared CvT data consisting of over 200 unique compounds (Kamiya *et al.*, 2020; Kamiya *et al.*, 2021).
- If YOU have CvT data, please consider collaborating with us! Visit https://github.com/USEPA/CompTox-PK-CvTdb or email Wambaugh.John@epa.gov

2 Toxicokinetic Trends Across Chemicals


- We can explore trends across chemicals and references using the
- "In PBPK modeling, predictions that are, on average, within a factor of 2 of the experimental data have frequently been considered adequate" (WHO, 2010).
- In Figure 2, 85.1% of replicate timepoint data in the CvTdb is within a factor of two of their respective mean. The "factor of two" rule may not be appropriate if the data itself is spread beyond a factor of two.




3 Introducing invivoPKfit

3a) What is *invivoPKfit*?

 R package that fits 1- and 2- compartment models to CvT data and extrapolates TK parameter values (e.g., half-life, elimination rate, and volume of distribution)



3b) TK Model Descriptions



ullet Parameters include V_d (volume of distribution), k_{elim} (elimination rate), k_{abs} (absorption rate; oral only), and F_{bio} (fraction bioavailable; oral




Figure 5: 2-Compartment Model. Adapted from Wambaugh et al. (2018)


$$C_{iv}(dose,\,t)\,=\,A\,st\,e^{-lpha\,st\,t}\,+\,B\,st\,e^{-eta\,st\,t} \ C_{oral}(dose,\,t)\,=\,A\,st\,e^{-lpha\,st\,t}\,+\,B\,st\,e^{-eta\,st\,t}\,+\,C\,st\,e^{-ka\,st\,t}$$

ullet Parameters include V1 (volume of primary compartment), k_{elim} , k_{12} (distribution rate to deep compartment), k_{21} distribution rate from deep compartment), k_{abs} (oral only), F_{bio} (oral only).

3c) Statistical Model Description

ullet invivoPKfit calculates the probability of C_{obs} given a PK model $M(p,\ t)$ and a statistical model $S(\sigma)$, where $P=L(C_{obs},\ M,\ S)$, and assumes that C_{obs} are log-normally distributed around the concentrations predicted by M(p, t) (Cox and Hinkley, 1979).


$$ln~L(x,~\mu,~\sigma)~=~\Sigma_{i=1,n}lnrac{1}{x\sigma\sqrt{2\pi}}e^{-rac{1}{2}\left(rac{ln~x-\mu)}{\sigma}
ight)^2} \ ln~L(x,~\mu,~\sigma)~=~\Sigma_{i=1,n}lnrac{1}{C_{obs,i}\sigma\sqrt{2\pi}}e^{-rac{1}{2}\left(rac{ln~C_{obs,i}-M(p,t_i)}{\sigma}
ight)^2}$$



4 invivoPKfit Analyses

4a) Predicted Concentrations


- 63.3% of 16,675 concentrations predicted by the 1-compartment model fit within a factor of two of the observed concentrations.
- 69.1% of 13,097 concentrations predicted by the 2-compartment model fit within a factor of two of the observed concentrations.



observed concentrations, such that $C_{\text{pred}} / C_{\text{obs}}$. The red dashed lines mark the 'factor of two' bounds. This plot does not include


4b) invivoPKfit Performance versus Data Variability

• Can start to gauge how data variability influences the performance of a



concentrations normalized by experiment-timpoint specific replicates (see Figure 2). The dashed lines mark the 'factor of 2' bounds to each axis. Sections were assigned based on both the ability of a model to accurately predict an observed concentration and the spread of the observed data. Points colored pink represent concentrations significantly underpredicted by the model.

• Plotting model predictions against observed data can help determine why a model might fit poorly. A poor fit could indicate 1) a deficiency in the model, 2) an inaccurate curation of data, and/or 3) a deficiency in the experimental collection of data.



represent non-detects that were below the limit of quantitation (L.O.Q.).

References

and Biopharmaceutics 2 (2): 123-48. https://doi.org/10.1007/BF01061504

Cox, D. R., and D. V. Hinklev. 1979. *Theoretical Statistics*. CRC Press

Estimation of Statistical Properties of Parameters." Applied Mathematics Letters 4 (1): 87–90. https://doi.org/https://doi.org/10.1016/089

Based Pharmacokinetic Models to Estimate Plasma and Tissue Concentrations After Oral Doses in Rats." Chemical Research in Toxicology 34 (10)

Pharmacokinetic Models Predicting Renal and Hepatic Concentrations of Industrial Chemicals After Virtual Oral Doses in Rats." Chemical Research in Toxicology 33 (7): 1736-51. https://doi.org/10.1021/acs.chemrestox.0c00009

Kudo. Naomi. Masanori Katakura. Yasunori Sato. and Yoichi Kawashima. 2002. "Sex Hormone-Regulated Renal Transport of Perfluorooctanoic Acid." Chemico-Biological Interactions 139 (3): 301-16. https://doi.org/10.1016/s0009-2797(02)00006-6.

Nash, John C. 2014. "On Best Practice Optimization Methods in R." Journal of Statistical Software 60 (2): 1-14. https://doi.org/10.18637/jss.v060.i02.

O'Flaherty, E. J. 1981. *Toxicants and Drugs: Kinetics and Dynamics*. John Wiley & Sons.

43 (9): 1-14. https://doi.org/10.18637/jss.v043.i09

Sayre, Risa R., John F. Wambaugh, and Christopher M. Grulke. 2020. "Database of Pharmacokinetic Time-Series Data and Parameters for 144 Environmental Chemicals." *Scientific Data* 7 (1): 122. https://doi.org/10.1038/s41597-020-0455-1.

Sheiner, L. B. 1985. "Analysis of Pharmacokinetic Data Using Parametric Models. II. Point Estimates of an Individual's Parameters." *Journal of* Pharmacokinetics and Biopharmaceutics 13 (5): 515-40. https://doi.org/10.1007/BF01059333.


Thorne, W. Brent. 2019. Posterdown: An R Package Built to Generate Reproducible Conference Posters for the Academic and Professional World

"Evaluating in Vitro-in Vivo Extrapolation of Toxicokinetics." Toxicological Sciences: An Official Journal of the Society of Toxicology 163 (1): 152–69.

World Health Organization. 2010. "Characterization and Application of Physiologically Based Pharmacokinetic Models in Risk Assessment." WHO Document Production Services, Geneva, Switzerland.

Acknowledgements

Employees of the U.S. EPA were funded through EPA's Office of Research and Development. This project was supported in part by an appointment to the Research Participation Program at the Center for Computational Toxicology, U.S. Environmental Protection Agency, administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and EPA. *This work does not*



necessarily reflect U.S. EPA policy.		