



RESEARCH ARTICLE

Associations between chlorophyll *a* and various microcystin-LR health advisory concentrations

Jeff Hollister, Betty Kreakie

Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, US Environmental Protection Agency, Narragansett, RI, USA

v1 First published: N/A, N/A: N/A (doi: N/A)
Latest published: N/A, N/A: N/A (doi: N/A)

Abstract

Cyanobacteria harmful algal blooms (cHABs) are associated with a wide range of adverse health effects that stem mostly from the presence of cyanotoxins. To help protect against these impacts, several health advisory levels have been set for some toxins. In particular, one of the more common toxins, microcystin-LR, has several advisory levels set for drinking water and recreational use. However, compared to other water quality measures, field measurements of microcystin-LR are not commonly available due to cost and advanced understanding required to interpret results. Addressing these issues will take time and resources. Thus, there is utility in finding indicators of microcystin-LR that are already widely available, can be estimated quickly and *in situ*, and used as a first defense against high concentrations of microcystin-LR. Chlorophyll *a* is commonly measured, can be estimated *in situ*, and has been shown to be positively associated with microcystin-LR. In this paper, we use this association to provide estimates of chlorophyll *a* concentrations that are indicative of a higher probability of exceeding select health advisory concentrations for microcystin-LR. Using the 2007 National Lakes Assessment and a conditional probability approach, we identify chlorophyll *a* concentrations that are more likely than not to be associated with an exceedance of a microcystin-LR health advisory level. We look at the recent US EPA health advisories for drinking water as well as the World Health Organization levels for drinking water and recreational use and identify a range of chlorophyll *a* thresholds. A 50% chance of exceeding one of the specific advisory microcystin-LR concentrations of 0.3, 1, 1.6, and 2 g/L is associated with chlorophyll *a* concentration thresholds of 24.64, 65.6, 89.71, and 94.93, respectively. When managing for these various microcystin-LR levels, exceeding these reported chlorophyll *a* concentrations should be a trigger for further testing and possible management action.

Open Peer Review

Referee Status: AWAITING PEER

REVIEW

Discuss this article

Comments (0)

Corresponding author: Jeff Hollister (hollister.jeff@epa.gov)**How to cite this article:** Hollister J and Kreakie B. Associations between chlorophyll *a* and various microcystin-LR health advisory concentrations F1000Research, : (doi:)**Copyright:** © 2016 Hollister J and Kreakie B. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**Grant information:** The author(s) declared that no grants were involved in supporting this work.**Competing interests:** No competing interests were disclosed.**First published:** N/A, N/A: N/A (doi: N/A)

Introduction

Over the last decade, numerous events and legislative activities have raised the public awareness of harmful algal blooms^{1–3}. In response the US Environmental Protection Agency (US EPA) has recently released suggested microcystin-LR (one of the more common toxins) concentrations that would trigger health advisories^{4–6}. Additionally, the World Health Organization (WHO) has proposed microcystin advisory levels for drinking water and a range of recreational risk levels^{7,8}. While these levels and associated advisories are likely to help mitigate the impacts from harmful algal blooms, they are not without complications.

One of these complications is that they rely on available measurements of microcystin-LR. While laboratory testing (e.g., chromatography) remains the gold standard for quantifying microcystin-LR concentrations in water samples, several field test kits have been developed. Even though field tests provide a much needed means for rapid assessment, they are not yet widely used and are moderately expensive (approximately \$150–\$200 depending on the specific kit) with a limited shelf life (typically one year)^{9,10}. Additionally, each technique requires nuanced understanding of the detection method (e.g., limit of detection, specific microcystin variants being measured, and sampling protocol).

Fortunately, microcystin-LR has been shown to be associated with several **other**, more commonly measured and well understood components of water quality that are readily assessed in the field. For instance, there are small or hand held fluorometers that measure chlorophyll *a*. Additionally, chlorophyll *a* is a very commonly measured component of water quality that is also known to be positively associated with microcystin-LR concentrations^{11,12}. **Yuan et al.**¹³ explore these associations in detail and control for other related variables. In their analysis they find that total nitrogen and chlorophyll *a* show the strongest association with microcystin. Furthermore, they identify chlorophyll *a* and total nitrogen concentrations that are associated with exceeding 1 µg/L of microcystin. Given these facts, it should be possible to identify chlorophyll *a* concentrations that would be associated with the new US EPA microcystin-LR health advisory levels for drinking water. Identifying these associations would provide another tool for water resource managers to help manage the threat to public health posed by cHABs and would be especially useful in the absence of measured microcystin-LR concentrations.

In this paper we build on past efforts and utilize the National Lakes Assessment (NLA) data and identify chlorophyll *a* concentrations that are associated with higher probabilities of exceeding several microcystin-LR health advisory concentrations^{6,8,13}. We add to past studies by exploring associations with newly announced advisory levels and by also applying a different method, conditional probability analysis. Utilizing different methods strengthens the evidence for suggested chlorophyll *a* levels that are associated with increased risk of exceeding the health advisory levels as those levels are not predicated on a single analytical method. So that others may repeat or adjust this analysis, the data, code, and this manuscript are freely available via <https://github.com/USEPA/microcystinchla>.

Methods

Data

We used the 2007 NLA chlorophyll *a* and microcystin-LR concentration data¹³. These data represent a snapshot of water quality from the summer of 2007 for the conterminous United States and were collected as part of an ongoing probabilistic monitoring program¹³. Data on chlorophyll *a* and microcystin-LR concentrations are available for 1028 lakes. These data are available for download from the [National Lake Assessment Data Site](#).

Analytical methods

We used a conditional probability analysis (CPA) approach to explore associations between chlorophyll *a* concentrations and World Health Organization (WHO) and US EPA microcystin-LR health advisory levels¹⁴. Many health advisory levels have been suggested (**Table 1**), but lakes with higher microcystin-LR concentrations in the NLA were rare. Only 1.16% of lakes sampled had a concentration greater than 10 µg/L. Thus, for this analysis we focused on the microcystin concentrations that are better represented in the NLA data. These were the US EPA children's drinking water advisory level of 0.3 µg/L (US EPA child), the WHO drinking water advisory level of 1 µg/L (WHO drinking), the US EPA adult drinking water advisory level of 1.6 µg/L (US EPA adult), and the WHO recreational, low probability of effect advisory level of 2 µg/L (WHO recreational)^{6–8}. **Conditional probability analysis provides** information about the probability of observing one event given another event has also occurred. For this analysis, we used CPA to examine how the conditional probability of exceeding one of the health advisories changes as chlorophyll *a* increases in a lake. We expect to find higher chlorophyll *a* concentrations to be associated

Table 1. Various suggested microcystin-LR health advisory concentrations from the US EPA and World Health Organization.

Source	Type	Concentration
US EPA	Adult drinking water advisory	1.6 µg/L
US EPA	Child drinking water advisory	0.3 µg/L
WHO	Drinking water	1 µg/L
WHO	Recreational: High probability of effect	20–2000 µg/L
WHO	Recreational: Low probability of effect	2–4 µg/L
WHO	Recreational: Moderate probability of effect	10–20 µg/L
WHO	Recreational: Very high probability of effect	>2000 µg/L

with higher probabilities of exceeding the microcystin-LR health advisory levels. We also calculated bootstrapped 95% confidence intervals (CI) using 1000 bootstrapped samples. Thus, to identify chlorophyll *a* concentrations of concern we identify the value of the upper 95% CI across a range of conditional probabilities of exceeding each health advisory level. Using the upper confidence limit to identify a threshold is justified as it ensures that a given threshold is unlikely to miss a microcystin exceedance.

As both microcystin-LR and chlorophyll *a* values were highly skewed right, a log base 10 transformation was used. Additional details of the specific implementation are available at <https://github.com/USEPA/microcystinchla>. A more detailed discussion of CPA is beyond the scope of this paper, but see Paul *et al.*¹⁵ and Hollister *et al.*¹⁶ for greater detail. Lastly, all analyses were conducted using R version 3.2.2 and code and data from this analysis are freely available as an R package at <https://github.com/USEPA/microcystinchla>.

Lastly, we assess the ability of these chlorophyll *a* thresholds to predict microcystin exceedance. We use error matrices and calculate total accuracy as well as the proportion of false negatives. Total accuracy is the total number of correct predictions divided by total observations. The proportion of false negatives is the total number of lakes that were predicted to not exceed the microcystin guidelines but actually did, divided by the total number of observations.

Results

In the 2007 NLA, microcystin-LR concentrations ranged from 0.05 to 225 $\mu\text{g/L}$. Microcystin-LR concentrations of 0.05 $\mu\text{g/L}$ represent the detection limits. Any value greater than that indicates the presence of microcystin-LR. Of those lakes with microcystin, the median concentration was 0.51 $\mu\text{g/L}$ and the mean was 3.17 $\mu\text{g/L}$. Of all lakes sampled, 21% of lakes exceeded the US EPA child level, 8.8% of lakes exceeded the US EPA adult level, 11.7% of lakes exceeded the WHO drinking level, and 7.3% of lakes exceeded the WHO recreational level. For chlorophyll *a*, the range was 0.07 to 936 $\mu\text{g/L}$. All lakes had reported chlorophyll *a* concentrations that exceeded detection limits. The median concentration was 7.79 $\mu\text{g/L}$ and the mean was 29.63 $\mu\text{g/L}$. The associations between chlorophyll *a* and the upper confidence interval across a range of conditional probability values are shown in Table 2. Specific chlorophyll *a* concentrations that are associated with greater than even odds of exceeding the advisory levels were 0.07, 0.07, 2.98, and 10.67 $\mu\text{g/L}$ for 0.3, 1.0, 1.6, and 2.0 $\mu\text{g/L}$ advisory levels, respectively (Table 2 & Figure 1).

The chlorophyll *a* cutoffs may be used to predict whether or not a lake exceeds the microcystin-LR health advisories. Doing so allows us to compare the accuracy of the prediction as well as evaluate false negatives. Total accuracy of the four cutoffs predicting microcystin-LR exceedances were 19% for the US EPA children's advisory, 21% for the WHO drinking water advisory, 22% for the US EPA adult advisory, and 23% for the WHO recreational advisory (Table 3–Table 6). However, total accuracy is only one part of the prediction performance with which we are concerned.

When using the chlorophyll *a* cutoffs as an indicator of microcystin-LR exceedances, the error that should be avoided is predicting that no exceedance has occurred when in fact it has. In other words, we would like to avoid Type II errors and minimize the proportion of

Table 2. Chlorophyll *a* concentrations that are associated with a 50 microcystin-LR health advisory concentration.

Conditional probability	US EPA child	WHO drink	US EPA adult	WHO recreational
0.10	0.07	0.07	0.07	1.22
0.20	0.07	4.43	11.80	20.00
0.30	2.98	20.43	31.40	53.64
0.40	10.67	42.24	66.96	82.22
0.50	23.36	66.96	83.52	105.84
0.60	38.30	103.20	133.20	155.52
0.70	65.60	166.63	871.20	216.00
0.80	117.50	338.40	871.20	871.20
0.90	167.04	516.00	871.20	871.20

false negatives. For the four chlorophyll *a* cut-offs we had a proportion of false negatives of 2%, 2%, 1%, and 1% for the US EPA children's, the WHO drinking water, the US EPA adult, and the WHO recreational advisories, respectively. In each case we missed less than 10% of the lakes that in fact exceeded the microcystin-LR advisory. While this method performs well with regard to the false negative percentage, it is possible that is a relic of the NLA dataset and testing with additional data would allow us to confirm this result.

Discussion

The association between Log10 microcystin-LR and Log10 chlorophyll *a* shows a wedge pattern (Figure 2). This indicates that, in general, higher concentrations of microcystin-LR almost always co-occur with higher concentrations of chlorophyll *a* yet the inverse is not true. Higher chlorophyll *a* is not necessarily predictive of higher microcystin-LR concentrations; however, chlorophyll *a* may be predictive of the probability of exceeding a certain threshold.

This is the case as the probability of exceeding each of the four tested health advisory levels increases as a function of chlorophyll *a* concentration (Figure 1). We used this association to identify chlorophyll *a* concentrations that are associated with a range of probabilities of exceeding a given health advisory level (Table 2). For the purposes of this discussion we focus on a conditional probability of 50% or greater (i.e., greater than even odds to exceed a health advisory level). The 50% conditional probability chlorophyll *a* thresholds represent 27.9%, 11.3%, 9%, and 6.9% of sample lakes for the US EPA child, the WHO drinking, the US EPA adult, and the WHO recreational levels, respectively.

There are numerous possible uses for the chlorophyll *a* and microcystin-LR advisory cut-off values. First, in the absence of microcystin-LR measurements, exceedance of the chlorophyll *a* concentrations could be a trigger for further actions. Given that there is uncertainty around these chlorophyll *a* cutoffs the best case scenario would be to monitor for chlorophyll *a* and in the event of exceeding a target concentration take water samples and have those samples tested for microcystin-LR.




Figure 1. Conditional probability plots showing association between the probability of exceeding various microcystin-LR (MLR) health advisory levels. **A.** Plot for US EPA child ($0.3 \mu\text{g/L}$). **B.** Plot for WHO drinking ($1 \mu\text{g/L}$). **C.** Plot for US EPA adult ($1.6 \mu\text{g/L}$). **D.** Plot for WHO recreational ($2 \mu\text{g/L}$).



Table 3. Confusion matrix comparing chlorophyll a predicted exceedances (rows) versus real exceedances (columns) for the US EPA children's drinking water advisory.

	Not exceed	Exceed	Row totals
Not exceed	649	96	745
Exceed	162	121	283
Column totals	811	217	1028

Table 4. Confusion matrix comparing chlorophyll a predicted exceedances (rows) versus real exceedances (columns) for the WHO drinking water advisory.

	Not exceed	Exceed	Row totals
Not exceed	839	77	916
Exceed	68	44	112
Column totals	907	121	1028

Table 5. Confusion matrix comparing chlorophyll a predicted exceedances (rows) versus real exceedances (columns) for the US EPA adult drinking water advisory.

	Not exceed	Exceed	Row totals
Not exceed	883	57	940
Exceed	54	34	88
Column totals	937	91	1028

Table 6. Confusion matrix comparing chlorophyll a predicted exceedances (rows) versus real exceedances (columns) for the WHO recreational water advisory.

	Not exceed	Exceed	Row totals
Not exceed	910	52	962
Exceed	43	23	66
Column totals	953	75	1028




Figure 2. Scatterplot showing association between chlorophyll *a* and microcystin-LR.

A second potential use is to identify past bloom events from historical data. As harmful algal blooms are made up of many species and have various mechanisms responsible for adverse impacts (e.g., toxins, hypoxia, odors), there is no single definition of a bloom. For cHABs, one approach has been to identify an increase over a baseline concentration of phycocyanin¹⁷. This is a useful approach for targeted studies, but phycocyanin is also not always available and measures the predominance of cyanobacterial pigments and not toxins. Using our chlorophyll *a* cutoffs provides a value that is more directly associated with microcystin-LR and can be used to classify lakes, from past surveys, as having bloomed.

Lastly, using chlorophyll *a* is not meant as a replacement for testing of microcystin-LR or other toxins. It should be used when other, direct measurements of cyanotoxins are not available. In those cases, which are likely to be common at least in the near future, using a more ubiquitous measurement, such as chlorophyll *a* will provide a reasonable proxy for the probability of exceeding a microcystin-LR health advisory level and provide better protection against adverse effects in both drinking and recreational use cases.

Data and software availability

Data and latest source code

<https://github.com/USEPA/microcystinchla>

Archived data and source code at time of publication

<http://dx.doi.org/10.5281/zenodo.45317>¹⁸

License

Creative Commons Zero 1.0: <http://creativecommons.org/public-domain/zero/1.0/>

Author contributions

JH and BK conceived of and conducted the analysis. Both authors reviewed the results and contributed to writing the manuscript.

Competing interests

No competing interests were disclosed.

Grant information

The author(s) declared that no grants were involved in supporting this work.

Acknowledgements

We would like to thank Anne Kuhn, Bryan Milstead, John Kiddon, Joe LiVolsi, Tim Gleason, and Wayne Munns for constructive reviews of this paper. This paper has not been subjected to Agency review. Therefore, it does not necessarily reflect the views of the Agency. Mention of trade names or commercial products does not constitute endorsement or recommendation for use. This contribution is identified by the tracking number ORD-015143 of the Atlantic Ecology Division, Office of Research and Development, National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency.

References

1. Jetoo S, Grover VI, Krantzberg G: **The Toledo drinking water advisory: Suggested application of the water safety planning approach.** *Sustainability*. 2015; 7(8): 9787–9808.
[Publisher Full Text](#)
2. Rinta-Kanto JM, Konopko EA, DeBruyn JM, *et al.*: **Lake Erie *Microcystis*: Relationship between microcystin production, dynamics of genotypes and environmental parameters in a large lake.** *Harmful Algae*. 2009; 8(5): 665–673.
[Publisher Full Text](#)
3. Harmful Algal Bloom and Hypoxia Research and Control Amendments Act: **Harmful Algal Bloom and Hypoxia Research and Control Amendments Act of 2014.** 2014; S. 1254.
[Reference Source](#)
4. McElhiney J, Lawton LA: **Detection of the cyanobacterial hepatotoxins microcystins.** *Toxicol Appl Pharmacol*. 2005; 203(3): 219–230.
[PubMed Abstract](#) | [Publisher Full Text](#)
5. Zurawell RW, Chen H, Burke JM, *et al.*: **Hepatotoxic cyanobacteria: a review of the biological importance of microcystins in freshwater environments.** *J Toxicol Environ Health B Crit Rev*. 2005; 8(1): 1–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. US EPA: **Drinking water health advisory for the cyanobacterial microcystin toxins.** EPA-820-R-15100. 2015.
[Reference Source](#)
7. World Health Organization: **Cyanobacterial toxins: Microcystin-LR in drinking-water.** Background document for development of WHO *guidelines for drinking-water quality*. Geneva, Switzerland. World Health Organization, 2nd ed. Geneva. 2003.
[Reference Source](#)
8. Chorus I, Bartram J: **Toxic cyanobacteria in water: A guide to their public health consequences, monitoring and management.** World Health Organization. 1999.
[Reference Source](#)
9. James R, Gregg A, Dindal A, *et al.*: **Environmental technology verification report: Abraxis microcystin test kits.** Online document. Accessed online: June 22, 2011.
[Reference Source](#)
10. Aranda-Rodríguez R, Jin Z, Harvie J, *et al.*: **Evaluation of three field test kits to detect microcystins from a public health perspective.** *Harmful Algae*. 2015; 42: 34–42.
[Publisher Full Text](#)
11. Pip E, Bowman L: **Microcystin and algal chlorophyll in relation to nearshore nutrient concentrations in Lake Winnipeg, Canada.** *Environ Pollut*. 2014; 3(2): p36.
[Publisher Full Text](#)
12. Yuan LL, Pollard AI, Pather S, *et al.*: **Managing microcystin: Identifying national-scale thresholds for total nitrogen and chlorophyll a.** *Freshwater Biol*. 2014; 59(9): 1970–1981.
[Publisher Full Text](#)
13. US EPA: **National lakes assessment: A collaborative survey of the nation's lakes.** EPA 841-R-09-001. 2009.
[Reference Source](#)
14. Paul JF, Munns WR Jr: **Probability surveys, conditional probability, and ecological risk assessment.** *Environ Toxicol Chem*. 2011; 30(6): 1488–1495.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. Paul JF, McDonald ME: **Development of empirical, geographically specific water quality criteria: A conditional probability analysis approach.** *J Am Water Resour Assoc*. 2005; 41(5): 1211–1223.
[Publisher Full Text](#)
16. Hollister JW, Walker HA, Paul JF: **CProb: a computational tool for conducting conditional probability analysis.** *J Environ Qual*. 2008; 37(6): 2392–2396.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Miller TR, Beversdorf L, Chaston SD, *et al.*: **Spatiotemporal molecular analysis of cyanobacteria blooms reveals *Microcystis*–*Aphanizomenon* interactions.** *PLoS One*. 2013; 8(9): e74933.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Hollister JW, Kreakie BJ: **Associations between chlorophyll a and various microcystin-LR.** *Zenodo*. 2016.
[Data Source](#)