

Associations between Chlorophyll *a* and various Microcystin-LR Health Advisory Concentrations

Jeffrey W. Hollister^{*} ¹ Betty J. Kreakie¹

¹US Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, 27 Tarzwell Drive Narragansett, RI, 02882, USA

^{*} corresponding author: hollister.jeff@epa.gov

Cyanobacteria harmful algal blooms (cHABs) are associated with a wide range of adverse health effects that stem mostly from the presence of cyanotoxins. To help protect against these impacts, several health advisory levels have been set for some toxins. In particular, one of the more common toxins, microcystin, has several advisory levels set for drinking water and recreational use and managing water bodies to meet those levels could have far reaching benefits. However, compared to other water quality measures, measurements of microcystin are not common and current field measurement techniques have limited precision and accuracy. Addressing these issues will take time and resources. Thus, there is utility in finding indicators of microcystin that are already widely available, can be estimated quickly and *in situ*, and used as a first defense against high levels of microcystin. In particular, chlorophyll *a* is very commonly measured, can be estimated *in situ*, and has been shown to be positively associated with microcystin. In this paper we use this association to provide estimates of chlorophyll *a* that if exceeded would be indicative of a higher probability of exceeding select health advisory concentrations for microcystin-LR. Using the 2007 National Lakes Assessment and a conditional probability approach that has been used in other water quality settings, we identify chlorophyll *a* concentrations that are more likely than not to be associated with an exceedance of a microcystin health advisory level. We look at the recent US EPA standards for drinking water as well as the World Health Organization levels for drinking water and recreational use. For microcystin concentrations of 0.3, 1, 1.6, and 2 we find chlorophyll *a* concentrations of 23.68, 63.94, 89.71, and 102.6, respectively. When managing for these various microcystin levels exceeding these reported chlorophyll *a* concentrations should be a trigger for further testing and possible management action.

1 Introduction

In the summer of 2014, the city of Toledo, OH was forced to shut down their municipal water supply due in part to an excess of microcystin-LR that resulted from a ongoing cyanobacteria harmful algal bloom (cHAB) in Lake Erie [REFS]. Since this event, significant legislation has been passed in the United States and the US Environmental Protection Agency (USEPA) has released suggested microcystin-LR concentrations that would trigger health advisories. MORE ON THE LEVELS. While these levels and associated advisories are likely to help mitigate the impacts from harmful algal blooms, they are not

without complications.

One of these complications is that they rely on available measurements of microcystin-LR. This toxin can be measured in the field using test strips but these are a coarse measure at best and currently available test strips focus on 1 and 10 $\mu\text{g/L}$ [REFS]. Measurements with greater accuracy and precision require taking water samples and processing those in a lab to determine the toxin concentration [REFS]. Additionally, microcystin-LR is, currently, not a routinely collected water quality parameter, thus, availability of microcystin-LR data may limit our ability to screen water bodies for exceedances of the various health advisories concentrations. Until microcystin-LR concentrations are more widely collected an alternative measure is needed. Fortunately, microcystin-LR has been shown to be associated with several other, more easily measured components of water quality.

Chlorophyll *a* is a very commonly measured components of water quality that is also known to be associated with Microcystin-LR concentrations [REFS]. Additionally there are many rapid measurements for assessing chlorophyll *a* levels *in situ*. For instance, there are small or hand held flourometers that provide reliable measurements [REFS]. Given these facts, it might be possible to identify chlorophyll *a* concentrations that would be associated with the various Microcystin-LR health advisory levels. Identifying these associations would provide another reliable tool for water resource managers to use to help manage the threat to public health posed by cHABs and would be especially useful in the absence of microcystin-LR concentrations. Thus, the goal of this paper is to utilize the National Lakes Assessment data and identify chlorophyll *a* concentrations that are associated with higher probabilities of exceeding several microcystin-LR health advisory concentrations [NLA REF]. So that others may repeat or adjust this analysis, the data, code, and this manuscript are freely available via <https://github.com/USAPE/microcystinchla>.

2 Methods

2.1 Data

We used the 2007 National Lakes Assessment (NLA) water quality and microcystin-LR concentration data [REF]. These data represent a snapshot of water quality from the summer of 2007 and data on chlorophyll *a* and microcystin-LR concentrations are available for lakes.

2.2 Conditional Probability Analysis

We used a conditional probability analysis (CPA) approach to explore associations between chlorophyll *a* concentrations and World Health Organization (WHO) and U.S. Environmental Protection Agency (U.S. EPA) microcystin-LR health advisory levels (Paul and Munns 2011). Many levels have been suggested (Table 1), but lakes with higher microcystin-LR concentrations in the NLA were rare. Only 1.16 % of lakes sampled had a concentration greater than 10. Thus, for this analysis we focus on the microcystin concentrations that are better represented in the NLA data. These were 0.3, 1, 1.6, and 2 $\mu\text{g/L}$.

A detailed discussion of CPA is beyond the scope of this paper, but see Paul et al. [REF] and Hollister et al. [REF] for details. For this analysis, we used CPA to examine how the conditional probability of exceeding one of the health advisory changes as chlorophyll *a* increases in a lake. The 95% confidence intervals were calculated from 1000 bootstrapped samples. To identify chlorophyll *a* concentrations of concern we used a 50% conditional probability of exceeding each health advisory level and extracted the minimum chlorophyll *a* concentration that was associated with the upper confidence level being 50% or greater. As both microcystin-LR and chlorophyll *a* values were both highly skewed right, a log base 10 transformation was used. Additional details of the specific implementation are available at <https://github.com/USEPA/microcystinchla>.

3 Results

In the 2007 NLA, microcystin-LR concentrations ranged from 0.05 to 225. Microcystin-LR concentrations of 0.05 $\mu\text{g/L}$ represent the detection limits. Any value greater than that indicates the presence of microcystin-LR. Of those lakes with microcystin, the median concentration was 0.51 and the mean was 0.51. Lastly, of all lakes sampled, 21% of lakes exceeded the U.S. EPA childrens drinking water standard, 8.8% of lakes exceeded the U.S. EPA adult drinking water standard, 11.7% of lakes exceeded the WHO drinking water standard, and 7.3% of lakes exceeded the WHO recreational standard. For chlorophyll *a*, the range was 0.07 to 936. All lakes had reported chlorophyll *a* concentrations that exceeded detection limits. The median concentration was 7.79 and the mean was 29.6301946. The associations between chlorophyll *a* and the upper confidence interval with a conditional probability of 50% were 23.68, 63.94, 89.71, and 102.6 for 0.3, 1.0, 1.6 and 2.0 $\mu\text{g/L}$, respectively (Table 2 & Figure 2).

4 Discussion

The association between Log 10 microcystin-LR and Log 10 chlorophyll *a* show a wedge pattern (Figure 1). This indicates that higher concentrations of microcystin-LR almost always co-occur with higher concentrations of chlorophyll *a* yet the inverse is not true. Higher chlorophyll *a* is not necessarily predictive of higher microcystin-LR concentrations; however, chlorophyll *a* may be predictive of the probability of exceeding a certain concentration.

This is the case as the probability of exceeding each of the four tested health advisory levels increases as a function of chlorophyll *a* concentration (Figure 2). We use this association to identify chlorophyll *a* concentrations that are associated with greater than even odds of exceeding a given health advisory level (Table 2). These represent 27.8%, 11.9%, 8.3%, and 7.4% of sample lakes for the U.S EPA childrens drinking water, the WHO drinking water, the U.S. EPA Adult drinking water, and the WHO recreational standards, respectively.

There are numerous possible uses for the chlorophyll *a* and microcystin-LR advisory cut-off values. First, in the absence of microcystin-LR measurements, exceedence of the chlorophyll *a* concentrations could be

a trigger for further actions. Given that there is uncertainty around these chlorophyll *a* cutoffs the best case scenario would be to monitor for chlorophyll and in the event of exceeding a target concentration take water samples and have those samples tested in a lab for microcystin-LR. A second potential use is to identify possible bloom events from historical data. Currently, there are several suggestions on how to identify a bloom event, but there is variation across the definitions and quantitative links to measures of cyanobacteria are not always included [REFS]. Using our chlorophyll *a* cutoffs provides a value that is directly associated with microcystin-LR and can be used to classify lakes, from past surveys, as havin bloomed.

These values are conservative: greater than even odds for the upper CI. Most protective.

Not meant as a replacement for testing of microcystin, but provides another useful tool to help understand the probable extent of microcystin occurence.

5 Figures




Figure 1: Scatterplot showing association between chlorophyll *a* and microcystin-LR.

```
## Loading required package: grid
```




Figure 2: Conditional probability plots showing association between the probability of exceeding various microcystin-LR (MLR) health advisory Levels. A.) Plot for U.S. EPA childrens drinking water advisory (0.3 $\mu\text{g/L}$). B.) Plot for WHO drinking water advisory (1 $\mu\text{g/L}$). C.) Plot for U.S. EPA adult drinking water advisory (1.6 $\mu\text{g/L}$). D.) Plot for WHO recreational advisory (2 $\mu\text{g/L}$).

6 Tables

Table 1: Various suggested microcystin-LR health advisory concentrations.

Source	Type	Concentration
WHO	Drinking	1 $\mu\text{g/L}$
U.S. EPA	Drinking	0.3 $\mu\text{g/L}$
U.S. EPA	Drinking	1.6 $\mu\text{g/L}$
WHO	Recreational	2-4 $\mu\text{g/L}$
WHO	Recreational	10-20 $\mu\text{g/L}$
WHO	Recreational	20-2000 $\mu\text{g/L}$
WHO	Recreational	>2000 $\mu\text{g/L}$

Table 2: Chlorophyll *a* concentrations that are associated with a 50% probability of exceeding a microcystin-LR health advisory concentration.

Source	Type	Microcystin	Chlorophyll
U.S. EPA	Drinking	0.3	23.68
WHO	Drinking	1.0	63.94
U.S. EPA	Drinking	1.6	89.71
WHO	Recreational	2.0	102.60

References

Paul, J. F., and W. R. Munns. 2011. Probability surveys, conditional probability, and ecological risk assessment. *Environmental Toxicology and Chemistry* 30:1488–1495.