

# Associations between chlorophyll *a* and various microcystin health advisory Concentrations

Jeffrey W. Hollister<sup>\*</sup> <sup>1</sup> Betty J. Kreakie<sup>1</sup>

<sup>1</sup>US Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, 27 Tarzwell Drive Narragansett, RI, 02882, USA

<sup>\*</sup> corresponding author: [hollister.jeff@epa.gov](mailto:hollister.jeff@epa.gov)

Cyanobacteria harmful algal blooms (cHABs) are associated with a wide range of adverse health effects that stem mostly from the presence of cyanotoxins. To help protect against these impacts, several health advisory levels have been set for some toxins. In particular, one of the more common toxins, microcystin, has several advisory levels set for drinking water and recreational use. However, compared to other water quality measures, field measurements of microcystin are not commonly available due to cost and advanced understanding required to interpret results. Addressing these issues will take time and resources. Thus, there is utility in finding indicators of microcystin that are already widely available, can be estimated quickly and *in situ*, and used as a first defense against high levels of microcystin. Chlorophyll *a* is commonly measured, can be estimated *in situ*, and has been shown to be positively associated with microcystin. In this paper, we use this association to provide estimates of chlorophyll *a* concentrations that are indicative of a higher probability of exceeding select health advisory concentrations for microcystin. Using the 2007 National Lakes Assessment and a conditional probability approach, we identify chlorophyll *a* concentrations that are more likely than not to be associated with an exceedance of a microcystin health advisory level. We look at the recent US EPA health advisories for drinking water as well as the World Health Organization levels for drinking water and recreational use and identify a range of chlorophyll *a* thresholds. A 50% chance of exceeding one of the specific advisory microcystin concentrations of 0.3, 1, 1.6, and 2  $\mu\text{g/L}$  is associated with chlorophyll *a* concentration thresholds of 23, 68, 84, and 104  $\mu\text{g/L}$ , respectively. When managing for these various microcystin levels, exceeding these reported chlorophyll *a* concentrations should be a trigger for further testing and possible management action.

## 1 Introduction

Over the last decade, numerous events and legislative activities have raised the public awareness of harmful algal blooms [1–3]. In response the US Environmental Protection Agency (USEPA) has recently released suggested microcystin (one of the more common toxins) concentrations that would trigger health advisories [4–6]. Additionally, the World Health Organization (WHO) has proposed microcystin advisory levels for drinking water and a range of recreational risk levels [7,8]. While these levels and associated advisories are likely to help mitigate the impacts from harmful algal blooms, they are not

36 without complications.

37 One of these complications is that they rely on available measurements of microcystin. While laboratory  
38 testing (e.g., chromatography) remains the gold standard for quantifying microcystin concentrations in  
39 water samples, several field test kits have been developed. Even though field tests provide a much needed  
40 means for rapid assessment, they are not yet widely used and are moderately expensive (approximately  
41 \$150-\$200 depending on specific kit) with a limited shelf life (typically one year) [9,10]. Additionally,  
42 each technique requires nuanced understanding of the detection method (e.g., limit of detection, specific  
43 microcystin variants being measured, and sampling protocol).

44 Fortunately, cyanobacteria and microcystin-LR has been shown to be associated with several other,  
45 more commonly measured and well understood components of water quality that are readily assessed  
46 in the field [11]. For instance, there are small or hand held fluorometers that measure chlorophyll  
47 *a*. Additionally, chlorophyll *a* is a very commonly measured component of water quality that is also  
48 known to be positively associated with microcystin-LR concentrations [12,13]. Recently, Yuan et. al [13]  
49 explored these associations in detail and controlled for other related variables. In their analysis they  
50 find that total nitrogen and chlorophyll *a* show the strongest association with microcystin. Furthermore,  
51 they identify chlorophyll *a* and total nitrogen concentrations that are associated with exceeding 1  $\mu\text{g/L}$  of  
52 microcystin. These findings suggest that chlorophyll *a* concentrations could also track the new USEPA  
53 microcystin health advisory levels for drinking water. Identifying this association would provide an  
54 important tool for water resource managers to help manage the threat to public health posed by CHABs  
55 and would be especially useful in the absence of measured microcystin concentrations.

56 In this paper we build on past efforts and utilize the National Lakes Assessment (NLA) data and  
57 identify chlorophyll *a* concentrations that are associated with higher probabilities of exceeding several  
58 microcystin health advisory concentrations [6,8,14]. We build on past studies by exploring associations  
59 with the newly announced advisory levels and by also applying a different method, conditional probability  
60 analysis. Utilizing different methods strengthens the evidence for suggested chlorophyll *a* levels that are  
61 associated with increased risk of exceeding the health advisory levels as those levels are not predicated  
62 on a single analytical method. So that others may repeat or adjust this analysis, the data, code, and  
63 this manuscript are freely available via <https://github.com/USEPA/microcystinchla>.

## 2 Methods

### 2.1 Data

We used the 2007 NLA chlorophyll *a* and microcystin-LR concentration data [14]. These data represent a snapshot of water quality from the summer of 2007 for the conterminous United States and were collected as part of an ongoing probabilistic monitoring program [14]. Water quality data, including chlorophyll *a* and microcystin-LR were obtained via an integrated sample taken from the surface of the lake down to 2 meters. Samples were taken at the same time from the index site (e.g. near the centroid of the lake) and these provide the source for both chlorophyll *a* and microcystin-LR [14].

For our analysis we only used samples that were part of the probability sampling design (i.e. no reference samples) and from the first visit to the lake (e.g. some lakes were sampled multiple times). The detection limit for microcystin-LR was 0.05  $\mu\text{g/L}$ . Approximately 67% of lakes reported microcystin-LR at the detection limit. For this analysis we retained these values as removing them would erroneously reduce the confidence intervals around the conditional probabilities. Data on chlorophyll *a* and microcystin-LR concentrations are available for 1028 lakes.

### 2.2 Analytical Methods

We used a conditional probability analysis (CPA) approach to explore associations between chlorophyll *a* concentrations and World Health Organization (WHO) and USEPA microcystin health advisory levels [16]. Many health advisory levels have been suggested (Table 1), but lakes with higher microcystin-LR concentrations in the NLA were rare. Only 1.16% of lakes sampled had a concentration greater than 10  $\mu\text{g/L}$ . Thus, for this analysis we focused on the microcystin concentrations that are better represented in the NLA data. These were the USEPA children's (i.e. bottle fed infants to pre-school age children) drinking water advisory level of 0.3  $\mu\text{g/L}$  (USEPA Child), the WHO drinking water advisory level of 1  $\mu\text{g/L}$  (WHO Drinking), the USEPA adult (i.e. beyond pre-school aged individuals) drinking water advisory level of 1.6  $\mu\text{g/L}$  (USEPA Adult), and the WHO recreational, low probability of effect advisory level of 2  $\mu\text{g/L}$  (WHO Recreational).

89 Conditional probability analysis provides information about the probability of observing one event  
90 given another event has also occurred. For this analysis, we used CPA to examine how the conditional  
91 probability of exceeding one of the health advisories changes as chlorophyll *a* increases in a lake. We  
92 expect to find higher chlorophyll *a* concentrations to be associated with higher probabilities of exceeding  
93 the microcystin health advisory levels. We also calculated bootstrapped 95% confidence intervals (CI)  
94 using 1000 bootstrapped samples. Thus, to identify chlorophyll *a* concentrations of concern we identified  
95 the value of the upper 95% CI across a range of conditional probabilities of exceeding each health  
96 advisory level. Using the upper confidence limit to identify a threshold is justified as it ensures that a  
97 given threshold is unlikely to miss a microcystin exceedance.

98 As both microcystin-LR and chlorophyll *a* values were highly right skewed, a log base 10 transformation  
99 was used. Additional details of the specific implementation are available at [https://github.com/](https://github.com/USEPA/microcystinchla)  
100 [USEPA/microcystinchla](https://github.com/USEPA/microcystinchla). A more detailed discussion of CPA is beyond the scope of this paper, but  
101 see Paul et al. [17] and Hollister et al. [18] for greater detail. Lastly, all analyses were conducted  
102 using R version 3.2.2 and code and data from this analysis are freely available as an R package at  
103 <https://github.com/USEPA/microcystinchla>.

104 Lastly, we assessed the ability of these chlorophyll *a* thresholds to predict microcystin exceedance. We  
105 used error matrices and calculate total accuracy as well as the proportion of false negatives. Total  
106 accuracy is the total number of correct predictions divided by total observations. The proportion of  
107 false negatives is the total number of lakes that were predicted to not exceed the microcystin guidelines  
108 but actually did, divided by the total number of observations.

### 109 3 Results

110 In the 2007 NLA, microcystin-LR concentrations ranged from 0.05 to 225  $\mu\text{g/L}$ . Microcystin-LR  
111 concentrations of 0.05  $\mu\text{g/L}$  represent the detection limits. Any value greater than that indicates the  
112 presence of microcystin-LR. Of those lakes with microcystin, the median concentration was 0.51  $\mu\text{g/L}$   
113 and the mean was 3.17  $\mu\text{g/L}$ . Of all lakes sampled, 21% of lakes exceeded the USEPA Child level, 8.8%  
114 of lakes exceeded the USEPA Adult level, 11.7% of lakes exceeded the WHO Drinking level, and 7.3%

of lakes exceeded the WHO Recreational level. Chlorophyll *a*, ranged from 0.07 to 936  $\mu\text{g/L}$  and this captures the range of trophic states from oligotrophic to hypereutrophic. All lakes had detectable levels of chlorophyll *a*. The median concentration was 7.79  $\mu\text{g/L}$  and the mean was 29.63  $\mu\text{g/L}$ . The association between chlorophyll *a* and the upper confidence interval across a range of conditional probability values are shown in Table 2. Specific chlorophyll *a* that are associated with greater than even odds of exceeding the advisory levels were 23, 68, 84, and 104  $\mu\text{g/L}$  for 0.3, 1.0, 1.6, and 2.0  $\mu\text{g/L}$  advisory levels, respectively (Table 2 & Figure ??).

The chlorophyll *a* cutoffs may be used to predict whether or not a lake exceeds the microcystin health advisories. Doing so allows us to compare the accuracy of the prediction as well as evaluate false negatives. Total accuracy of the four cutoffs predicting microcystin exceedances were 74% for the USEPA children's drinking water advisory, 86% for the WHO drinking water advisory, 89% for the USEPA adult drinking water advisory, and 91% for the WHO recreational advisory (Tables 3, 4, 5, & 6). However, total accuracy is only one part of the prediction performance with which we are concerned.

When using the chlorophyll *a* cutoffs as an indicator of microcystin exceedances, the error that should be avoided is predicting that no exceedance has occurred when in fact it has. In other words, we would like to avoid Type II errors and minimize the proportion of false negatives. For the four chlorophyll *a* cut-offs we had a proportion of false negatives of 9%, 8%, 6%, and 5% for the USEPA children's, the WHO drinking water, the USEPA adult, and the WHO recreational advisories, respectively. In each case we missed less than 10% of the lakes that in fact exceeded the microcystin advisory. While this method performs well with regard to the false negative percentage, it is possible that is a relic of the NLA dataset and testing with additional data would allow us to confirm this result.

## 4 Discussion

The log-log association between microcystin-LR and chlorophyll *a* indicates that, in general, higher concentrations of microcystin-LR almost always co-occur with higher concentrations of chlorophyll *a* yet the inverse is not true (Figure ??). Higher chlorophyll *a* is not necessarily predictive of higher microcystin-LR concentrations; however, chlorophyll *a* may be predictive of the probability of exceeding

141 a certain threshold.

142 Indeed, the probability of exceeding each of the four tested health advisory levels increased as a  
143 function of chlorophyll *a* concentration (Figure ??). We used this association to identify chlorophyll *a*  
144 concentrations that were associated with a range of probabilities of exceeding a given health advisory  
145 level (Table 2). For the purposes of this discussion we focus on a conditional probability of 50% or  
146 greater (i.e., greater than even odds to exceed a health advisory level). The 50% conditional probability  
147 chlorophyll *a* thresholds represents 28.6%, 11%, 8.9%, and 7.2% of sample lakes for the USEPA Child,  
148 the WHO Drinking, the USEPA Adult, and the WHO recreational levels, respectively.

149 There are numerous possible uses for the chlorophyll *a* and microcystin advisory cut-off values. First, in  
150 the absence of microcystin-LR measurements, exceedence of the chlorophyll *a* concentrations could be a  
151 trigger for further actions. Given that there is uncertainty around these chlorophyll *a* cutoffs the best  
152 case scenario would be to monitor for chlorophyll *a* and in the event of exceeding a target concentration  
153 take water samples and have those samples tested for microcystin-LR.

154 A second potential use is to identify past bloom events from historical data. As harmful algal blooms  
155 are made up of many species and have various mechanisms responsible for adverse impacts (e.g., toxins,  
156 hypoxia, odors), there is no single definition of a bloom. For cHABs, one approach has been to utilize  
157 phycocyanin to screen for or identify bloom events [19]. This is a useful approach, but phycocyanin  
158 is not always available, thus limiting its utility especially for examining historical data. Using our  
159 chlorophyll *a* cutoffs provides a value that is also associated with microcystin-LR and can be used to  
160 classify lakes, from past surveys, as having bloomed.

161 The values we propose are national and may miss regional variation in water quality, including,  
162 chlorophyll *a* and microcystin-LR [21]. A set of regional conditional probabilities would be interesting;  
163 however, limiting the analysis to the data available per region would make interpretation difficult. The  
164 sample size for each of the conditional probabilities would be reduced (it ranges from 67 to 155) and  
165 the number of lakes in each region that exceed the microcystin values is also reduced. The result is that  
166 our confidence in the conditional probabilities would be less (i.e. greatly increased confidence intervals)  
167 and the relationships less pronounced as we have fewer lakes on which to base the probabilities. Thus,  
168 this dataset is best for making national scale recommendations.

169 Two other limitations with the 2007 National Lakes Assessment data are that they represent a single  
170 sample from that lake and do not capture temporal dynamics and without subsetting the data do not  
171 provide us the ability to validate the presence of microcystin-LR. As of this writing, the 2012 National  
172 Lakes Assessment data are not public. When these data are released, a validation of this approach can  
173 be completed then.

174 Lastly, using chlorophyll *a* is not meant as a replacement for testing of microcystin-LR or other toxins.  
175 It should be used when other, direct measurements of cyanotoxins are not available. In those cases,  
176 which are likely to be common at least in the near future, using a more ubiquitous measurement, such  
177 as chlorophyll *a* will provide a reasonable proxy for the probability of exceeding a microcystin health  
178 advisory level and provide better protection against adverse effects in both drinking and recreational  
179 use cases.

## 180 5 Acknowledgements

181 We would like to thank Anne Kuhn, Bryan Milstead, John Kiddon, Joe LiVolsi, Tim Gleason, Wayne  
182 Munns, and Leslie D'Anglada for constructive reviews of this paper. Special thanks to Jason Marion,  
183 Alan Wilson, and Zofia Taranu for reviews of the submitted manuscript. This paper has not been  
184 subjected to Agency review. Therefore, it does not necessary reflect the views of the Agency. Mention  
185 of trade names or commercial products does not constitute endorsement or recommendation for use.  
186 This contribution is identified by the tracking number ORD-015143 of the Atlantic Ecology Division,  
187 Office of Research and Development, National Health and Environmental Effects Research Laboratory,  
188 US Environmental Protection Agency.

## 189 6 Figures

190 ## Saving 6 x 6.8 in image



Table 1: Various suggested microcystin health advisory concentrations.

Source	Type	Concentration
USEPA	Child Drinking Water Advisory	0.3 $\mu\text{g/L}$
WHO	Drinking Water	1 $\mu\text{g/L}$
USEPA	Adult Drinking Water Advisory	1.6 $\mu\text{g/L}$
WHO	Recreational: Low Prob. of Effect	2-4 $\mu\text{g/L}$
WHO	Recreational: Moderate Prob. of Effect	10-20 $\mu\text{g/L}$
WHO	Recreational: High Prob. of Effect	20-2000 $\mu\text{g/L}$
WHO	Recreational: Very High Prob. of Effect	>2000 $\mu\text{g/L}$

Table 2: Chlorophyll *a* concentrations that are associated with a 50% probability of exceeding a microcystin health advisory concentration.

Cond. Probability	USEPA Child (0.3 $\mu\text{g/L}$ )	WHO Drink (1 $\mu\text{g/L}$ )	USEPA Adult (1.6 $\mu\text{g/L}$ )	WHO Recreation
0.1	0.07	0.07	0.07	1
0.2	0.07	4.00	12.00	17
0.3	3.00	17.00	32.00	45
0.4	11.00	37.00	68.00	77
0.5	23.00	68.00	84.00	104
0.6	39.00	97.00	115.00	185
0.7	66.00	126.00	871.00	871
0.8	116.00	271.00	871.00	871
0.9	170.00	516.00	871.00	871

Table 3: Confusion matrix comparing chlorophyll *a* predicted exceedences (rows) versus real exceedances (columns) for the USEPA childrens drinking water advisory.

Not Exceed	Exceed	Row Totals
Not Exceed	643	95 738
Exceed	168	122 290
Column Totals	811	217 1028

Table 4: Confusion matrix comparing chlorophyll *a* predicted exceedences (rows) versus real exceedances (columns) for the WHO drinking water advisory.

Not Exceed	Exceed	Row Totals
Not Exceed	841	78 919
Exceed	66	43 109
Column Totals	907	121 1028

Table 5: Confusion matrix comparing chlorophyll *a* predicted exceedences (rows) versus real exceedances (columns) for the USEPA adult drinking water advisory.

Not Exceed	Exceed	Row Totals
Not Exceed	884	57 941
Exceed	53	34 87
Column Totals	937	91 1028

Table 6: Confusion matrix comparing chlorophyll *a* predicted exceedences (rows) versus real exceedances (columns) for the WHO recreational water advisory.

Not Exceed	Exceed	Row Totals
Not Exceed	908	51 959
Exceed	45	24 69
Column Totals	953	75 1028

## References

1. Jetoo S, Grover VI, Krantzberg G (2015) The toledo drinking water advisory: Suggested application of the water safety planning approach. *Sustainability* 7: 9787–9808.
2. Rinta-Kanto JM, Konopko EA, DeBruyn JM, Bourbonniere RA, Boyer GL, et al. (2009) Lake erie microcystis: Relationship between microcystin production, dynamics of genotypes and environmental parameters in a large lake. *Harmful Algae* 8: 665–673.
3. Bloom HA, Research H, Act CA (2014) Harmful Algal Bloom and Hypoxia Research and Control Amendments Act of 2014. (S. 1254). Available: <https://www.congress.gov/113/bills/s1254/BILLS-113s1254enr.pdf>.
4. McElhiney J, Lawton LA (2005) Detection of the cyanobacterial hepatotoxins microcystins. *Toxicology and Applied Pharmacology* 203: 219–230.
5. Zurawell RW, Chen H, Burke JM, Prepas EE (2005) Hepatotoxic cyanobacteria: A review of the biological importance of microcystins in freshwater environments. *Journal of Toxicology and Environmental Health, Part B* 8: 1–37.
6. USEPA (2015) Drinking water health advisory for the cyanobacterial microcystin toxins. EPA-820-R-15100. Available: <http://www2.epa.gov/sites/production/files/2015-06/documents/microcystins-report-2015.pdf>.
7. World Health Organization (2003) Cyanobacterial toxins: Microcystin-LR in drinking-water. Background document for development of WHO guidelines for drinking-water quality. Geneva, Switzerland. World Health Organization, 2nd ed. Geneva.
8. Chorus EI, Bartram J (1999) Toxic cyanobacteria in water: A guide to their public health consequences, monitoring and management. World Health Organization.
9. James R, Gregg A, Dindal A, McKernan J (2011) Environmental technology verification report:

215 Abraxis microcystin test kits. Online document. Accessed online: June 22.

216 10. Aranda-Rodriguez R, Jin Z, Harvie J, Cabecinha A (2015) Evaluation of three field test kits to  
217 detect microcystins from a public health perspective. *Harmful Algae* 42: 34–42.

218 11. Yuan LL, Pollard AI (2015) Deriving nutrient targets to prevent excessive cyanobacterial densities  
219 in uS lakes and reservoirs. *Freshwater Biology* 60: 1901–1916.

220 12. Pip E, Bowman L (2014) Microcystin and algal chlorophyll in relation to nearshore nutrient  
221 concentrations in Lake Winnipeg, Canada. *Environment and Pollution* 3: p36.

222 13. Yuan LL, Pollard AI, Pather S, Oliver JL, D’Anglada L (2014) Managing microcystin: Identifying  
223 national-scale thresholds for total nitrogen and chlorophyll a. *Freshwater Biology* 59: 1970–1981.

224 14. USEPA (2009) National lakes assessment: A collaborative survey of the nation’s lakes. EPA  
225 841-R-09-001.

226 15. USEPA (2007) Survey of the nation’s lakes field operations manual EPA841-B-07-004.

227 16. Paul JF, Munns WR (2011) Probability surveys, conditional probability, and ecological risk  
228 assessment. *Environmental Toxicology and Chemistry* 30: 1488–1495.

229 17. Paul JF, McDonald ME (2005) Development of empirical, geographically specific water quality  
230 criteria: A conditional probability analysis approach. *Journal of the American Water Resources*  
231 *Association* 41: 1211–1223.

232 18. Hollister JW, Walker HA, Paul JF (2008) CProb: A computational tool for conducting conditional  
233 probability analysis. *Journal of Environmental Quality* 37: 2392–2396.

234 19. Miller TR, Beversdorf L, Chaston SD, McMahon KD (2013) Spatiotemporal molecular analysis of  
235 cyanobacteria blooms reveals microcystis-aphanizomenon interactions. *PloS one* 8: e74933.

236 20. Marion JW, Lee J, Wilkins III J, Lemeshow S, Lee C, et al. (2012) In vivo phycocyanin flourometry  
237 as a potential rapid screening tool for predicting elevated microcystin concentrations at eutrophic lakes.



238 Environmental science & technology 46: 4523–4531.

239 21. Beaver JR, Manis EE, Loftin KA, Graham JL, Pollard AI, et al. (2014) Land use patterns, ecoregion,  
240 and microcystin relationships in uS lakes and reservoirs: A preliminary evaluation. Harmful Algae 36:  
241 57–62.

242 22. Wagner T, Soranno PA, Webster KE, Cheruvilil KS (2011) Landscape drivers of regional variation in  
243 the relationship between total phosphorus and chlorophyll in lakes. Freshwater Biology 56: 1811–1824.