

Choose Results File

Browse...

NEOSresult_Upper Connecticut River_2026-05-07.txt

Upload complete

Choose the corresponding `01\_UserSpecs\_loadingtargets\_selected.csv` file

Browse...

01_UserSpecs_loadingtargets_selected.csv

Upload complete

Choose WWTP File

Browse...

WWTP_COMIDs.csv

Upload complete

Choose SPARROW Inputs File

Browse...

ne_sparrow_model_input_short.csv

Upload complete

Choose State Cropland Streamcat Files

Browse...

CT_NH_MA_VT_NY_RI_StreamCatData_2019.csv

Upload complete

Important: Selecting 'Preview Uploads' or 'View NEOS Results' before all uploads have completed may produce unexpected results. Previews can be found under the 'File Preview' tab, and generated reports can be found under the 'View Results' tab.

Preview Uploads

View NEOS Results

Postprocessing Step

RBEROST Results Postprocessing Step

File Preview

View Results

View available scenarios

1

▼

Scaled-up Optimization Results

To save this report, open the RBEROST postprocessor in a web browser and use the print to pdf functionality.

User Notes:

Your model has successfully solved.

The total cost to reduce loads to the limit you provided is \$74,674.12.

The model did not implement any WWTP retrofits.

The model chose to implement the total area of the following agricultural BMPs:

Contour Farming (ac)	Cover crops (ac)	Fert 20 (ac)	Filterstrip (ac)	MIN TILL (ac)	Ponds (ac)	Terrace Waterway (ac)	Waterway Only (ac)
529.2	0.0	3,843.9	456.5	3,539.2	0.0	2.6	0.0

Download Agricultural BMPs by COMID (session/9ed3b9f299a2681f9eb9aeb03e50b5fb/download/downloadagdat?w=)

The model chose to implement the total area of the following urban BMPs.

Enhanced Biofiltration w ISR (ac)	Gravel Wetland (ac)	Infiltration Basin (ac)	Infiltration Trench (ac)	Porous Pavement w subsurface infiltration (ac)
0.22	0.22	383.2	0.93	0.94

Download Urban BMPs by COMID (session/9ed3b9f299a2681f9eb9aeb03e50b5fb/download/downloadurbandat?w=)

The model did not implement any riparian buffer BMPs.

The model did not implement any septic BMPs.