

StreamCatTools: An R package for working with StreamCat and LakeCat watershed data in R

Marc H. Weber¹, Ryan A. Hill¹, Selia Markley⁴, Travis Hudson³, and Alan Brookes²

¹ Office of Water, United States Environmental Protection Agency ² United States Environmental Protection Agency Retired ³ Oak Ridge Associated Universities Student Services Contractor c/o United States Environmental Protection Agency ⁴ Oak Ridge Institute for Science and Education Fellow c/o United States Environmental Protection Agency

DOI:

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Submitted:

Published:

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

Summary

StreamCatTools provides functions for easily working with, visualizing and analyzing StreamCat (Hill et al., 2016) and LakeCat (Hill et al., 2018) watershed metrics within **R**. The StreamCat and LakeCat datasets provide hundreds of landscape metrics for both the local catchment (e.g. landscape draining to a particular stream reach) and full watershed (i.e., the local catchment and all upstream catchments) for every stream reach and lake depicted in the medium resolution National Hydrography Dataset Plus Version 2.1 (NHDPlusV21)(McKay et al., 2012) for the contiguous United States (CONUS). **StreamCatTools** functions wrap the application programming interface (API) for the StreamCat and LakeCat online database and facilitate their simple, straightforward access and use within R.

Statement of Need

Easily accessible, robust, and consistent watershed data underpin hydrology research, water quality monitoring programs, and predictive modelling applications, to name just a few examples. The StreamCat (Hill et al., 2016) and LakeCat (Hill et al., 2018) datasets fill this need by providing nationally consistent curated watershed data for CONUS that have had stringent quality control applied. The data encompass hundreds of watershed metrics for every stream reach and lake feature represented in the NHDPlusV21 (McKay et al., 2012). **StreamCatTools** provides easily accessible watershed metrics for CONUS via: (1) a simple interface to the StreamCat and LakeCat web services in the R programming language; (2) convenient functionality to find available StreamCat and LakeCat metric names and information about variables; and (3) extraction of StreamCat and LakeCat metrics by COMID (a unique identifier in the NHDPlusV21 framework), state, county, NHD hydro-region, or for all of CONUS. Providing these valuable watershed data via web services in R follows the FAIR principles (Fair, Accessible, Interoperable, and Reusable) laid out in (Wilkinson et al., 2016).

Package Overview

StreamCatTools provides a simple streamlined set of functions to easily query and ingest watershed landscape metrics into an R session. Figure 1 shows the overall framework




Figure 1: Diagram of the StreamCat and StreamCatTools framework. The backend Oracle database and REST web service are exposed through api.gov to functions in StreamCatTools that simplify access and analysis of the data in R via the application programming interface (API). The example here shows the general categories of Oracle database tables in the StreamCat Oracle database.

of the StreamCat database, application programming interface, and functionality in the package that simplifies data access in R using web services for the StreamCat and LakeCat datasets.

The core functions in `StreamCatTools` leverage the `httr2` library (Wickham, 2025) for a modern, pipeable method for working with APIs. Specifically, `StreamCatTools` simplifies calls to the API for StreamCat and LakeCat data within the R programming language to allow a user to: (1) access details on available StreamCat and LakeCat metrics; (2) match users' sites to NHDPlusV21 waterbodies (streams or lakes) to access site-relevant StreamCat and LakeCat data; and (3) retrieve StreamCat and LakeCat watershed metrics by COMID, state, county, hydroregion, or for all of CONUS. Additionally, several convenience functions to retrieve year-specific data for variables with time series data, such as the National Land Cover Database (NLCD) (U.S. Geological Survey, 2019) and the National Nutrient Inventory (NNI) (Brehob et al., 2025). Lastly, there is plotting functionality for use with NNI metrics as well as a function to retrieve the watersheds for specific lake features as an `sf` object. Users also can access watersheds for specific COMIDs using the `hydrogeofetch` package for visualization and plotting of watersheds along with StreamCat or LakeCat metrics.

You can install the most recent version of `StreamCatTools` from *CRAN* by running:

```
install.packages("StreamCatTools")
```

You can install the most recent package version from *GitHub* by running the following code (note we use `pak` as the replacement for `remotes` for package installation from GitHub):

```
install.packages("pak")
library(pak)
pkg_install("github::USEPA/StreamCatTools")
```

`StreamCatTools` is loaded into an **R** session:

```
library(StreamCatTools)
```

`StreamCatTools` includes several functions to retrieve basic metric metadata. First, users can list metric names and find out more about StreamCat and LakeCat data available in StreamCat using the `sc_get_params()` and `lc_get_params()` functions. The parameters `aoi` and `metric_names` provide a user with the available areas of interest for metrics and names of metrics. The areas of interest available for all metrics in StreamCat and LakeCat are 'cat' and 'ws' (catchment and watershed). Some metrics in StreamCat are also available at the 100 meter riparian buffer scale ('catrp100' and 'wsrp100'), and for certain metrics the designation of 'other' needs to be used (where the metric is in-stream or derived and not part of the other listed areas of interest). Extracting this information in `StreamCatTools` looks like this:

```
aois <- sc_get_params(param='aoi')
aois
```

```
#> [1] "cat"          "catrp100" "other"      "ws"          "wsrp100"
```

```
names <- sc_get_params(param='metric_names')
names[1:10]
```

```
#> [1] "agkffact"      "al2o3"        "bankfulldepth" "bankfullwidth"
#> [5] "bfi"           "canaldens"    "cao"           "cbnf"
#> [9] "chem"          "clay"
```

And the same information for LakeCat metrics is derived in similar fashion:

```
aois <- lc_get_params(param='aoi')
aois
```

```
#> [1] "cat" "ws"
```

```
names <- lc_get_params(param='metric_names')
names[1:10]
```

```
#> [1] "agkffact"      "al2o3"        "bfi"           "canaldens"    "cao"
#> [6] "cbnf"          "clay"         "coalminedens" "compstrgth"   "damdens"
```

StreamCat and LakeCat are built around the concepts of local drainage area (i.e catchment) and watershed (i.e., the local drainage area and all upstream catchments) (Hill et al., 2016). This approach uses the NHDPlusV21 hydrographic framework of catchments as the building block for summarizing information represented in landscape data. Watershed data are produced by combining both the catchment landscape summary and all upstream catchment summaries using either the weighted average of metrics (in most cases) or a sum or count (for certain metrics). `sc_get_params()` and `lc_get_params()` also include a 'variable_info' parameter to return more detailed metadata for metrics including both short and long metric descriptions, years available (if applicable), units, and the metric category (Table 1).

Table 1: Example of variable information returned by the `variable_info` parameter in the `sc_get_params` function.

Metric	Short Description	Year	Units
agkffact[AOI]	Ag Soil Erodibility Kf Factor	NA	Unitless
bfi[AOI]	Base Flow Index	NA	Percent
huden[Year][AOI]	Mean Housing Density	2010	Count/Square Kilometer
inorgnwetdep[Year][AOI]	Mean Annual Precipitation-Weighted Nitrogen	2008	Kilogram/Hectare/Year
n_ags_[Year][AOI]	Nitrogen Agricultural Surplus	1987-2017	Kilograms
pcthighsev[Year][AOI]	Percent High Burn Severity Class For Year	1984-2018	Percent

Additional functions for getting metadata about the underlying StreamCat and LakeCat data include the `sc_fullname()` and `lc_fullname()` functions which provide the descriptive full name for any given metric in StreamCat or LakeCat:

```
sc_fullname(metric='pctgrs2019')
```

```
#> [1] "Grassland/Herbaceous Percentage 2019"
```

Users can also filter metric names and information by the metric year(s), the indicator categories for metrics, the metric dataset names, or the areas of interest available for a given metric using the `sc_get_metric_names` or `lc_get_metric_names` functions (Table 2).

```
metrics <- sc_get_metric_names(category = c('Deposition', 'Climate'),
                              aoi=c('Cat', 'Ws'))
my_data <- head(metrics[,c('Category', 'Metric', 'AOI')], 10)
```

Table 2: Example of metric names returned by the `sc_get_metric_names` function.

Category	Metric	AOI
Climate	bfi[AOI]	Cat, Ws
Deposition	inorgnwetdep[Year][AOI]	Cat, Ws
Deposition	nh4[Year][AOI]	Cat, Ws
Deposition	no3[Year][AOI]	Cat, Ws
Climate	precip8110[AOI]	Cat, Ws
Climate	precip9120[AOI]	Cat, Ws
Climate	precip[Year][AOI]	Cat, Ws
Deposition	sn[Year][AOI]	Cat, Ws
Climate	tmax8110[AOI]	Cat, Ws
Climate	tmax9120[AOI]	Cat, Ws

More details on these functions can be found at the [package introduction page](#).

The primary package functionality is in the `sc_get_data` and `lc_get_data` functions which allow users to extract catchment or watershed metrics of interest by providing NHDPlusV21 COMIDs for streams or lakes within the database. The following example shows a request for two catchment and watershed metrics for three stream reaches:

```
df <- sc_get_data(metric='pcturbmd2019,damdens',
                  aoi='cat,ws',
                  comid='179,1337,1337420')
```

The first line requests percent of NLCD medium intensity developed land cover in 2019 and the density of dams. The second line specifies that the geographic area of interest (aoi) is the drainage to the local reach scale (i.e. 'cat', short for catchment) and the full watershed. The final line requests these data for three NHDPlusV21 stream segments specified by their unique COMIDs. A similar request for LakeCat metrics for data on lakes might look like this, where a user can ask for data for a particular county using the county FIPS code rather than for a set of identifying COMIDs:

```
df <- lc_get_data(metric='pctwdwet2006', aoi='ws', county='41003')
head(df)
```

```
#>      comid pctwdwet2006ws
#> 1 23769033      0.0000000
#> 2 23769029      0.1701645
#> 3 23769021      2.4697951
#> 4 23769003      0.0000000
#> 5 23768999      8.1185071
#> 6 23768983      0.0000000
```

Users can also request data for one or more state(s), county(ies), hydroregion(s), or all of CONUS (though asking for data for all of CONUS may time out if asking for too many metrics at once). The following example demonstrates a request data for a particular county in Oregon:

```
df <- sc_get_data(metric='pctwdwet2006', aoi='ws', county='41003')
```

StreamCatTools provides a convenience function in either `sc_get_params` or `lc_get_params` to discover the FIPS codes for states and counties to make it easier for users to query data by these parameters. For example, the following code chunk shows how to get the FIPS code for Benton County, Oregon to query data for that county:

```
df <- sc_get_params(param='county') |> dplyr::filter(state == 'OR' &
  ↪ county_name == 'Benton County') |> dplyr::pull(fips)
```

Or we can ask for several different metrics for all of a state:

```
df <- sc_get_data(metric='clay,agkffact', aoi='ws', state='CT')
head(df)
```

```
#>      comid agkffactws  clayws
#> 1 7702768      0.0177 6.100716
#> 2 7702200      0.0009 6.243493
#> 3 7702202      0.0011 6.368938
#> 4 7701112      0.0177 6.080299
#> 5 7701168      0.0054 6.092434
#> 6 7701214      0.0040 6.082292
```

Additionally, convenience functions are provided for accessing the NLCD and NNI datasets, respectively, using `sc_get_nlcd` and `lc_get_nlcd` and `sc_get_nni` and `lc_get_nni`:

```
df <- sc_get_nlcd(comid='1337420', year='2019', aoi='ws')

df <- sc_get_nni(year='1987, 1990, 2005, 2017', aoi='cat,ws',
  comid='179,1337,1337420')
```

Applications and Discussion

One can easily summarize the data returned from `sc_get_data` or `lc_get_data` for a set of rivers or lakes or for the watersheds for an entire county or state in a number of ways. The following example calculates the mean percent of agriculture on slopes > 10% within a watershed for all watersheds in Benton County, Oregon:

```
df <- sc_get_data(metric='pctagslphigh2019', aoi='ws', county='41003') |>
  dplyr::summarise(mean_pctagslphigh2019ws = mean(pctagslphigh2019ws,
    na.rm=TRUE))
df

#>   mean_pctagslphigh2019ws
#> 1                4.406472
```

Watershed metrics from `StreamCatTools` can also be easily visualized with functions from `hydrogeofetch` (Blodgett, 2026) and `ggplot2` (Wickham, 2016).

```
library(hydrogeofetch)
library(ggplot2)
library(ggspatial)
library(StreamCatTools)

start_comid = 23763517
nldi_feature <- list(featureSource = "comid", featureID = start_comid)
flowline_nldi <- hydrogeofetch::navigate_nldi(nldi_feature, mode = "UT",
  data_source = "flowlines", distance=5000)
df <- sc_get_data(metric='pctimp2019', aoi='cat',
  comid=flowline_nldi$UT_flowlines$nhdplus_comid)
flowline_nldi <- flowline_nldi$UT_flowlines
flowline_nldi$PCTIMP2019 <-
  df$pctimp2019cat[match(flowline_nldi$nhdplus_comid, df$comid)]
basin <- hydrogeofetch::get_nldi_basin(nldi_feature = nldi_feature)
```

Figure 2 plots the NLCD percent imperiousness (percentage of area covered by constructed, artificial surfaces) for the the local drainage (catchment in NHDPlusV21 syntax) and displays the values mapped to each stream reach and to the overall basin boundary.

Watersheds for lakes can also be retrieved using the `lc_get_watershed` function to visualize LakeCat metrics along with the plotted watersheds for lake features as shown in Figure 3.

```
library(ggplot2)
library(patchwork)
library(ggforce)
```




Figure 2: Map of NLCD percent imperviousness for each catchment for the Calapooia River watershed in Oregon.

```
df <- lc_get_nlcd(comid='19334077', year='2019', aoi='ws')
lake <- hydrogeofetch::get_waterbodies(id = 19334077)
ws <- lc_get_watershed(comid = 19334077, huc2 = "01", huc2_filter = "01",
  threads = 2, retries = 5, verbose=FALSE,
  ↪ progress=FALSE)
```

Functions to specifically plot NNI metrics in StreamCat have been developed and are also available (Markley et al., 2026), such as Figure 4 showing the annual time series of nitrogen and phosphorus budget data for a given watershed such as the Mississippi-Atchafalaya River Basin:

```
library(StreamCatTools)
library(ggplot2)
library(ggpattern)
com <- '22812041'
sc_plotnni(comid = com, include.nue = TRUE)
```

Potential future functionality for **StreamCatTools** includes expanding the scope of plotting functions as well as expanding the range of metrics and ease of querying these metrics. Additionally, a complementary package, **StreamCatR** is being developed that will allow users to quickly and easily process their own landscape metrics to the NHDPlus framework or other hydrologic frameworks with network topology. **StreamCatTools** currently facilitates easy ingestion of StreamCat and LakeCat watershed landscape metrics into workflows in R which is of great use to state watershed planners, researchers and non-governmental organizations and borne out by the over 5000 package downloads and over 350 citations of the underlying StreamCat and LakeCat data that are served by the **StreamCatTools** package.




Figure 3: NLCD land cover proportions with an example lake watershed.




Figure 4: Annual time series of nitrogen and phosphorus budget data for the Mississippi=Atchafalaya river basin.

Acknowledgements

Examples of using StreamCat and LakeCat make extensive use of `hydrogeofetch` (Blodgett, 2026) and the functions for accessing the API are facilitated through use of `httr2` (Wickham, 2025). Figures were created using `ggplot2` (Wickham, 2016).

We would like to sincerely thank the editor and reviewers for all of their helpful feedback which greatly improved both the software and the manuscript.

The United States Environmental Protection Agency (EPA) GitHub project code is provided on an “as is” basis and the user assumes responsibility for its use. EPA has relinquished control of the information and no longer has responsibility to protect the integrity, confidentiality, or availability of the information. Any reference to specific commercial products, processes, or services by service mark, trademark, manufacturer, or otherwise, does not constitute or imply their endorsement, recommendation or favoring by EPA. The EPA seal and logo shall not be used in any manner to imply endorsement of any commercial product or activity by EPA or the United States Government. The information in this document has been funded entirely by the United States Environmental Protection Agency (USEPA), in part through appointments to the USEPA’s Internship/Research Participation Program at the Office of Research and Development administered by the Oak Ridge Institute for Science and Education through an interagency agreement. The views expressed in this article are those of the authors and do not necessarily represent the views or policies of the USEPA.

References

- Blodgett, D. (2026). *Hydrogeofetch: Hydrologic geospatial fabric data extraction tool chain* (Version 2.0.0). U.S. Geological Survey; U.S. Geological Survey. [TBD](#)
- Brehob, M. M., Pennino, M. J., Compton, J. E., Zhang, Q., Weber, M. H., Hill, R. A., Markley, S., Pickard, B., Keefer, M., Stackpoole, S. M., Knose, L. A., Ruiz-Mercado, G. J., Clark, C. M., Rea, A. W., Carleton, J. N., Lin, J., Bash, J. O., Foley, K. M., Hogrefe, C., & Sabo, R. D. (2025). The US EPA’s national nutrient inventory: Critical shifts in US nutrient pollution sources from 1987 to 2017. *Environmental Science & Technology*, 59(51), 27836–27852. <https://doi.org/10.1021/acs.est.5c08196>
- Hill, R. A., Weber, M. H., Debbout, R. M., Leibowitz, S. G., & Olsen, A. R. (2018). The lake-catchment (LakeCat) dataset: Characterizing landscape features for lake basins within the conterminous USA. *Freshwater Science*, 37(2), 208–221. <https://doi.org/10.1086/697966>
- Hill, R. A., Weber, M. H., Leibowitz, S. G., Olsen, A. R., & Thornbrugh, D. J. (2016). The stream-catchment (StreamCat) dataset: A database of watershed metrics for the conterminous united states. *JAWRA Journal of the American Water Resources Association*, 52(1), 120–128. <https://doi.org/10.1111/1752-1688.12372>
- Markley, S., Weber, M. H., Hill, R. A., Brehob, M., Alford, S., Sabo, R., Pennino, M., & Compton, J. (2026). StreamCatNNI and LakeCatNNI: Watershet and local catchment nitrogen and phosphorus budgets for US streams and lakes. *Nature Data*.
- McKay, L., Bondelid, T., Dewald, T., Johnston, J., Moore, R., & Reah, A. (2012). *NHDPlus version 2: User guide*. http://www.horizon-systems.com/NHDPlus/NHDPlusV2_home.php
- U.S. Geological Survey. (2019). *National land cover database*. <https://www.mrlc.gov/.%E2%80%AFhttps://doi.org/10.5066/P94UXNTS>
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York. <https://doi.org/10.1007/978-0-387-98141-3>
- Wickham, H. (2025). *httr2: Perform HTTP requests and process the responses*. <https://httr2.r-lib.org>

Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., Silva Santos, L. B. da, Bourne, P. E., et al. (2016). The FAIR guiding principles for scientific data management and stewardship. *Scientific Data*, 3(1), 1–9. <https://doi.org/10.1038/sdata.2016.18>