

1) Definition

$f(n[])$:

$g(n[])$

$k(n[])$

$v[, , , ,]$

$n[, , , ,]$


$n[, , , ,]$

$sum++ \mid V_i > V_n$

$sum++ \mid n_i \% 2$

$sum++ \mid n_i \% 2$

2) Analysis



$n[]$

$sum1$

$sum2$

$sum3$

max

$for(i: MAX)$

$si \ n_i \% 2$

$sum1++$

$sum1$

$sum2$

$sum3$

$for(i: MAX)$

$si \ n_i \% 2$

$sum2++$

$max \leftarrow n_0$

$for(i: MAX)$

$si \ n_i < n;$

$max \leftarrow n_i$

$for(j: MAX)$


$si \ n_i == max$

$sum3++$

1) Definition

 $F(n):$
 $V[1..n]$
 $\forall x \in V \quad \text{rand}() \% 75, x < 25 : x = 25$
 $g(n, m):$
 $M[1..n][1..m] \quad \forall x \in M \{$
 $\text{if } j \% 2 = 0$
 $\text{rand} \% 50 + 50$
 $K(n, m)$
 $M[1..n][1..m] \quad \forall x \in M \{$
 $\text{if } j \% 2 = 1$
 $\text{rand} \% 30$

2) Analysis



$n1$ for($i: n1$)
 $n1$ $M1[i][1]$ $V[i] \leftarrow \text{rand}() \% 75$
 $n1$ $\text{if } V[i] < 25$
 $n1$ $V[i] = 25$

$n1$ $M1[i][2]$
 $n1$ for($i: n2$)
 $n1$ for($j: m1$)

 $\text{if } j \% 2 = 0$
 $M[i][j] \leftarrow (\text{rand}() \% 50) + 50$
 else
 0
 $\text{for } (i: n3)$
 $\text{for } (j: m2)$
 $\text{if } j \% 2 = 1$
 $M[i][j] \leftarrow \text{rand}() \% 50$
 else
 0

 Master $f(n)$

 Master $g(n, m)$

 Master $K(n, m)$