Southeast Conservation Blueprint Summary

for Kentucky

Created 10/16/2026

Table of Contents

About the Southeast Blueprint 3

Southeast Blueprint Priorities 4

Hubs and Corridors 6

Indicator Summary 8

More Information 50

The Southeast Conservation Blueprint 2026

Southeast Conservation Adaptation Strategy

[THIS PAGE INTENTIONALLY LEFT BLANK]

About the Southeast Blueprint

The Southeast Conservation Blueprint is the primary product of the Southeast Conservation Adaptation Strategy (SECAS). It is a living, spatial plan to achieve the SECAS vision of a connected network of lands and waters across the Southeast and Caribbean. The Blueprint is regularly updated to incorporate new data, partner input, and information about on-the-ground conditions.

The Blueprint identifies priority areas based on a suite of natural and cultural resource indicators representing terrestrial, freshwater, and marine ecosystems. A connectivity analysis identifies corridors that link coastal and inland areas and span climate gradients.

For more information:

- Visit the Blueprint webpage
- Review the Blueprint 2026 Development Process
- View and download the Blueprint data and make maps on the Blueprint page of the SECAS Atlas

We're here to help!


- Do you have a question about the Blueprint?
- Would you like help using the Blueprint to support a proposal or inform a decision?
- Do you have a suggestion on how to improve the Blueprint? The Blueprint and its inputs are regularly revised based on input from people like you.
- Do you have feedback on how to improve the Blueprint Explorer interface?

If you need help or have questions, contact Southeast Blueprint staff by reaching out to a member of the user support team.

We're here to support you. We really mean it. It's what we do!


This report was generated by the Southeast Conservation Blueprint Explorer, which was developed by Astute Spruce, LLC in partnership with the U.S. Fish and Wildlife Service under the Southeast Conservation Adaptation Strategy.

Southeast Blueprint Priorities



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Priorities for a connected network of lands and waters

- Highest priority
- High priority
- Medium priority
- Priority connections

Priority Categories

For a connected network of lands and waters

In total, Blueprint priorities and priority connections cover roughly 50% of the Southeast Blueprint geography.

Highest priority

Areas where conservation action would make the biggest impact, based on a suite of natural and cultural resource indicators. This class covers roughly 10% of the Southeast Blueprint geography.

High priority

Areas where conservation action would make a big impact, based on a suite of natural and cultural resource indicators. This class covers roughly 15% of the Southeast Blueprint geography.

Medium priority

Areas where conservation action would make an above-average impact, based on a suite of natural and cultural resource indicators. This class covers roughly 20% of the Southeast Blueprint geography.

Priority connections

Connections between priority areas that cover the shortest distance possible while routing through as much Blueprint priority as possible. This class covers roughly 5% of the Southeast Blueprint geography.


Table 1: Extent of each Blueprint priority category within Kentucky.

Priority Category	Acres	Percent of Area
Highest priority	2,375,522	9.2%
High priority	3,684,460	14.2%
Medium priority	5,564,503	21.5%
Priority connections	1,547,616	6.0%
Lower priority	12,685,112	49.1%
Total area	**25,857,213**	**100%**

Hubs and Corridors


The Blueprint uses a least-cost path connectivity analysis to identify corridors that link hubs across the shortest distance possible, while also routing through as much Blueprint priority as possible.

In the continental Southeast, hubs are large patches (~5,000+ acres) of highest priority Blueprint areas and/or protected lands.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Hubs
Corridors

Table 2: Extent of hubs and corridors within Kentucky.

Type	Acres	Percent of Area
Hubs	2,402,302	9.3%
Corridors	5,234,879	20.2%
Not a hub or corridor	18,220,032	70.5%
Total area	**25,857,213**	**100%**

Indicator Summary

Table 3: Terrestrial indicators.

Indicator	Present
East Coastal Plain open pine birds	-
Fire frequency	✓
Grassland & savanna extent	✓
Grassland & savanna restoration	✓
Greenways & trails	✓
Imperiled amphibians & reptiles	✓
Imperiled mammals	✓
Intact habitat cores	✓
Landscape condition	✓
Mississippi Alluvial Valley forest birds - protection	✓
Mississippi Alluvial Valley forest birds - reforestation	✓
Potential access to parks	✓
Resilient terrestrial sites	✓
River cane restoration	✓
South Atlantic forest birds	-
South Atlantic low-urban historic landscapes	-
Urban park size	✓
West Coastal Plain & Ouachitas forested wetland birds	-
West Coastal Plain & Ouachitas open pine birds	-
West Gulf Coast mottled duck nesting	-


Table 4: Freshwater indicators.

Indicator	Present
Atlantic migratory fish habitat	-
Floodplain inundation	✓
Gulf migratory fish connectivity	-
Imperiled aquatic species	✓
Lakes & reservoirs	✓
Natural landcover in floodplains	✓
Network complexity	✓
Permeable surface	✓

Terrestrial


Fire frequency

This indicator uses remote sensing, fire treatment data, and historical landcover models to assess whether an area matches its historical fire regime now and in the recent past (1984-2024). It focuses on fire-adapted forests and grasslands that historically burned at least once every 20 years, which rely on regular, low-intensity fires to maintain habitat, support native plant growth, and reduce wildfire risk. It evaluates departure from historical conditions based on the U.S. Forest Service's fire regime condition classes. This indicator combines Landsat-based fire history metrics from the U.S. Geological Survey, fire treatment data from LANDFIRE, and historical fire regime estimates from LANDFIRE. Development of fire history metrics was supported by Southeast FireMap.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



How well fire-adapted areas match their historic fire regime under current conditions vs. recent past (1984-2024)

- Currently in historical fire regime, low (<33%) departure in recent past
- Currently in historical fire regime, medium (33-66%) departure in recent past
- Currently in historical fire regime, high (>66%) departure in recent past
- Not currently in historical fire regime, low (<33%) departure in recent past
- Not currently in historical fire regime, medium (33-66%) departure in recent past
- Not currently in historical fire regime, high (>66%) departure in recent past
- Urban, row crop, or not a fire-adapted area (>20-year fire regime)

Table 5: Indicator values for fire frequency within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: How well fire-adapted areas match their historic fire regime under current conditions vs. recent past (1984-2024)	Acres	Percent of Area	
↑ High	Currently in historical fire regime, low (<33%) departure in recent past	16,532	<0.1%	
	Currently in historical fire regime, medium (33-66%) departure in recent past	45,652	0.2%	↑ In good condition
	Currently in historical fire regime, high (>66%) departure in recent past	58,131	0.2%	↓ Not in good condition
	Not currently in historical fire regime, low (<33%) departure in recent past	954	<0.1%	
	Not currently in historical fire regime, medium (33-66%) departure in recent past	49,014	0.2%	
	Not currently in historical fire regime, high (>66%) departure in recent past	10,213,205	39.5%	
↓ Low	Urban, row crop, or not a fire-adapted area (>20-year fire regime)	15,473,726	59.8%	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Grassland & savanna extent

This indicator represents grasslands and savannas in the southeastern United States, which support important plants, reptiles, amphibians, mammals, birds, and pollinators. It considers known grassland and savanna locations, predicted locations within protected areas, and surrounding pollinator buffers. It also incorporates other potential grassland and savanna locations within natural and altered landscapes, and restoration opportunities within historical locations based on past fire intervals and historic ecosystem predictions. This indicator combines data from multiple sources, including the Southeastern Grasslands Institute, the National Land Cover Database, LANDFIRE, Oklahoma and Texas ecological systems maps, and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- Known grassland/savanna
- Likely grassland/savanna >10 acres
- Likely grassland/savanna ≤10 acres
- Pollinator buffer around known or likely grassland/savanna
- Potential grassland/savanna in grassland/savanna hub
- Potential grassland/savanna outside grassland/savanna hub
- Historic grassland/savanna
- Not identified as grassland/savanna

Table 6: Indicator values for grassland & savanna extent within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Known grassland/savanna	620	<0.1%	
	Likely grassland/savanna >10 acres	73,252	0.3%	
	Likely grassland/savanna ≤10 acres	25,697	<0.1%	↑ In good condition
	Pollinator buffer around known or likely grassland/savanna	308,749	1.2%	↓ Not in good condition
	Potential grassland/savanna in grassland/savanna hub	1,547,830	6.0%	
	Potential grassland/savanna outside grassland/savanna hub	2,387,657	9.2%	
	Historic grassland/savanna	8,756,008	33.9%	
↓ Low	Not identified as grassland/savanna	12,757,399	49.3%	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Grassland & savanna restoration

This indicator represents potential restoration opportunities for grasslands and savannas within their historical range in the southeastern United States. Grasslands support important plants, reptiles, amphibians, mammals, birds, and pollinators, but have significantly declined from their current extent. It considers proximity to clusters of existing grasslands and savannas, protected areas with management potential, landcover classes with good restoration potential, priority areas for restoration of longleaf and shortleaf pine, and historical grassland and savanna locations. This indicator combines data from multiple sources, including the Southeast Blueprint grassland and savanna extent indicator, the Longleaf Suitability Analysis, shortleaf pine suitability, landcover from the National Land Cover Database and LANDFIRE, and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Restoration priority

- Highest (in a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)
- Very high (in a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)
- High (outside a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)
- Medium (outside a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)
- Low (other historic grassland/savanna)
- Very low (already known or likely grassland/savanna)
- Lowest (not identified as historic grassland/savanna)

Table 7: Indicator values for grassland & savanna restoration within Kentucky. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Restoration priority	Acres	Percent of Area
↑ High	Highest (in a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)	582,269	2.3%
	Very high (in a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)	1,282,977	5.0%
	High (outside a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)	141,330	0.5%
	Medium (outside a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)	2,128,367	8.2%
	Low (other historic grassland/savanna)	8,728,624	33.8%
	Very low (already known or likely grassland/savanna)	99,569	0.4%
↓ Low	Lowest (not identified as historic grassland/savanna)	12,894,077	49.9%
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Greenways & trails

This cultural resource indicator measures both the natural condition and connected length of greenways and trails to characterize the quality of the recreational experience. Natural condition is based on the amount of impervious surface surrounding the path. Connected length captures how far a person can go without leaving a dedicated path, based on common distances for walking, running, and biking. This indicator originates from OpenStreetMap data and the National Land Cover Database.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- Mostly natural and connected for ≥40 km
- Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km
- Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km
- Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km
- Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km
- Developed and connected for <1.9 km
- Sidewalk
- Not identified as a trail, sidewalk, or other path

Table 8: Indicator values for greenways & trails within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Mostly natural and connected for ≥40 km	3,994	<0.1%	
	Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km	8,838	<0.1%	
	Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km	6,592	<0.1%	
	Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km	5,179	<0.1%	↑ In good condition
	Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km	2,829	<0.1%	↓ Not in good condition
	Developed and connected for <1.9 km	3,454	<0.1%	
	Sidewalk	15,705	<0.1%	
↓ Low	Not identified as a trail, sidewalk, or other path	25,810,622	99.8%	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Imperiled amphibians & reptiles

This indicator identifies potential habitat to support amphibian and reptile Regional Species of Greatest Conservation Need (RSGCN). Using a rarity-weighted richness approach, it prioritizes places important for species that are restricted to smaller ranges, such as narrow endemics, which are therefore more vulnerable to habitat loss or change. RSGCN are regional priority species derived from the list of SGCN identified in Southeast State Wildlife Action Plans as most in need of conservation action. RSGCN were chosen based on consistent criteria, such as level of conservation concern, regional stewardship responsibility, and ecological significance. This indicator uses 190 species and subspecies models from the U.S. Geological Survey Gap Analysis Project that are based on known ranges, habitat relationships, and LANDFIRE landcover.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Importance for range-restricted species

- Highest importance (top 5%)
- Very high importance (upper 5-10%)
- High importance (upper 10-20%)
- Medium importance (above average)
- Low importance (below average)
- Developed landcover or no potential habitat

Table 9: Indicator values for imperiled amphibians & reptiles within Kentucky. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Importance for range-restricted species	Acres	Percent of Area
↑ High	Highest importance (top 5%)	362,965	1.4%
	Very high importance (upper 5-10%)	1,831,804	7.1%
	High importance (upper 10-20%)	4,252,634	16.4%
	Medium importance (above average)	5,105,244	19.7%
	Low importance (below average)	8,492,313	32.8%
↓ Low	Developed landcover or no potential habitat	5,812,252	22.5%
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Imperiled mammals

This indicator identifies potential habitat to support mammal Regional Species of Greatest Conservation Need (RSGCN). Using a rarity-weighted richness approach, it prioritizes places important for species that are restricted to smaller ranges, such as narrow endemics, which are therefore more vulnerable to habitat loss or change. RSGCN are regional priority species derived from the list of SGCN identified in Southeast State Wildlife Action Plans as most in need of conservation action. RSGCN were chosen based on consistent criteria, such as level of conservation concern, regional stewardship responsibility, and ecological significance. This indicator uses 58 species models from the U.S. Geological Survey Gap Analysis Project that are based on known ranges, habitat relationships, and LANDFIRE landcover.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Importance for range-restricted species

- Highest importance (top 5%)
- Very high importance (upper 5-10%)
- High importance (upper 10-20%)
- Medium importance (above average)
- Low importance (below average)
- Developed landcover or no potential habitat

Table 10: Indicator values for imperiled mammals within Kentucky. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Importance for range-restricted species	Acres	Percent of Area
↑ High	Highest importance (top 5%)	2,444,774	9.5%
	Very high importance (upper 5-10%)	4,090,578	15.8%
	High importance (upper 10-20%)	3,615,552	14.0%
	Medium importance (above average)	4,793,941	18.5%
	Low importance (below average)	7,552,503	29.2%
↓ Low	Developed landcover or no potential habitat	3,359,866	13.0%
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Intact habitat cores

This indicator represents the size of large, unfragmented patches of natural habitat. It identifies minimally disturbed natural areas at least 100 acres in size and greater than 200 meters wide. Large areas of intact natural habitat are important for many wildlife species, including reptiles and amphibians, birds, and large mammals. This indicator originates from Esri's green infrastructure data.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- Large core (>10,000 acres)
- Medium core (>1,000-10,000 acres)
- Small core (>100-1,000 acres)
- Not a core

Table 11: Indicator values for intact habitat cores within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Large core (>10,000 acres)	2,619,469	10.1%	
	Medium core (>1,000-10,000 acres)	5,640,142	21.8%	
	Small core (>100-1,000 acres)	1,995,665	7.7%	↑ In good condition
↓ Low	Not a core	15,601,937	60.3%	↓ Not in good condition
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Landscape condition

This indicator represents natural areas with limited human alteration while also considering the naturalness of the surrounding landscape. Examples of human alteration include urban development and intense agricultural use. The degree of naturalness across the landscape is a key ecological condition for sustaining species and ecosystem services that are sensitive to habitat fragmentation at multiple scales. This indicator combines data from multiple sources, including the National Land Cover Database; various data on grasslands, mines, quarries, and reservoirs; ideas from the Florida Critical Lands and Waters Identification Project's approach for evaluating land use intensity; and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- Very natural landscape
- Natural landscape
- Mostly natural landscape
- Partly natural landscape
- Altered landscape
- Heavily altered landscape

Table 12: Indicator values for landscape condition within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Very natural landscape	1,149,577	4.4%	
	Natural landscape	5,929,104	22.9%	
	Mostly natural landscape	7,640,311	29.5%	↑ In good condition
	Partly natural landscape	10,215,079	39.5%	↓ Not in good condition
	Altered landscape	788,387	3.0%	
↓ Low	Heavily altered landscape	134,755	0.5%	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Mississippi Alluvial Valley forest birds - protection

This indicator prioritizes areas for new land protection within the Mississippi Alluvial Valley (MAV) based on benefits to forest breeding birds that need large interior cores of bottomland hardwood habitat (Swainson's warbler, cerulean warbler, swallow-tailed kite). The model considers core size, the amount of existing protected land within a forest patch, proximity to reforestation priorities, and risk of conversion to agriculture based on flooding frequency. The highest scores go to drier, unprotected forest patches with cores at least 2,000 ha (~5,000 ac) in size that are adjacent to complementary reforestation priority areas also identified by the Lower Mississippi Valley Joint Venture (LMVJV). This indicator originates from the LMVJV MAV forest breeding bird protection priorities.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Priority of forest breeding bird habitat patch for future protection

- Score >90-100 (highest priority)
- Score >80-90
- Score >70-80
- Score >60-70
- Score >50-60
- Score >40-50
- Score >30-40
- Score >20-30
- Score >10-20
- Score >0-10 (low priority)
- Score 0 (not a priority)

Table 13: Indicator values for Mississippi Alluvial Valley forest birds - protection within Kentucky. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Priority of forest breeding bird habitat patch for future protection	Acres	Percent of Area
↑ High	Score >90-100 (highest priority)	4,325	<0.1%
	Score >80-90	0	0%
	Score >70-80	3,640	<0.1%
	Score >60-70	3,377	<0.1%
	Score >50-60	3,861	<0.1%
	Score >40-50	0	0%
	Score >30-40	4,122	<0.1%
	Score >20-30	0	0%
	Score >10-20	1,955	<0.1%
	Score >0-10 (low priority)	46	<0.1%
↓ Low	Score 0 (not a priority)	128,989	0.5%
	Area not evaluated for this indicator	*25,706,900*	*99.4%*
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Mississippi Alluvial Valley forest birds - reforestation

This indicator prioritizes areas for reforestation within the Mississippi Alluvial Valley (MAV) based on benefits to three species of forest breeding birds that need large interior cores of bottomland hardwood habitat (Swainson's warbler, cerulean warbler, swallow-tailed kite). The model considers the core size, number of cores, and percent of local forest cover that would result from reforestation, as well as risk of conversion to agriculture based on flooding frequency. The highest scores go to drier areas where reforestation would create new forest patches containing interior cores at least 2,000 ha (~5,000 ac) in size. It originates from the Lower Mississippi Valley Joint Venture MAV forest breeding bird reforestation priorities.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Likelihood that reforestation will contribute to forest breeding bird habitat needs

- Most likely (≥90th percentile)
- Most likely (80th to <90th percentile)
- More likely (70th to <80th percentile)
- Less likely (60th to <70th percentile)
- Least likely (50th to <60th percentile)
- Least likely (40th to <50th percentile)
- Least likely (30th to <40th percentile)
- Least likely (20th to <30th percentile)
- Least likely (10th to <20th percentile)
- Least likely (<10th percentile)
- Not a priority for reforestation

Table 14: Indicator values for Mississippi Alluvial Valley forest birds - reforestation within Kentucky. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Likelihood that reforestation will contribute to forest breeding bird habitat needs	Acres	Percent of Area
↑ High	Most likely (≥90th percentile)	15,272	<0.1%
	Most likely (80th to <90th percentile)	25,760	<0.1%
	More likely (70th to <80th percentile)	12,031	<0.1%
	Less likely (60th to <70th percentile)	3,885	<0.1%
	Least likely (50th to <60th percentile)	9,778	<0.1%
	Least likely (40th to <50th percentile)	1,754	<0.1%
	Least likely (30th to <40th percentile)	125	<0.1%
	Least likely (20th to <30th percentile)	0	0%
	Least likely (10th to <20th percentile)	0	0%
	Least likely (<10th percentile)	0	0%
↓ Low	Not a priority for reforestation	81,561	0.3%
	Area not evaluated for this indicator	*25,707,049*	*99.4%*
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Potential access to parks

This cultural resource indicator prioritizes places to create new parks that would fill gaps in access to open space within socially vulnerable communities in urban areas. It identifies areas where residents currently lack access to parks within a 10-minute walk (accounting for walkable road networks and access barriers like highways and fences), then prioritizes based on park need using demographic and environmental metrics. Parks help improve public health, foster a conservation ethic by providing opportunities for people to connect with nature, and support critical ecosystem services. This indicator originates from the Trust for Public Land's ParkServe park priority areas and the Center for Disease Control's Social Vulnerability Index.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Priority for a new park to serve people who lack nearby park access

- Very high priority
- High priority
- Moderate priority
- Not identified as a priority (within urban areas)

Table 15: Indicator values for potential access to parks within Kentucky. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Priority for a new park to serve people who lack nearby park access	Acres	Percent of Area
↑ High	Very high priority	42,222	0.2%
	High priority	53,027	0.2%
	Moderate priority	43,455	0.2%
↓ Low	Not identified as a priority (within urban areas)	25,718,509	99.5%
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Resilient terrestrial sites

This indicator depicts an area's capacity to maintain species diversity and ecosystem function in the face of climate change. It measures two factors that influence resilience. The first, landscape diversity, reflects the number of microhabitats and climatic gradients created by topography, elevation, and hydrology. The second, local connectedness, reflects the degree of habitat fragmentation and strength of barriers to species movement. Highly resilient sites contain many different habitat niches that support biodiversity, and allow species to move freely through the landscape to find suitable microclimates as the climate changes. This indicator originates from The Nature Conservancy's Resilient Land data.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- Most resilient
- More resilient
- Slightly more resilient
- Average/median resilience
- Slightly less resilient
- Less resilient
- Least resilient
- Developed

Table 16: Indicator values for resilient terrestrial sites within Kentucky. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Most resilient	554,683	2.1%
	More resilient	7,099,962	27.5%
	Slightly more resilient	4,943,250	19.1%
	Average/median resilience	5,019,082	19.4%
	Slightly less resilient	2,548,304	9.9%
	Less resilient	2,266,902	8.8%
	Least resilient	239,992	0.9%
↓ Low	Developed	2,757,712	10.7%
	Area not evaluated for this indicator	*427,327*	*1.7%*
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


River cane restoration

This indicator represents priority places for river cane restoration and co-stewardship near the lands of federally recognized Tribes. River cane is a bamboo species native to the Southeast. Historically, it was abundant and widespread, forming dense patches called canebrakes that could stretch for miles within the floodplain. Today, it has been reduced to less than 2% of its former extent. River cane not only provides significant habitat value and ecosystem services, but many Southeastern Native American Tribes consider it a cultural keystone species important to the continuity of traditional lifeways. This indicator combines data from multiple sources, including Natural Resources Conservation Service soils data, the Environmental Protection Agency's estimated floodplain, lands of federally recognized Tribes lands from the U.S. Census Bureau, and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- Potential river cane restoration/management area on Tribal land
- Potential river cane restoration/management area on protected land near Tribal lands
- Potential river cane restoration/management area near Tribal lands
- Potential river cane restoration/management area on protected land
- Potential river cane restoration/management area
- Not identified as a rivercane restoration/management area

Table 17: Indicator values for river cane restoration within Kentucky. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Potential river cane restoration/management area on Tribal land	0	0%
	Potential river cane restoration/management area on protected land near Tribal lands	920	<0.1%
	Potential river cane restoration/management area near Tribal lands	2,707	<0.1%
	Potential river cane restoration/management area on protected land	213,140	0.8%
	Potential river cane restoration/management area	3,006,541	11.6%
↓ Low	Not identified as a rivercane restoration/management area	22,633,905	87.5%
	Area not evaluated for this indicator	*0.22*	*<0.1%*
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Urban park size

This cultural resource indicator measures the size of parks larger than 5 acres in the urban environment. Protected natural areas in urban environments provide urban residents a nearby place to connect with nature, and offer refugia for some species. This indicator complements the equitable access to potential parks indicator by capturing the value of existing parks. It originates from the Protected Areas Database of the United States, U.S. Census Bureau urban areas, and the National Land Cover Database.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- 75+ acre urban park
- 50 to <75 acre urban park
- 30 to <50 acre urban park
- 10 to <30 acre urban park
- 5 to <10 acre urban park
- <5 acre urban park
- Not identified as an urban park

Table 18: Indicator values for urban park size within Kentucky. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	75+ acre urban park	42,966	0.2%
	50 to <75 acre urban park	2,424	<0.1%
	30 to <50 acre urban park	3,648	<0.1%
	10 to <30 acre urban park	4,750	<0.1%
	5 to <10 acre urban park	1,097	<0.1%
	<5 acre urban park	1,346	<0.1%
↓ Low	Not identified as an urban park	25,800,982	99.8%
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Floodplain inundation

This indicator uses the frequency of inundation to measure the hydrologic function of the floodplain. In a natural floodplain, water can move freely from mainstem rivers and lakes onto different parts of the floodplain. This cycle of frequent, but not persistent, flooding supports many aquatic species and habitats, like fish, waterfowl and swamp forests. It also provides essential ecosystem services such as nutrient retention and downstream flood control. When parts of the floodplain stay constantly wet or dry, this may signal negative impacts from structures like dikes, dams, and levees, which disconnect waterbodies from their adjacent floodplains. This indicator is based on Sentinel-2 satellite imagery and methods developed by Yvonne Allen of the U.S. Fish and Wildlife Service.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Frequency of inundation within the floodplain

- Frequent inundation (flooded in 21-90% of days with available data)
- Regular inundation (flooded in 6-20% of days with available data)
- Occasional inundation (flooded in 2-5% of days with available data)
- Persistent inundation (flooded in 91-100% of days with available data)
- No detected inundation (flooded in 0% of days with available data)
- Not identified as a floodplain

Table 19: Indicator values for floodplain inundation within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: Frequency of inundation within the floodplain	Acres	Percent of Area	
↑ High	Frequent inundation (flooded in 21-90% of days with available data)	136,957	0.5%	
	Regular inundation (flooded in 6-20% of days with available data)	253,266	1.0%	↑ In good condition
	Occasional inundation (flooded in 2-5% of days with available data)	267,311	1.0%	↓ Not in good condition
	Persistent inundation (flooded in 91-100% of days with available data)	325,287	1.3%	
	No detected inundation (flooded in 0% of days with available data)	2,071,107	8.0%	
↓ Low	Not identified as a floodplain	22,803,286	88.2%	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Imperiled aquatic species

This indicator measures the number of aquatic animal Regional Species of Greatest Conservation Need (RSGCN) observed within each 12-digit HUC subwatershed, including fish, mussels, snails, crayfish, and amphibians. It captures patterns of rare and endemic aquatic species diversity. RSGCN are regional priority species derived from the list of SGCN identified in Southeast State Wildlife Action Plans as most in need of need of conservation action. RSGCN were chosen based on consistent criteria, such as level of conservation concern, regional stewardship responsibility, and ecological significance. This indicator originates from state Natural Heritage Program data collected by the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Number of aquatic animal Regional Species of Greatest Conservation Need (RSGCN) observed

- 8+ species
- 7 species
- 6 species
- 5 species
- 4 species
- 3 species
- 2 species
- 1 species
- 0 species
- Not identified as a floodplain

Table 20: Indicator values for imperiled aquatic species within Kentucky. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Number of aquatic animal Regional Species of Greatest Conservation Need (RSGCN) observed	Acres	Percent of Area
↑ High	8+ species	347,007	1.3%
	7 species	44,341	0.2%
	6 species	49,592	0.2%
	5 species	101,720	0.4%
	4 species	96,784	0.4%
	3 species	214,446	0.8%
	2 species	296,811	1.1%
	1 species	512,396	2.0%
	0 species	1,390,829	5.4%
↓ Low	Not identified as a floodplain	22,803,286	88.2%
	Area not evaluated for this indicator	*0.22*	*<0.1%*
	Total area	**25,857,213**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Lakes & reservoirs

This indicator assesses the condition of lakes and reservoirs based on the amount of natural landcover within the upstream watershed. Higher scores go to lakes and reservoirs with less surrounding urban and agricultural development (including intensive forest management for timber production), which negatively impact drinking water quality and quantity and fish habitat. This indicator uses waterbody and watershed data from the LAGOS-LOCUS dataset, as well as landcover from the National Land Cover Database and LANDFIRE.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



- Lake or reservoir with low disturbance (0-25%) in upstream watershed
- Lake or reservoir with medium disturbance (>25-60%) in upstream watershed
- Lake or reservoir with high disturbance (>60%) in upstream watershed
- Not identified as a lake or reservoir

Table 21: Indicator values for lakes & reservoirs within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Lake or reservoir with low disturbance (0-25%) in upstream watershed	33,848	0.1%	↑ In good condition
	Lake or reservoir with medium disturbance (>25-60%) in upstream watershed	187,294	0.7%	↓ Not in good condition
	Lake or reservoir with high disturbance (>60%) in upstream watershed	27,685	0.1%	
↓ Low	Not identified as a lake or reservoir	25,608,387	99.0%	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Natural landcover in floodplains

This indicator measures the amount of natural landcover in the estimated floodplain of rivers and streams within each catchment. It assesses the stream channel and its surrounding riparian buffer, measuring the percent of unaltered habitat like forests, wetlands, or open water (rather than agriculture or development). Intact vegetated buffers within the floodplain of rivers and streams provide aquatic habitat, improve water quality, reduce erosion and flooding, recharge groundwater, and more. This indicator originates from the National Land Cover Database and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Percent natural landcover within the estimated floodplain, by catchment

- >90% natural landcover
- >80-90% natural landcover
- >70-80% natural landcover
- >60-70% natural landcover
- ≤60% natural landcover
- Not identified as a floodplain

Table 22: Indicator values for natural landcover in floodplains within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: Percent natural landcover within the estimated floodplain, by catchment	Acres	Percent of Area	
↑ High	>90% natural landcover	544,856	2.1%	
	>80-90% natural landcover	222,945	0.9%	↑ In good condition
	>70-80% natural landcover	222,438	0.9%	↓ Not in good condition
	>60-70% natural landcover	230,343	0.9%	
	≤60% natural landcover	1,833,344	7.1%	
↓ Low	Not identified as a floodplain	22,803,286	88.2%	
	Area not evaluated for this indicator	*0.22*	*<0.1%*	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Network complexity

This indicator depicts the number of connected stream size classes in a river network between dams or waterfalls. River networks with a variety of connected stream classes help retain aquatic biodiversity in a changing climate by allowing species to access climate refugia and move between habitats. This indicator originates from the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Number of connected stream size classes

- 7 size classes
- 6 size classes
- 5 size classes
- 4 size classes
- 3 size classes
- 2 size classes
- 1 size class
- Not identified as a floodplain

Table 23: Indicator values for network complexity within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: Number of connected stream size classes	Acres	Percent of Area	
↑ High	7 size classes	591,956	2.3%	
	6 size classes	1,169,870	4.5%	
	5 size classes	604,775	2.3%	
	4 size classes	375,437	1.5%	↑ In good condition
	3 size classes	110,943	0.4%	↓ Not in good condition
	2 size classes	114,233	0.4%	
	1 size class	79,699	0.3%	
↓ Low	Not identified as a floodplain	22,810,301	88.2%	
	Area not evaluated for this indicator	*0.22*	*<0.1%*	
	Total area	**25,857,213**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Permeable surface

This indicator measures the average percent of non-impervious cover within each catchment. High levels of impervious surface degrade water quality and alter freshwater flow, impacting both aquatic species communities and ecosystem services for people, like the availability of clean drinking water. This indicator originates from the National Land Cover Database.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Percent of catchment permeable

- >95% permeable (likely high water quality and supporting most sensitive aquatic species)
- >90-95% permeable (likely declining water quality and supporting most aquatic species)
- >70-90% permeable (likely degraded water quality and not supporting many aquatic species)
- ≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)

Table 24: Indicator values for permeable surface within Kentucky. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Percent of catchment permeable	Acres	Percent of Area	
↑ High	>95% permeable (likely high water quality and supporting most sensitive aquatic species)	23,763,319	91.9%	↑ In good condition
	>90-95% permeable (likely declining water quality and supporting most aquatic species)	987,128	3.8%	↓ Not in good condition
	>70-90% permeable (likely degraded water quality and not supporting many aquatic species)	848,718	3.3%	
↓ Low	≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)	258,048	1.0%	
	Area not evaluated for this indicator	*0.22*	*<0.1%*	
	Total area	**25,857,213**	**100%**	


To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

More Information

Priority Amphibian and Reptile Conservation Areas

Priority Amphibian and Reptile Conservation Areas (PARCAs) are an expert-driven, nonregulatory designation that includes places capable of supporting viable amphibian and reptile populations, places occupied by rare or imperiled species, and places rich in biodiversity or species unique to that geographic area. Reptiles and amphibians are a critical part of the Southeast region's rich biodiversity and many populations are declining in the face of threats like habitat loss, invasive species, and climate change. The PARCA dataset is maintained by the Amphibian and Reptile Conservancy and does not yet include Virginia or Kentucky.





Priority Amphibian and Reptile Conservation Area (PARCA)

Not a PARCA (excluding Kentucky and Virginia)

Table 25: Extent of Priority Amphibian and Reptile Conservation Areas within Kentucky.

Priority Amphibian and Reptile Conservation Areas status	Acres	Percent of Area
Priority Amphibian and Reptile Conservation Area (PARCA)	809	<0.1%
Not a PARCA (excluding Kentucky and Virginia)	25,856,404	100.0%
Total area	**25,857,213**	**100%**

Priority Amphibian and Reptile Conservation Areas at this location:

Cumberland Mountains

The Cumberland Mountains PARCA occurs within the Central Appalachians ecoregion and contains the highest elevations of the Cumberland Plateau and some of largest tracts of contiguous forest in Tennessee. Much of this PARCA is mountainous and much more dissected than the adjacent Cumberland Plateau, with elevations ranging from 360-1,060 m with local relief up to 600 m. It’s located primarily on publicly owned land, including North Cumberland Wildlife Management Area, Tackett Creek Wildlife Management Area, Frozen Head State Park, and Indian Mountain State Park. It contains a very rich diversity of herpetofauna, with a big portion of that diversity consisting of salamanders. One is the Wehrle's salamander, with a subspecies population only occurring on the Cumberland Plateau. Unfortunately, a lot of primary habitat has been disturbed and negatively impacted by coal mining, feral hogs, and incompatible forestry practices. However, recent conservation focus and effort have been aimed toward mitigating the impacts of mining and conducting inventory and monitoring of the Wehrle's salamander.

Sand Prairie Bottoms

The Sand Prairie Bottoms PARCA is situated along the Mississippi Alluvial Valley and encompasses the meander belt of the Mississippi River. Point bars, oxbows, natural levees, and abandoned channels are all components of this ecosystem. This area was once extremely diverse, but due to draining of wetlands, channelizing of streams, and clearing of bottomland forests for agriculture, much of the land has been severely altered and tarnished.

Wayne

The Wayne PARCA, encompassing almost all of Wayne County, West Virginia, is predominantly characterized by rolling hills and river valleys, with the Big Sandy River and its tributaries like Twelvepole Creek weaving through its terrain. These waterways not only provide vital habitats for aquatic species, but also contribute to the PARCA’s overall biodiversity. Wayne County is heavily forested, primarily with hardwood species such as oak, hickory, maple, and poplar, which support a variety of wildlife, including the eastern hognose snake, a rare species found in the region’s diverse habitats.


Western Tennessee River Valley

The Western Tennessee River Valley PARCA is an ecologically diverse region characterized by its rich alluvial soils and a mix of hardwood forests, wetlands, and riverine habitats. This area, which stretches along the Tennessee River, supports a variety of plant and animal life adapted to its moist, fertile environment. The floodplains and riverbanks are home to lush bottomland forests with species such as oak, cypress, and tupelo trees, while the wetlands host a range of amphibians, fish, and migratory birds. This valley is also critical for maintaining water quality and providing essential ecosystem services, such as flood regulation and nutrient cycling.

Names and descriptions of public Priority Amphibian and Reptile Areas were provided by the Amphibian and Reptile Conservancy on August 30, 2024 and edited slightly for clarity and consistency.


Protected Areas

Protected areas include a variety of public lands owned or managed by federal, state, and local agencies and nonprofit organizations, as well as some private conservation lands and conservation easements. Protected areas are derived from the Protected Areas Database of the United States (PAD-US v4.1) and include Fee, Designation, Easement, Marine, and Proclamation (Dept. of Defense lands only) boundaries.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Within a protected area

Not within a protected area

Table 26: Extent of protected areas within Kentucky.

Protected area status	Acres	Percent of Area
Not within a protected area	23,688,369	91.6%
Within a protected area	2,168,844	8.4%
Total area	**25,857,213**	**100%**

Protected areas at this location:

- Daniel Boone National Forest (USDA Forest Service; 563,865 acres)
- Redbird Purchase Unit (USDA Forest Service; 146,434 acres)
- Fort Knox (109,089 acres)
- Land Between The Lakes National Recreation Area (107,587 acres)
- Land Between the Lakes Other (USDA Forest Service; 107,548 acres)
- Cumberland Recreation Area (90,149 acres)
- Corrigan Wildlife Mangement Area (Molpus; 54,818 acres)
- Barkley Recreation Area (52,513 acres)
- Lake Cumberland Wildlife Management Area (US Army Corps of Engineers; 51,668 acres)
- Mammoth Cave National Park (National Park Service; 50,624 acres)
- Lake Cumberland (46,069 acres)
- Peabody Wildlife Management Area (Kentucky Department of Fish and Wildlife Resources & Western Land Services & Patriot Coal Corporation; 41,327 acres)
- Fort Campbell (37,150 acres)
- Lake Barkley (36,798 acres)
- Green Rv Recreation Area (32,392 acres)
- Boone Forestlands Wildlife Management Area (Begley Properties, LLC; 31,666 acres)
- Big South Fork National River and Recreation Area (National Park Service; 30,307 acres)
- Bluegrass Conservancy Easements (Private Individual & Bluegrass Land Conservancy; 28,357 acres)
- Redbird Wildlife Management Area (U.S. Forest Service; 24,015 acres)
- Green River Lake Wildlife Management Area (US Army Corps of Engineers; 21,039 acres)
- Barren River Recreation Area (20,297 acres)
- Yatesville Recreation Area (20,111 acres)
- Beaver Creek Wildlife Management Area (U.S. Forest Service; 17,751 acres)
- Elk Forest Wildlife Management Area (Private Individual; 16,801 acres)
- Bernheim Arboretum and Research Forest (Bernheim Arboretum and Research Forest; 16,164 acres)
- ... and 2,439 more protected areas ...

Note: areas are listed based on name, ownership, and boundary information in the Protected Areas Database of the United States, which may include overlapping and duplicate areas. Only protected areas with at least 1 acre of overlap with this area are included in the list above.


Sea Level Rise

Sea level rise unlikely to be a threat (inland counties).

Urban Growth


The FUTURES urban growth model predicts the likelihood that an area will urbanize at every decade from 2020 to 2100. Developed areas from the 2021 National Landcover Database serve as the baseline for current urban areas. The model simulates landscape change based on trends in population growth, local development suitability factors, and an urban patch-growing algorithm. It considers environmental drivers like distance to floodplain, slope, and available infrastructure, and even socio-economic status. The probability of urbanization for each area reflects how many times it urbanized out of 50 model runs.

To explore maps for additional time periods, click here.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Probability of urbanization by 2060

- Urban in 2021
- Very high likelihood of urbanization (>50% probability)
- High likelihood of urbanization (25 - 50% probability)
- Moderate likelihood of urbanization (2 - 25% probability)
- Not likely to urbanize

Table 27: Extent of projected urbanization by decade within Kentucky. Values from FUTURES model projections for the contiguous United States developed by the Center for Geospatial Analytics, NC State University. 2060 corresponds to the SECAS goal: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.


Decade	Acres	Percent of Area
Urban in 2021	2,135,970	8.3%
2030 projected extent	2,187,745	8.5%
2040 projected extent	2,216,985	8.6%
2050 projected extent	2,239,416	8.7%
2060 projected extent	2,260,626	8.7%
2070 projected extent	2,280,093	8.8%
2080 projected extent	2,295,552	8.9%
2090 projected extent	2,305,825	8.9%
2100 projected extent	2,312,583	8.9%
Not projected to urbanize by 2100	*23,544,630*	*91.1%*
Total area	**25,857,213**	**100%**

8.3% of this area is already urban in 2021, and an additional 9.4% has at least a moderate probability of urbanizing by 2060.

By 2060, the size of the urban footprint is projected to increase **5.8%** over 2021 levels.


Wildfire Likelihood

Wildfire likelihood data originate from the Wildfire Risk to Communities project developed by the U.S. Forest Service. This layer depicts the probability of wildfire burning in a specific location in any given year. Annual burn probabilities in the United States range from 0-14%, but do not exceed 8% in the Southeast. Wildfire likelihood is based on fire behavior modeling across thousands of simulations of possible fire seasons. In each simulation, factors contributing to the probability of a fire occurring (such as weather, topography, and ignitions) vary based on patterns derived from observations in recent decades. Wildfire likelihood is not predictive and does not reflect any forecasted future weather or fire danger conditions. It also does not say anything about the intensity of fire if it occurs. To explore additional wildfire risk information, please see the [Wildfire Risk to Communities] website.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

48 96 192 miles



Wildfire likelihood (annual burn probability)

- High
- Moderate-high
- Moderate
- Low-moderate
- Low
- Not predicted to experience wildfire

Table 28: Area in each wildfire probability category within Kentucky. Values from the Wildire Risk To Communities project developed by the USDA Forest Service.

Wildfire likelihood (annual burn probability)	Acres	Percent of Area
Not predicted to experience wildfire (0% probability)	2,166,771	8.4%
Low (>0 - 0.01% probability)	8,735,975	33.8%
Low-moderate (>0.01 - 0.02154% probability)	4,728,494	18.3%
Low-moderate (>0.02154 - 0.04642% probability)	2,713,270	10.5%
Moderate (>0.04642 - 0.1% probability)	1,540,076	6.0%
Moderate (>0.1 - 0.21544% probability)	1,177,272	4.6%
Moderate (>0.21544 - 0.46416% probability)	1,778,051	6.9%
Moderate-high (>0.46416 - 1% probability)	2,756,692	10.7%
Moderate-high (>1 - 2.15443% probability)	259,668	1.0%
High (>2.15443 - 4.64159% probability)	849	<0.1%
High (>4.64159% probability)	0	0%
No wildfire risk data available	*95*	*<0.1%*
Total area	**25,857,213**	**100%**