Southeast Conservation Blueprint Summary

for Mississippi

Created 10/21/2026

Table of Contents

About the Southeast Blueprint 3

Southeast Blueprint Priorities 4

Hubs and Corridors 6

Indicator Summary 8

More Information 74

The Southeast Conservation Blueprint 2026

Southeast Conservation Adaptation Strategy

[THIS PAGE INTENTIONALLY LEFT BLANK]

About the Southeast Blueprint

The Southeast Conservation Blueprint is the primary product of the Southeast Conservation Adaptation Strategy (SECAS). It is a living, spatial plan to achieve the SECAS vision of a connected network of lands and waters across the Southeast and Caribbean. The Blueprint is regularly updated to incorporate new data, partner input, and information about on-the-ground conditions.

The Blueprint identifies priority areas based on a suite of natural and cultural resource indicators representing terrestrial, freshwater, and marine ecosystems. A connectivity analysis identifies corridors that link coastal and inland areas and span climate gradients.

For more information:

- Visit the Blueprint webpage
- Review the Blueprint 2026 Development Process
- View and download the Blueprint data and make maps on the Blueprint page of the SECAS Atlas

We're here to help!


- Do you have a question about the Blueprint?
- Would you like help using the Blueprint to support a proposal or inform a decision?
- Do you have a suggestion on how to improve the Blueprint? The Blueprint and its inputs are regularly revised based on input from people like you.
- Do you have feedback on how to improve the Blueprint Explorer interface?

If you need help or have questions, contact Southeast Blueprint staff by reaching out to a member of the user support team.

We're here to support you. We really mean it. It's what we do!


This report was generated by the Southeast Conservation Blueprint Explorer, which was developed by Astute Spruce, LLC in partnership with the U.S. Fish and Wildlife Service under the Southeast Conservation Adaptation Strategy.

Southeast Blueprint Priorities



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Priorities for a connected network of lands and waters

- Highest priority
- High priority
- Medium priority
- Priority connections

Priority Categories

For a connected network of lands and waters

In total, Blueprint priorities and priority connections cover roughly 50% of the Southeast Blueprint geography.

Highest priority

Areas where conservation action would make the biggest impact, based on a suite of natural and cultural resource indicators. This class covers roughly 10% of the Southeast Blueprint geography.

High priority

Areas where conservation action would make a big impact, based on a suite of natural and cultural resource indicators. This class covers roughly 15% of the Southeast Blueprint geography.

Medium priority

Areas where conservation action would make an above-average impact, based on a suite of natural and cultural resource indicators. This class covers roughly 20% of the Southeast Blueprint geography.

Priority connections

Connections between priority areas that cover the shortest distance possible while routing through as much Blueprint priority as possible. This class covers roughly 5% of the Southeast Blueprint geography.


Table 1: Extent of each Blueprint priority category within Mississippi.

Priority Category	Acres	Percent of Area
Highest priority	1,961,417	6.3%
High priority	2,953,328	9.5%
Medium priority	5,400,001	17.4%
Priority connections	2,162,517	7.0%
Lower priority	18,521,620	59.7%
Total area	**30,998,884**	**100%**

Hubs and Corridors


The Blueprint uses a least-cost path connectivity analysis to identify corridors that link hubs across the shortest distance possible, while also routing through as much Blueprint priority as possible.

In the continental Southeast, hubs are large patches (~5,000+ acres) of highest priority Blueprint areas and/or protected lands.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Hubs
Corridors

Table 2: Extent of hubs and corridors within Mississippi.

Type	Acres	Percent of Area
Hubs	2,760,085	8.9%
Corridors	5,327,730	17.2%
Not a hub or corridor	22,911,070	73.9%
Total area	**30,998,884**	**100%**

Indicator Summary

Table 3: Terrestrial indicators.

Indicator	Present
East Coastal Plain open pine birds	✓
Fire frequency	✓
Grassland & savanna extent	✓
Grassland & savanna restoration	✓
Greenways & trails	✓
Imperiled amphibians & reptiles	✓
Imperiled mammals	✓
Intact habitat cores	✓
Landscape condition	✓
Mississippi Alluvial Valley forest birds - protection	✓
Mississippi Alluvial Valley forest birds - reforestation	✓
Potential access to parks	✓
Resilient terrestrial sites	✓
River cane restoration	✓
South Atlantic forest birds	-
South Atlantic low-urban historic landscapes	-
Urban park size	✓
West Coastal Plain & Ouachitas forested wetland birds	-
West Coastal Plain & Ouachitas open pine birds	-
West Gulf Coast mottled duck nesting	-

Table 4: Freshwater indicators.

Indicator	Present
Floodplain inundation	✓
Gulf migratory fish connectivity	✓
Imperiled aquatic species	✓
Lakes & reservoirs	✓
Natural landcover in floodplains	✓
Network complexity	✓
Permeable surface	✓


Table 5: Coastal & marine indicators.

Indicator	Present
Coastal shoreline condition	✓
Estuarine coastal condition	✓
Gulf coral & hardbottom	✓
Gulf deep-sea coral richness	-
Gulf marine birds	✓
Gulf marine mammals	✓
Gulf sea turtles	✓
Island habitat	✓
Marine highly migratory fish	-
Resilient coastal sites	✓
Seagrass	✓
South Atlantic maritime forest	-
Stable coastal wetlands	✓

Terrestrial


East Coastal Plain open pine birds

This indicator identifies areas within the historical longleaf pine range east of the Mississippi River where creating or maintaining open pine habitat would most benefit six focal species of birds (Bachman's sparrow, red-cockaded woodpecker, Henslow's sparrow, red-headed woodpecker, Northern bobwhite, brown-headed nuthatch). It prioritizes areas for open pine conservation based on suitability for longleaf pine, feasibility of prescribed burning, proximity to protected lands, habitat suitability for focal bird species, and proximity to bird source populations. It originates from the East Gulf Coastal Plain Joint Venture's prioritization of areas for open pine ecosystem restoration.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Priority for open pine conservation for focal bird species

- High priority (score >80-100)
- Medium-high priority (score >60-80)
- Medium priority (score >40-60)
- Medium-low priority (score >20-40)
- Low priority (score 0-20)
- Not a priority (not identified as upland pine)

Table 6: Indicator values for East Coastal Plain open pine birds within Mississippi. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Priority for open pine conservation for focal bird species	Acres	Percent of Area
↑ High	High priority (score >80-100)	7,318	<0.1%
	Medium-high priority (score >60-80)	423,081	1.4%
	Medium priority (score >40-60)	1,140,979	3.7%
	Medium-low priority (score >20-40)	3,694,783	11.9%
	Low priority (score 0-20)	485,691	1.6%
↓ Low	Not a priority (not identified as upland pine)	6,959,944	22.5%
	Area not evaluated for this indicator	*18,287,089*	*59.0%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Fire frequency

This indicator uses remote sensing, fire treatment data, and historical landcover models to assess whether an area matches its historical fire regime now and in the recent past (1984-2024). It focuses on fire-adapted forests and grasslands that historically burned at least once every 20 years, which rely on regular, low-intensity fires to maintain habitat, support native plant growth, and reduce wildfire risk. It evaluates departure from historical conditions based on the U.S. Forest Service's fire regime condition classes. This indicator combines Landsat-based fire history metrics from the U.S. Geological Survey, fire treatment data from LANDFIRE, and historical fire regime estimates from LANDFIRE. Development of fire history metrics was supported by Southeast FireMap.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



How well fire-adapted areas match their historic fire regime under current conditions vs. recent past (1984-2024)

- Currently in historical fire regime, low (<33%) departure in recent past
- Currently in historical fire regime, medium (33-66%) departure in recent past
- Currently in historical fire regime, high (>66%) departure in recent past
- Not currently in historical fire regime, low (<33%) departure in recent past
- Not currently in historical fire regime, medium (33-66%) departure in recent past
- Not currently in historical fire regime, high (>66%) departure in recent past
- Urban, row crop, or not a fire-adapted area (>20-year fire regime)

Table 7: Indicator values for fire frequency within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: How well fire-adapted areas match their historic fire regime under current conditions vs. recent past (1984-2024)	Acres	Percent of Area	
↑ High	Currently in historical fire regime, low (<33%) departure in recent past	79,653	0.3%	
	Currently in historical fire regime, medium (33-66%) departure in recent past	296,759	1.0%	↑ In good condition
	Currently in historical fire regime, high (>66%) departure in recent past	357,055	1.2%	↓ Not in good condition
	Not currently in historical fire regime, low (<33%) departure in recent past	8,825	<0.1%	
	Not currently in historical fire regime, medium (33-66%) departure in recent past	471,656	1.5%	
	Not currently in historical fire regime, high (>66%) departure in recent past	8,875,494	28.6%	
↓ Low	Urban, row crop, or not a fire-adapted area (>20-year fire regime)	20,903,514	67.4%	
	Area not evaluated for this indicator	*5,928*	*<0.1%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Grassland & savanna extent

This indicator represents grasslands and savannas in the southeastern United States, which support important plants, reptiles, amphibians, mammals, birds, and pollinators. It considers known grassland and savanna locations, predicted locations within protected areas, and surrounding pollinator buffers. It also incorporates other potential grassland and savanna locations within natural and altered landscapes, and restoration opportunities within historical locations based on past fire intervals and historic ecosystem predictions. This indicator combines data from multiple sources, including the Southeastern Grasslands Institute, the National Land Cover Database, LANDFIRE, Oklahoma and Texas ecological systems maps, and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Known grassland/savanna
- Likely grassland/savanna >10 acres
- Likely grassland/savanna ≤10 acres
- Pollinator buffer around known or likely grassland/savanna
- Potential grassland/savanna in grassland/savanna hub
- Potential grassland/savanna outside grassland/savanna hub
- Historic grassland/savanna
- Not identified as grassland/savanna

Table 8: Indicator values for grassland & savanna extent within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Known grassland/savanna	3,194	<0.1%	
	Likely grassland/savanna >10 acres	33,719	0.1%	
	Likely grassland/savanna ≤10 acres	19,737	<0.1%	↑ In good condition
	Pollinator buffer around known or likely grassland/savanna	204,950	0.7%	↓ Not in good condition
	Potential grassland/savanna in grassland/savanna hub	418,108	1.3%	
	Potential grassland/savanna outside grassland/savanna hub	1,613,347	5.2%	
	Historic grassland/savanna	9,402,992	30.3%	
↓ Low	Not identified as grassland/savanna	19,302,483	62.3%	
	Area not evaluated for this indicator	*354*	*<0.1%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Grassland & savanna restoration

This indicator represents potential restoration opportunities for grasslands and savannas within their historical range in the southeastern United States. Grasslands support important plants, reptiles, amphibians, mammals, birds, and pollinators, but have significantly declined from their current extent. It considers proximity to clusters of existing grasslands and savannas, protected areas with management potential, landcover classes with good restoration potential, priority areas for restoration of longleaf and shortleaf pine, and historical grassland and savanna locations. This indicator combines data from multiple sources, including the Southeast Blueprint grassland and savanna extent indicator, the Longleaf Suitability Analysis, shortleaf pine suitability, landcover from the National Land Cover Database and LANDFIRE, and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

257 miles
64 128



Restoration priority

- Highest (in a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)
- Very high (in a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)
- High (outside a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)
- Medium (outside a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)
- Low (other historic grassland/savanna)
- Very low (already known or likely grassland/savanna)
- Lowest (not identified as historic grassland/savanna)

Table 9: Indicator values for grassland & savanna restoration within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Restoration priority	Acres	Percent of Area
↑ High	Highest (in a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)	647,876	2.1%
	Very high (in a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)	1,327,860	4.3%
	High (outside a grassland/savanna hub, in a protected area with management potential, in a historic patch >10 acres)	133,588	0.4%
	Medium (outside a grassland/savanna hub, unprotected landcover with good restoration potential, in a historic patch >10 acres)	5,195,345	16.8%
	Low (other historic grassland/savanna)	4,248,344	13.7%
	Very low (already known or likely grassland/savanna)	56,651	0.2%
↓ Low	Lowest (not identified as historic grassland/savanna)	19,388,866	62.5%
	Area not evaluated for this indicator	*354*	*<0.1%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Greenways & trails

This cultural resource indicator measures both the natural condition and connected length of greenways and trails to characterize the quality of the recreational experience. Natural condition is based on the amount of impervious surface surrounding the path. Connected length captures how far a person can go without leaving a dedicated path, based on common distances for walking, running, and biking. This indicator originates from OpenStreetMap data and the National Land Cover Database.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

257 miles
64 128



- Mostly natural and connected for ≥40 km
- Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km
- Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km
- Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km
- Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km
- Developed and connected for <1.9 km
- Sidewalk
- Not identified as a trail, sidewalk, or other path

Table 10: Indicator values for greenways & trails within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Mostly natural and connected for ≥40 km	1,535	<0.1%	
	Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km	2,903	<0.1%	
	Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km	2,040	<0.1%	
	Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km	1,885	<0.1%	↑ In good condition
	Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km	1,584	<0.1%	↓ Not in good condition
	Developed and connected for <1.9 km	2,590	<0.1%	
	Sidewalk	4,091	<0.1%	
↓ Low	Not identified as a trail, sidewalk, or other path	30,975,979	99.9%	
	Area not evaluated for this indicator	*6,278*	*<0.1%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Imperiled amphibians & reptiles

This indicator identifies potential habitat to support amphibian and reptile Regional Species of Greatest Conservation Need (RSGCN). Using a rarity-weighted richness approach, it prioritizes places important for species that are restricted to smaller ranges, such as narrow endemics, which are therefore more vulnerable to habitat loss or change. RSGCN are regional priority species derived from the list of SGCN identified in Southeast State Wildlife Action Plans as most in need of conservation action. RSGCN were chosen based on consistent criteria, such as level of conservation concern, regional stewardship responsibility, and ecological significance. This indicator uses 190 species and subspecies models from the U.S. Geological Survey Gap Analysis Project that are based on known ranges, habitat relationships, and LANDFIRE landcover.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Importance for range-restricted species

- Highest importance (top 5%)
- Very high importance (upper 5-10%)
- High importance (upper 10-20%)
- Medium importance (above average)
- Low importance (below average)
- Developed landcover or no potential habitat

Table 11: Indicator values for imperiled amphibians & reptiles within Mississippi. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Importance for range-restricted species	Acres	Percent of Area
↑ High	Highest importance (top 5%)	1,758,510	5.7%
	Very high importance (upper 5-10%)	994,587	3.2%
	High importance (upper 10-20%)	1,797,384	5.8%
	Medium importance (above average)	3,837,240	12.4%
	Low importance (below average)	12,843,637	41.4%
↓ Low	Developed landcover or no potential habitat	9,308,583	30.0%
	Area not evaluated for this indicator	*458,943*	*1.5%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Imperiled mammals

This indicator identifies potential habitat to support mammal Regional Species of Greatest Conservation Need (RSGCN). Using a rarity-weighted richness approach, it prioritizes places important for species that are restricted to smaller ranges, such as narrow endemics, which are therefore more vulnerable to habitat loss or change. RSGCN are regional priority species derived from the list of SGCN identified in Southeast State Wildlife Action Plans as most in need of conservation action. RSGCN were chosen based on consistent criteria, such as level of conservation concern, regional stewardship responsibility, and ecological significance. This indicator uses 58 species models from the U.S. Geological Survey Gap Analysis Project that are based on known ranges, habitat relationships, and LANDFIRE landcover.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Importance for range-restricted species

- Highest importance (top 5%)
- Very high importance (upper 5-10%)
- High importance (upper 10-20%)
- Medium importance (above average)
- Low importance (below average)
- Developed landcover or no potential habitat

Table 12: Indicator values for imperiled mammals within Mississippi. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Importance for range-restricted species	Acres	Percent of Area
↑ High	Highest importance (top 5%)	0	0%
	Very high importance (upper 5-10%)	100,832	0.3%
	High importance (upper 10-20%)	4,268,069	13.8%
	Medium importance (above average)	10,819,585	34.9%
	Low importance (below average)	11,390,772	36.7%
↓ Low	Developed landcover or no potential habitat	3,960,683	12.8%
	Area not evaluated for this indicator	*458,943*	*1.5%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Intact habitat cores

This indicator represents the size of large, unfragmented patches of natural habitat. It identifies minimally disturbed natural areas at least 100 acres in size and greater than 200 meters wide. Large areas of intact natural habitat are important for many wildlife species, including reptiles and amphibians, birds, and large mammals. This indicator originates from Esri's green infrastructure data.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Large core (>10,000 acres)
- Medium core (>1,000-10,000 acres)
- Small core (>100-1,000 acres)
- Not a core

Table 13: Indicator values for intact habitat cores within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Large core (>10,000 acres)	2,093,458	6.8%	
	Medium core (>1,000-10,000 acres)	9,931,000	32.0%	
	Small core (>100-1,000 acres)	4,192,752	13.5%	↑ In good condition
↓ Low	Not a core	14,775,752	47.7%	↓ Not in good condition
	Area not evaluated for this indicator	*5,923*	*<0.1%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Landscape condition

This indicator represents natural areas with limited human alteration while also considering the naturalness of the surrounding landscape. Examples of human alteration include urban development and intense agricultural use. The degree of naturalness across the landscape is a key ecological condition for sustaining species and ecosystem services that are sensitive to habitat fragmentation at multiple scales. This indicator combines data from multiple sources, including the National Land Cover Database; various data on grasslands, mines, quarries, and reservoirs; ideas from the Florida Critical Lands and Waters Identification Project's approach for evaluating land use intensity; and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Very natural landscape
- Natural landscape
- Mostly natural landscape
- Partly natural landscape
- Altered landscape
- Heavily altered landscape

Table 14: Indicator values for landscape condition within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Very natural landscape	2,685,522	8.7%	
	Natural landscape	10,377,437	33.5%	
	Mostly natural landscape	8,196,520	26.4%	↑ In good condition
	Partly natural landscape	8,300,899	26.8%	↓ Not in good condition
	Altered landscape	893,024	2.9%	
↓ Low	Heavily altered landscape	86,539	0.3%	
	Area not evaluated for this indicator	*458,943*	*1.5%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Mississippi Alluvial Valley forest birds - protection

This indicator prioritizes areas for new land protection within the Mississippi Alluvial Valley (MAV) based on benefits to forest breeding birds that need large interior cores of bottomland hardwood habitat (Swainson's warbler, cerulean warbler, swallow-tailed kite). The model considers core size, the amount of existing protected land within a forest patch, proximity to reforestation priorities, and risk of conversion to agriculture based on flooding frequency. The highest scores go to drier, unprotected forest patches with cores at least 2,000 ha (~5,000 ac) in size that are adjacent to complementary reforestation priority areas also identified by the Lower Mississippi Valley Joint Venture (LMVJV). This indicator originates from the LMVJV MAV forest breeding bird protection priorities.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Priority of forest breeding bird habitat patch for future protection

- Score >90-100 (highest priority)
- Score >80-90
- Score >70-80
- Score >60-70
- Score >50-60
- Score >40-50
- Score >30-40
- Score >20-30
- Score >10-20
- Score >0-10 (low priority)
- Score 0 (not a priority)

Table 15: Indicator values for Mississippi Alluvial Valley forest birds - protection within Mississippi. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Priority of forest breeding bird habitat patch for future protection	Acres	Percent of Area
↑ High	Score >90-100 (highest priority)	44,895	0.1%
	Score >80-90	65,238	0.2%
	Score >70-80	103,794	0.3%
	Score >60-70	133,961	0.4%
	Score >50-60	71,142	0.2%
	Score >40-50	8,733	<0.1%
	Score >30-40	41,094	0.1%
	Score >20-30	3,667	<0.1%
	Score >10-20	54,654	0.2%
	Score >0-10 (low priority)	1,164	<0.1%
↓ Low	Score 0 (not a priority)	4,408,566	14.2%
	Area not evaluated for this indicator	*26,061,978*	*84.1%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Mississippi Alluvial Valley forest birds - reforestation

This indicator prioritizes areas for reforestation within the Mississippi Alluvial Valley (MAV) based on benefits to three species of forest breeding birds that need large interior cores of bottomland hardwood habitat (Swainson's warbler, cerulean warbler, swallow-tailed kite). The model considers the core size, number of cores, and percent of local forest cover that would result from reforestation, as well as risk of conversion to agriculture based on flooding frequency. The highest scores go to drier areas where reforestation would create new forest patches containing interior cores at least 2,000 ha (~5,000 ac) in size. It originates from the Lower Mississippi Valley Joint Venture MAV forest breeding bird reforestation priorities.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Likelihood that reforestation will contribute to forest breeding bird habitat needs

- Most likely (≥90th percentile)
- Most likely (80th to <90th percentile)
- More likely (70th to <80th percentile)
- Less likely (60th to <70th percentile)
- Least likely (50th to <60th percentile)
- Least likely (40th to <50th percentile)
- Least likely (30th to <40th percentile)
- Least likely (20th to <30th percentile)
- Least likely (10th to <20th percentile)
- Least likely (<10th percentile)
- Not a priority for reforestation

Table 16: Indicator values for Mississippi Alluvial Valley forest birds - reforestation within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Likelihood that reforestation will contribute to forest breeding bird habitat needs	Acres	Percent of Area
↑ High	Most likely (≥90th percentile)	289,659	0.9%
	Most likely (80th to <90th percentile)	345,365	1.1%
	More likely (70th to <80th percentile)	302,638	1.0%
	Less likely (60th to <70th percentile)	305,718	1.0%
	Least likely (50th to <60th percentile)	329,817	1.1%
	Least likely (40th to <50th percentile)	352,571	1.1%
	Least likely (30th to <40th percentile)	398,328	1.3%
	Least likely (20th to <30th percentile)	357,563	1.2%
	Least likely (10th to <20th percentile)	413,587	1.3%
	Least likely (<10th percentile)	153,461	0.5%
↓ Low	Not a priority for reforestation	1,688,948	5.4%
	Area not evaluated for this indicator	*26,061,228*	*84.1%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Potential access to parks

This cultural resource indicator prioritizes places to create new parks that would fill gaps in access to open space within socially vulnerable communities in urban areas. It identifies areas where residents currently lack access to parks within a 10-minute walk (accounting for walkable road networks and access barriers like highways and fences), then prioritizes based on park need using demographic and environmental metrics. Parks help improve public health, foster a conservation ethic by providing opportunities for people to connect with nature, and support critical ecosystem services. This indicator originates from the Trust for Public Land's ParkServe park priority areas and the Center for Disease Control's Social Vulnerability Index.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Priority for a new park to serve people who lack nearby park access

- Very high priority
- High priority
- Moderate priority
- Not identified as a priority (within urban areas)

Table 17: Indicator values for potential access to parks within Mississippi. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Priority for a new park to serve people who lack nearby park access	Acres	Percent of Area
↑ High	Very high priority	91,006	0.3%
	High priority	135,844	0.4%
	Moderate priority	144,608	0.5%
↓ Low	Not identified as a priority (within urban areas)	30,139,121	97.2%
	Area not evaluated for this indicator	*488,306*	*1.6%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


Resilient terrestrial sites

This indicator depicts an area's capacity to maintain species diversity and ecosystem function in the face of climate change. It measures two factors that influence resilience. The first, landscape diversity, reflects the number of microhabitats and climatic gradients created by topography, elevation, and hydrology. The second, local connectedness, reflects the degree of habitat fragmentation and strength of barriers to species movement. Highly resilient sites contain many different habitat niches that support biodiversity, and allow species to move freely through the landscape to find suitable microclimates as the climate changes. This indicator originates from The Nature Conservancy's Resilient Land data.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Most resilient
- More resilient
- Slightly more resilient
- Average/median resilience
- Slightly less resilient
- Less resilient
- Least resilient
- Developed

Table 18: Indicator values for resilient terrestrial sites within Mississippi. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Most resilient	717,016	2.3%
	More resilient	3,290,373	10.6%
	Slightly more resilient	4,389,221	14.2%
	Average/median resilience	11,293,704	36.4%
	Slightly less resilient	3,833,349	12.4%
	Less resilient	2,851,283	9.2%
	Least resilient	294,474	0.9%
↓ Low	Developed	2,730,013	8.8%
	Area not evaluated for this indicator	*1,599,451*	*5.2%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial


River cane restoration

This indicator represents priority places for river cane restoration and co-stewardship near the lands of federally recognized Tribes. River cane is a bamboo species native to the Southeast. Historically, it was abundant and widespread, forming dense patches called canebrakes that could stretch for miles within the floodplain. Today, it has been reduced to less than 2% of its former extent. River cane not only provides significant habitat value and ecosystem services, but many Southeastern Native American Tribes consider it a cultural keystone species important to the continuity of traditional lifeways. This indicator combines data from multiple sources, including Natural Resources Conservation Service soils data, the Environmental Protection Agency's estimated floodplain, lands of federally recognized Tribes lands from the U.S. Census Bureau, and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Potential river cane restoration/management area on Tribal land
- Potential river cane restoration/management area on protected land near Tribal lands
- Potential river cane restoration/management area near Tribal lands
- Potential river cane restoration/management area on protected land
- Potential river cane restoration/management area
- Not identified as a rivercane restoration/management area

Table 19: Indicator values for river cane restoration within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Potential river cane restoration/management area on Tribal land	7,957	<0.1%
	Potential river cane restoration/management area on protected land near Tribal lands	382,889	1.2%
	Potential river cane restoration/management area near Tribal lands	4,348,487	14.0%
	Potential river cane restoration/management area on protected land	95,140	0.3%
	Potential river cane restoration/management area	1,792,153	5.8%
↓ Low	Not identified as a rivercane restoration/management area	23,871,925	77.0%
	Area not evaluated for this indicator	*500,333*	*1.6%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Urban park size


This cultural resource indicator measures the size of parks larger than 5 acres in the urban environment. Protected natural areas in urban environments provide urban residents a nearby place to connect with nature, and offer refugia for some species. This indicator complements the equitable access to potential parks indicator by capturing the value of existing parks. It originates from the Protected Areas Database of the United States, U.S. Census Bureau urban areas, and the National Land Cover Database.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

257 miles

64 128



- 75+ acre urban park
- 50 to <75 acre urban park
- 30 to <50 acre urban park
- 10 to <30 acre urban park
- 5 to <10 acre urban park
- <5 acre urban park
- Not identified as an urban park

Table 20: Indicator values for urban park size within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	75+ acre urban park	82,803	0.3%
	50 to <75 acre urban park	1,365	<0.1%
	30 to <50 acre urban park	1,221	<0.1%
	10 to <30 acre urban park	2,206	<0.1%
	5 to <10 acre urban park	976	<0.1%
	<5 acre urban park	610	<0.1%
↓ Low	Not identified as an urban park	30,869,837	99.6%
	Area not evaluated for this indicator	*39,867*	*0.1%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Floodplain inundation

This indicator uses the frequency of inundation to measure the hydrologic function of the floodplain. In a natural floodplain, water can move freely from mainstem rivers and lakes onto different parts of the floodplain. This cycle of frequent, but not persistent, flooding supports many aquatic species and habitats, like fish, waterfowl and swamp forests. It also provides essential ecosystem services such as nutrient retention and downstream flood control. When parts of the floodplain stay constantly wet or dry, this may signal negative impacts from structures like dikes, dams, and levees, which disconnect waterbodies from their adjacent floodplains. This indicator is based on Sentinel-2 satellite imagery and methods developed by Yvonne Allen of the U.S. Fish and Wildlife Service.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Frequency of inundation within the floodplain

- Frequent inundation (flooded in 21-90% of days with available data)
- Regular inundation (flooded in 6-20% of days with available data)
- Occasional inundation (flooded in 2-5% of days with available data)
- Persistent inundation (flooded in 91-100% of days with available data)
- No detected inundation (flooded in 0% of days with available data)
- Not identified as a floodplain

Table 21: Indicator values for floodplain inundation within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Frequency of inundation within the floodplain	Acres	Percent of Area	
↑ High	Frequent inundation (flooded in 21-90% of days with available data)	1,019,684	3.3%	
	Regular inundation (flooded in 6-20% of days with available data)	1,080,626	3.5%	↑ In good condition
	Occasional inundation (flooded in 2-5% of days with available data)	686,043	2.2%	↓ Not in good condition
	Persistent inundation (flooded in 91-100% of days with available data)	645,480	2.1%	
	No detected inundation (flooded in 0% of days with available data)	3,500,406	11.3%	
↓ Low	Not identified as a floodplain	23,955,840	77.3%	
	Area not evaluated for this indicator	*110,806*	*0.4%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Gulf migratory fish connectivity

This indicator captures how far upstream migratory fish in the Gulf of America have been observed. How far upstream migratory fish can travel reflects not just the presence of dams and other barriers, but also the presence of measures like fish ladders that allow specific species to access habitat upstream of dams. This indicator originates from The Nature Conservancy's Southeast Aquatic Connectivity Assessment Project and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Presence of Gulf sturgeon
- Presence of Alabama shad, American shad, or striped bass
- Not identified as Gulf migratory fish habitat (east of the Mississippi River)

Table 22: Indicator values for Gulf migratory fish connectivity within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Presence of Gulf sturgeon	882,689	2.8%	
	Presence of Alabama shad, American shad, or striped bass	1,418	<0.1%	↑ In good condition
↓ Low	Not identified as Gulf migratory fish habitat (east of the Mississippi River)	11,616,772	37.5%	↓ Not in good condition
	Area not evaluated for this indicator	*18,498,005*	*59.7%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Imperiled aquatic species

This indicator measures the number of aquatic animal Regional Species of Greatest Conservation Need (RSGCN) observed within each 12-digit HUC subwatershed, including fish, mussels, snails, crayfish, and amphibians. It captures patterns of rare and endemic aquatic species diversity. RSGCN are regional priority species derived from the list of SGCN identified in Southeast State Wildlife Action Plans as most in need of need of conservation action. RSGCN were chosen based on consistent criteria, such as level of conservation concern, regional stewardship responsibility, and ecological significance. This indicator originates from state Natural Heritage Program data collected by the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Number of aquatic animal Regional Species of Greatest Conservation Need (RSGCN) observed

- 8+ species
- 7 species
- 6 species
- 5 species
- 4 species
- 3 species
- 2 species
- 1 species
- 0 species
- Not identified as a floodplain

Table 23: Indicator values for imperiled aquatic species within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Number of aquatic animal Regional Species of Greatest Conservation Need (RSGCN) observed	Acres	Percent of Area
↑ High	8+ species	652,526	2.1%
	7 species	89,358	0.3%
	6 species	208,761	0.7%
	5 species	247,687	0.8%
	4 species	295,191	1.0%
	3 species	435,655	1.4%
	2 species	849,511	2.7%
	1 species	1,012,993	3.3%
	0 species	2,751,149	8.9%
↓ Low	Not identified as a floodplain	23,955,720	77.3%
	Area not evaluated for this indicator	*500,333*	*1.6%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Lakes & reservoirs

This indicator assesses the condition of lakes and reservoirs based on the amount of natural landcover within the upstream watershed. Higher scores go to lakes and reservoirs with less surrounding urban and agricultural development (including intensive forest management for timber production), which negatively impact drinking water quality and quantity and fish habitat. This indicator uses waterbody and watershed data from the LAGOS-LOCUS dataset, as well as landcover from the National Land Cover Database and LANDFIRE.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Lake or reservoir with low disturbance (0-25%) in upstream watershed
- Lake or reservoir with medium disturbance (>25-60%) in upstream watershed
- Lake or reservoir with high disturbance (>60%) in upstream watershed
- Not identified as a lake or reservoir

Table 24: Indicator values for lakes & reservoirs within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Lake or reservoir with low disturbance (0-25%) in upstream watershed	47,247	0.2%	↑ In good condition
	Lake or reservoir with medium disturbance (>25-60%) in upstream watershed	213,013	0.7%	↓ Not in good condition
	Lake or reservoir with high disturbance (>60%) in upstream watershed	80,842	0.3%	
↓ Low	Not identified as a lake or reservoir	30,651,859	98.9%	
	Area not evaluated for this indicator	*5,923*	*<0.1%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Natural landcover in floodplains

This indicator measures the amount of natural landcover in the estimated floodplain of rivers and streams within each catchment. It assesses the stream channel and its surrounding riparian buffer, measuring the percent of unaltered habitat like forests, wetlands, or open water (rather than agriculture or development). Intact vegetated buffers within the floodplain of rivers and streams provide aquatic habitat, improve water quality, reduce erosion and flooding, recharge groundwater, and more. This indicator originates from the National Land Cover Database and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Percent natural landcover within the estimated floodplain, by catchment

- >90% natural landcover
- >80-90% natural landcover
- >70-80% natural landcover
- >60-70% natural landcover
- ≤60% natural landcover
- Not identified as a floodplain

Table 25: Indicator values for natural landcover in floodplains within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: Percent natural landcover within the estimated floodplain, by catchment	Acres	Percent of Area	
↑ High	>90% natural landcover	3,213,937	10.4%	
	>80-90% natural landcover	745,778	2.4%	↑ In good condition
	>70-80% natural landcover	464,598	1.5%	↓ Not in good condition
	>60-70% natural landcover	360,807	1.2%	
	≤60% natural landcover	1,757,711	5.7%	
↓ Low	Not identified as a floodplain	23,955,840	77.3%	
	Area not evaluated for this indicator	*500,213*	*1.6%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Network complexity

This indicator depicts the number of connected stream size classes in a river network between dams or waterfalls. River networks with a variety of connected stream classes help retain aquatic biodiversity in a changing climate by allowing species to access climate refugia and move between habitats. This indicator originates from the Southeast Aquatic Resources Partnership and applies to the Environmental Protection Agency's estimated floodplain, which spatially defines areas estimated to be inundated by a 100-year flood (also known as the 1% annual chance flood).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Number of connected stream size classes

- 7 size classes
- 6 size classes
- 5 size classes
- 4 size classes
- 3 size classes
- 2 size classes
- 1 size class
- Not identified as a floodplain

Table 26: Indicator values for network complexity within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: Number of connected stream size classes	Acres	Percent of Area	
↑ High	7 size classes	1,480,702	4.8%	
	6 size classes	1,676,570	5.4%	
	5 size classes	1,725,824	5.6%	
	4 size classes	559,661	1.8%	↑ In good condition
	3 size classes	361,453	1.2%	↓ Not in good condition
	2 size classes	395,216	1.3%	
	1 size class	340,548	1.1%	
↓ Low	Not identified as a floodplain	23,956,531	77.3%	
	Area not evaluated for this indicator	*502,380*	*1.6%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater


Permeable surface

This indicator measures the average percent of non-impervious cover within each catchment. High levels of impervious surface degrade water quality and alter freshwater flow, impacting both aquatic species communities and ecosystem services for people, like the availability of clean drinking water. This indicator originates from the National Land Cover Database.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Percent of catchment permeable

- >95% permeable (likely high water quality and supporting most sensitive aquatic species)
- >90-95% permeable (likely declining water quality and supporting most aquatic species)
- >70-90% permeable (likely degraded water quality and not supporting many aquatic species)
- ≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)

Table 27: Indicator values for permeable surface within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Percent of catchment permeable	Acres	Percent of Area	
↑ High	>95% permeable (likely high water quality and supporting most sensitive aquatic species)	28,811,583	92.9%	↑ In good condition
	>90-95% permeable (likely declining water quality and supporting most aquatic species)	764,043	2.5%	↓ Not in good condition
	>70-90% permeable (likely degraded water quality and not supporting many aquatic species)	738,443	2.4%	
↓ Low	≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)	184,482	0.6%	
	Area not evaluated for this indicator	*500,333*	*1.6%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Coastal shoreline condition

This indicator assesses shoreline condition based on the presence of hardened structures like jetties, groins, and riprap, as well as other human development. By restricting the natural movement of sediment, shoreline armoring increases erosion, prevents the inland migration of coastal ecosystems in response to sea level rise, and degrades habitat for birds, sea turtles, fish, plants, and other species both on and offshore. Natural shorelines in harder-to-develop coastal areas receive the highest shoreline condition scores, while hardened shorelines receive the lowest scores. This indicator originates from the National Oceanic and Atmospheric Administration's Environmental Sensitivity Index dataset.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Natural and harder to develop
- Natural
- Partially armored and harder to develop
- Partially armored
- Armored

Table 28: Indicator values for coastal shoreline condition within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Natural and harder to develop	1,727	<0.1%	
	Natural	19,259	<0.1%	↑ In good condition
	Partially armored and harder to develop	6.2	<0.1%	↓ Not in good condition
	Partially armored	343	<0.1%	
↓ Low	Armored	2,438	<0.1%	
	Area not evaluated for this indicator	*30,975,111*	*99.9%*	
	Total area	**30,998,884**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Estuarine coastal condition

This indicator combines measures of water quality, sediment quality, contaminants in fish tissue, and benthic community condition to create an overall index of coastal estuarine condition. Estuaries serve as important nursery habitat for wildlife, including many species of fish and shellfish eaten as seafood. They also improve water quality by filtering out sediments and pollutants, provide recreational opportunities, and support coastal economies. This indicator originates from the Environmental Protection Agency's National Coastal Condition Assessment data.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Good
- Fair to good
- Fair
- Poor to fair
- Poor
- Shallow estuary not assessed for condition

Table 29: Indicator values for estuarine coastal condition within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values	Acres	Percent of Area	
↑ High	Good	127,599	0.4%	
	Fair to good	155,026	0.5%	↑ In good condition
	Fair	167,175	0.5%	↓ Not in good condition
	Poor to fair	0	0%	
	Poor	0	0%	
↓ Low	Shallow estuary not assessed for condition	18,912	<0.1%	
	Area not evaluated for this indicator	*30,530,172*	*98.5%*	
	Total area	**30,998,884**	**100%**	


To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Gulf coral & hardbottom

This indicator predicts the presence of coral and hardbottom in the Gulf of America based on direct observations, acoustic surveys, and known locations of artificial reefs and shipwrecks. Hardbottom provides an anchor for important seafloor habitats such as deep-sea corals, plants, and sponges, providing valuable structure that supports a wide range of invertebrate and fish species. Hardbottom is also sometimes associated with diverse chemosynthetic communities supported by micro-organisms that feed off of hydrocarbon seeps. This indicator combines data from multiple sources, including Bureau of Ocean Energy Management seismic water bottom anomalies, usSEABED sediment data, several National Oceanic and Atmospheric Administration datasets, various state layers, and more.





- Confirmed hardbottom-associated species (e.g., corals, sponges, patch reef, chemosynthetic communities)
- Confirmed natural hardbottom
- Artificial reefs
- Shipwrecks
- Probable natural hardbottom (fine resolution)
- Rock (coarse resolution)
- Gravel (coarse resolution)
- Not identified as hardbottom

Table 30: Indicator values for Gulf coral & hardbottom within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Confirmed hardbottom-associated species (e.g., corals, sponges, patch reef, chemosynthetic communities)	0	0%
	Confirmed natural hardbottom	0	0%
	Artificial reefs	1,870	<0.1%
	Shipwrecks	612	<0.1%
	Probable natural hardbottom (fine resolution)	0	0%
	Rock (coarse resolution)	17,910	<0.1%
	Gravel (coarse resolution)	7,154	<0.1%
↓ Low	Not identified as hardbottom	588,886	1.9%
	Area not evaluated for this indicator	*30,382,453*	*98.0%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Gulf marine birds

This indicator identifies important areas in the Gulf of America for birds that feed exclusively or mainly at sea. It uses habitat suitability models for 24 species of marine birds (black-capped petrel, black tern, Bonaparte's gull, brown booby, brown noddy, brown pelican, bridled tern, band-rumped storm-petrel, Cory's shearwater, common loon, common tern, great shearwater, herring gull, laughing gull, masked booby, magnificent frigatebird, northern gannet, parasitic jaeger, royal tern, sandwich tern, sargasso shearwater, sooty tern, Wilson's storm-petrel, and pomarine jaeger) based on sightings from vessel-based and aerial surveys. This indicator originates from the Gulf of Mexico Marine Assessment Program for Protected Species (GoMMAPPS) seabird models.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- High marine bird richness (9-16 of 24 modeled species)
- Medium marine bird richness (6-8 of 24 modeled species)
- Low marine bird richness (0-5 of 24 modeled species)

Table 31: Indicator values for Gulf marine birds within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	High marine bird richness (9-16 of 24 modeled species)	28,391	<0.1%
	Medium marine bird richness (6-8 of 24 modeled species)	13,503	<0.1%
↓ Low	Low marine bird richness (0-5 of 24 modeled species)	230,364	0.7%
	Area not evaluated for this indicator	*30,726,626*	*99.1%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Gulf marine mammals

This indicator identifies important areas in the Gulf of America for dolphins and whales. It incorporates monthly density predictions for 13 marine mammal species or species groups (Atlantic spotted dolphin, beaked whales, blackfish [which includes killer whale, melon-headed whale, false killer whale, pygmy killer whale], bottlenose dolphin, Bryde's whale, clymene dolphin, pantropical spotted dolphin, pilot whales, pygmy/dwarf sperm whales, Rice's whale, Risso's dolphin, sperm whale, spinner dolphin) based on sightings from boat-based and aerial surveys and data on oceanographic conditions. It uses marine mammal models developed by the National Oceanic and Atmospheric Administration as part of the Gulf of Mexico Marine Assessment Program for Protected Species (GoMMAPPS).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Percentile of importance for marine mammal index species (across larger analysis area)

- >90th percentile
- >80th-90th percentile
- >70th-80th percentile
- >60th-70th percentile
- >50th-60th percentile
- >40th-50th percentile
- >30th-40th percentile
- >20th-30th percentile
- >10th-20th percentile
- ≤10th percentile
- Land

Table 32: Indicator values for Gulf marine mammals within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for marine mammal index species (across larger analysis area)	Acres	Percent of Area
↑ High	>90th percentile	0	0%
	>80th-90th percentile	150	<0.1%
	>70th-80th percentile	14,016	<0.1%
	>60th-70th percentile	17,322	<0.1%
	>50th-60th percentile	12,608	<0.1%
	>40th-50th percentile	924	<0.1%
	>30th-40th percentile	3,683	<0.1%
	>20th-30th percentile	0	0%
	>10th-20th percentile	0	0%
	≤10th percentile	428,982	1.4%
↓ Low	Land	2,500	<0.1%
	Area not evaluated for this indicator	*30,518,700*	*98.5%*
	Total area	**30,998,884**	**100%**


To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Gulf sea turtles

This indicator identifies important areas in the Gulf of America for sea turtles. It incorporates monthly density predictions for four species (green, Kemp's ridley, leatherback, and loggerhead sea turtles) based on sightings from boat-based and aerial surveys and data on oceanographic conditions. It uses sea turtle models developed by the National Oceanic and Atmospheric Administration as part of the Gulf of Mexico Marine Assessment Program for Protected Species (GoMMAPPS).





Percentile of importance for sea turtle index species (across larger analysis area)

- >90th percentile
- >80th-90th percentile
- >70th-80th percentile
- >65th-70th percentile
- ≤65th percentile
- Land

Table 33: Indicator values for Gulf sea turtles within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values: Percentile of importance for sea turtle index species (across larger analysis area)	Acres	Percent of Area
↑ High	>90th percentile	6,655	<0.1%
	>80th-90th percentile	42,218	0.1%
	>70th-80th percentile	0	0%
	>65th-70th percentile	0	0%
	≤65th percentile	451,321	1.5%
↓ Low	Land	176,780	0.6%
	Area not evaluated for this indicator	*30,321,910*	*97.8%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Island habitat

This indicator represents important habitat for coastal island-dependent species across the Southeast. Because the isolation of islands can make them ecologically unique and protect them from disturbance and mainland predators, they often serve as important habitat for many species of mammals, plants, and insects, as well as breeding coastal birds and sea turtles. The highest scores go to island critical habitat for six threatened and endangered animal and plant species: piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, and Bartram's hairstreak butterfly. This indicator uses U.S. Fish and Wildlife Service critical habitat data and island boundaries from the U.S. Geological Survey and Esri.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



- Island critical habitat for any of six threatened and endangered species (piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, or Bartram's hairstreak butterfly)
- Other island area
- Not a coastal island

Table 34: Indicator values for island habitat within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Island critical habitat for any of six threatened and endangered species (piping plover, loggerhead sea turtle, Cape Sable thoroughwort, Florida semaphore cactus, silver rice rat, or Bartram's hairstreak butterfly)	7,736	<0.1%
	Other island area	4,801	<0.1%
↓ Low	Not a coastal island	1,962,709	6.3%
	Area not evaluated for this indicator	*29,023,638*	*93.6%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Resilient coastal sites

This indicator depicts the capacity of coastal habitats to migrate to adjacent lowlands in order to sustain biodiversity and natural services under increasing inundation from sea level rise. It is based on the physical and condition characteristics of current tidal complexes, their predicted migration space, and surrounding buffer areas. These characteristics include marsh complex size, shared edge with migration space, sediment balance, water quality, natural landcover, landform diversity, and more. This indicator originates from The Nature Conservancy's Resilient Coastal Sites project.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

257 miles
64 128



- Most resilient
- More resilient
- Slightly more resilient
- Average/median resilience
- Slightly less resilient
- Less resilient
- Least resilient
- Sea level rise area

Table 35: Indicator values for resilient coastal sites within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Most resilient	0	0%
	More resilient	94,572	0.3%
	Slightly more resilient	61,870	0.2%
	Average/median resilience	18,617	<0.1%
	Slightly less resilient	394	<0.1%
	Less resilient	307	<0.1%
	Least resilient	138	<0.1%
↓ Low	Sea level rise area	10,312	<0.1%
	Area not evaluated for this indicator	*30,812,674*	*99.4%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Seagrass

This indicator represents the presence of seagrass in the Atlantic Ocean and Gulf of America. Seagrasses provide food and habitat for a range of marine and estuarine wildlife, including fish, sea turtles, shrimp, crabs, oysters, and more. They also produce oxygen, filter water, control erosion, and buffer storms. Seagrasses serve as an important indicator of the overall health of coastal ecosystems because they are sensitive to water quality and require sufficiently clear water for sunlight to penetrate. This indicator originates from the National Oceanic and Atmospheric Administration's Marine Cadastre.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

257 miles
64 128



Seagrass present

Table 36: Indicator values for seagrass within Mississippi. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Seagrass present	339	<0.1%
	Area not evaluated for this indicator	*30,998,545*	*100.0%*
	Total area	**30,998,884**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine


Stable coastal wetlands

This indicator uses remote sensing to calculate the unvegetated-vegetated ratio of tidal wetlands, which compares how much of a wetland is not covered by plants (e.g., sediment, rocks, open water) to how much is covered by plants. Marshes that maintain a higher proportion of vegetation tend to be more stable and resilient to threats like sea level rise, erosion, and coastal development. This ratio, and how it changes over time, is a good surrogate for salt marsh degradation processes like sediment loss and conversion to open water. This indicator originates from a U.S. Geological Survey project on an unvegetated to vegetated ratio for coastal wetlands.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

257 miles
64
128



- Stable coastal wetlands
- Other coastal wetlands
- Not identified as coastal wetlands

Table 37: Indicator values for stable coastal wetlands within Mississippi. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Stable coastal wetlands	70,537	0.2%	↑ In good condition
	Other coastal wetlands	22,497	<0.1%	↓ Not in good condition
↓ Low	Not identified as coastal wetlands	727,651	2.3%	
	Area not evaluated for this indicator	*30,178,200*	*97.4%*	
	Total area	**30,998,884**	**100%**	


To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

More Information

Priority Amphibian and Reptile Conservation Areas

Priority Amphibian and Reptile Conservation Areas (PARCAs) are an expert-driven, nonregulatory designation that includes places capable of supporting viable amphibian and reptile populations, places occupied by rare or imperiled species, and places rich in biodiversity or species unique to that geographic area. Reptiles and amphibians are a critical part of the Southeast region's rich biodiversity and many populations are declining in the face of threats like habitat loss, invasive species, and climate change. The PARCA dataset is maintained by the Amphibian and Reptile Conservancy and does not yet include Virginia or Kentucky.





Priority Amphibian and Reptile Conservation Area (PARCA)

Not a PARCA (excluding Kentucky and Virginia)

Table 38: Extent of Priority Amphibian and Reptile Conservation Areas within Mississippi.

Priority Amphibian and Reptile Conservation Areas status	Acres	Percent of Area
Priority Amphibian and Reptile Conservation Area (PARCA)	6,309,181	20.4%
Not a PARCA (excluding Kentucky and Virginia)	24,689,348	79.6%
Total area	**30,998,884**	**100%**

Priority Amphibian and Reptile Conservation Areas at this location:

Gulf Coast

The Gulf Coast PARCA spans the southern portion of Mississippi, encompassing a diverse landscape influenced by its proximity to the Gulf of Mexico. This region includes coastal flatwoods and marshes, upland longleaf pine forests, and expansive riverine systems, with notable features such as the Pascagoula River, renowned for its biodiversity. Habitats range from saltwater and freshwater marshes to fire-maintained longleaf pine forests, supporting species like the dusky gopher frog, Gulf salt marsh snake, and Mississippi diamondback terrapin. The confluence of these subregions, coupled with the area's history of fluctuating sea levels and sediment deposition, has sculpted a unique set of conditions conducive to supporting a wide range of plant and animal life, each with a distinct subset of species, threats, and conservation needs, underpinning its importance for conservation. Urban development, invasive species, and fire suppression are major threats that collaborative conservation efforts with national forests, national wildlife refuges, and educational institutions are crucial to address.

Mann's Land

Named in honor of herpetologist Tom Mann, a longtime herpetologist in the region, the Mann's Land PARCA, located in east-central Mississippi, includes the Tombigbee National Forest and Black Prairie Wildlife Management Area. This PARCA features a mosaic of open prairies, chalk outcrops, and oak-hickory forests. The unique geology, including alkaline soils supporting prairie flora, provides habitats for species like the Webster's salamander, crawfish frog, and prairie kingsnake. Key threats are prairie conversion, invasive species like cogongrass, and incompatible forestry practices. Conservation efforts can be supported by partnerships with Mississippi State University, private landowners, and the Natural Resources Conservation Service (NRCS). Important research needs include mapping surface geology for salamanders and monitoring the effectiveness of prairie restoration efforts.

Mobile Bay

Alabama's largest PARCA, Mobile Bay, contains a largely developed metro area, and in a region of such high biodiversity, this has pushed many species to the brink of extirpation. This area is considered the world's center of turtle diversity, harboring beautiful species like the eastern chicken turtle, southern painted turtle, Alabama red-bellied turtle, and Mississippi diamondback terrapin. Often called "America's Amazon," it was the boyhood home of EO Wilson; his dream was that it would one day hold a national park to celebrate its amazing biodiversity. Undeveloped areas along the coast and inland swamps are key habitats for a large number of imperiled amphibian and reptile species that find refuge or breed here, including the Gulf Coast waterdog.

Pearl River Basin

Boasting more than 100 species of reptiles and amphibians, the Pearl River Basin is widely considered one of America's crown jewels for herpetological biodiversity. It consists primarily of upland forest dominated by evergreen/mixed hardwoods, pine flatwoods, and forested wetlands. Much of the historical longleaf pine component within the woodlands and flatwoods has been converted to loblolly pine. Ongoing efforts continue to restore longleaf and improve habitat conditions to these vital communities. This area faces a large number of human threats from urbanization, water pollution, and unsustainable timber practices. The natural areas here are also threatened by invasive species (wild hogs and cogongrass), as well as fire suppression and increased hurricane frequency. A significant effort will need to be made by cooperating

across the many individuals and organizations who occupy this land to control point source pollution, return pine forests to natural fire conditions, control invasive species, and restore many of these habitats.

South Delta
The South Delta PARCA, nestled between the Mississippi and Yazoo Rivers, is characterized by rich alluvial soils that have historically supported extensive agriculture. Despite this transformation, remnants of bottomland hardwood forests and cypress-tupelo swamps persist within protected areas like Panther Swamp and Hillside National Wildlife Refuges. These wetlands are vital for species such as the western chicken turtle, alligator snapping turtle, and small-mouthed salamander. The region faces threats from agricultural runoff, invasive species like feral hogs and Chinese tallow trees, and altered hydrology. Conservation strategies include working with the Army Corps of Engineers, refining agricultural practices, and restoring abandoned catfish ponds to benefit aquatic species. Research efforts are needed to assess the impacts of invasive species and monitor turtle populations.

Tishomingo
The Tishomingo PARCA, located in the northeastern corner of Mississippi, is a unique ecological region due to its position at the southwestern reach of the Appalachian foothills. This area, part of the Cumberland Plateau, features rugged topography and distinct geological formations that merge into the Fall Line Hills and Blackbelt Prairie. The mixed hardwood and pine forests provide critical habitats for species like the eastern hellbender, green salamander, and northern pine snake. Threats to this region include non-compatible timber practices, river sedimentation, and agricultural runoff. Conservation efforts can be bolstered through partnerships with regional authorities, state parks, and local universities. Key research needs include expanded surveys for hellbenders and studies on the impacts of quarry operations.

Tunica Hills
The Tunica Hills PARCA is unique in that it is predominantly composed of the southern mesophytic hardwood forest community type that developed on loess hills with steep ravines and intermittent or spring-fed streams. Currently, this community type is only recognized from this area of Louisiana. In addition to southern mesophytic hardwoods, other associated community types within the Tunica Hills PARCA include hardwood slope forests and mixed hardwood-loblolly pine forests. Threats to this PARCA include water pollution (from sand and gravel mining), habitat conversion from hardwoods to pine, unsustainable timber harvests, and invasive species (wild hogs). Major priorities for this PARCA include adhering to best management practices and streamside management zones, maintaining late stage hardwoods, and creating/restoring ephemeral wetlands.


Upper Pearl
The Upper Pearl PARCA stretches along the Pearl River from its meeting with the Gulf Coast to central Mississippi. This region includes dynamic aquatic ecosystems and dense riparian forests of oak, hickory, and pine. The Pearl River is crucial for freshwater turtles like the Pearl River map turtle, ringed sawback, and alligator snapping turtle. Threats include altered hydrology from impoundments, invasive species, and agricultural runoff, which can be mitigated through conservation opportunities involving collaboration with water management districts, local municipalities, and environmental organizations. Research priorities include long-term turtle monitoring and habitat usage studies for key species that rely

on the pristine waters that support one of the most diverse turtle populations in the country.

Names and descriptions of public Priority Amphibian and Reptile Areas were provided by the Amphibian and Reptile Conservancy on August 30, 2024 and edited slightly for clarity and consistency.


Protected Areas

Protected areas include a variety of public lands owned or managed by federal, state, and local agencies and nonprofit organizations, as well as some private conservation lands and conservation easements. Protected areas are derived from the Protected Areas Database of the United States (PAD-US v4.1) and include Fee, Designation, Easement, Marine, and Proclamation (Dept. of Defense lands only) boundaries.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Within a protected area

Not within a protected area

Table 39: Extent of protected areas within Mississippi.

Protected area status	Acres	Percent of Area
Not within a protected area	27,898,533	90.0%
Within a protected area	3,100,351	10.0%
Total area	**30,998,884**	**100%**

Protected areas at this location:

- Mississippi 16th Section Public School Trust Lands (646,178 acres)
- De Soto National Forest (USDA Forest Service; 532,431 acres)
- Homochitto National Forest (USDA Forest Service; 191,846 acres)
- Bienville National Forest (USDA Forest Service; 180,227 acres)
- Holly Springs National Forest (USDA Forest Service; 156,210 acres)
- Camp Shelby (138,435 acres)
- Tombigbee National Forest (USDA Forest Service; 67,449 acres)
- Delta National Forest (USDA Forest Service; 61,472 acres)
- Sunflower Wma (59,101 acres)
- Gulf Islands National Seashore (58,525 acres)
- Gulf Islands National Seashore (National Park Service; 57,793 acres)
- Upper Sardis Wma (49,322 acres)
- Sam D. Hamilton Noxubee National Wildlife Refuge (US Fish and Wildlife Service; 48,347 acres)
- Leaf River Wma (41,418 acres)
- Sardis Lake (39,413 acres)
- Pascagoula Wma (MDWFP; 37,758 acres)
- Panther Swamp National Wildlife Refuge (US Fish and Wildlife Service; 33,162 acres)
- Natchez Trace Parkway (National Park Service; 31,527 acres)
- Canal Section Wma (COE; 29,202 acres)
- Chickasawhay Wma (29,055 acres)
- Caney Creek Wma (28,447 acres)
- Caston Creek Wma (28,288 acres)
- Tallahala Wma (27,545 acres)
- Chickasaw Wma (26,802 acres)
- Bienville Wma (26,770 acres)
- ... and 928 more protected areas ...


Note: areas are listed based on name, ownership, and boundary information in the Protected Areas Database of the United States, which may include overlapping and duplicate areas. Only protected areas with at least 1 acre of overlap with this area are included in the list above.

Sea Level Rise

NOAA's sea level rise (SLR) inundation models represent areas likely to experience flooding at high tide based on each foot of SLR above current levels. Darker blue areas will experience flooding first, and at greater depth, compared to lighter blue areas. These models are not linked to a future timeframe; see the projections below. NOAA calculates the inundation footprint at "mean higher high water", or the average highest daily tide. The area covered in each SLR scenario includes areas projected to be inundated at lower levels. For example, the area inundated by 4 ft of SLR also includes areas inundated by 3 ft, 2 ft, 1 ft, and 0 ft of SLR (where 0 ft represents current levels).

To explore additional SLR information, please see NOAA's Sea Level Rise Viewer.





Flooding extent by projected sea level rise (ft)



Table 40: Extent of flooding by projected average highest daily tide due to sea level rise within Mississippi. Values from the NOAA sea level rise inundation data.

Feet of sea level rise	Acres	Percent of Area
0 feet	535,278	1.7%
1 foot	572,401	1.8%
2 feet	597,478	1.9%
3 feet	613,216	2.0%
4 feet	626,417	2.0%
5 feet	639,873	2.1%
6 feet	654,871	2.1%
7 feet	667,808	2.2%
8 feet	683,025	2.2%
9 feet	701,309	2.3%
10 feet	719,453	2.3%
Not projected to be inundated by up to 10 feet	*939,210*	*3.0%*
Sea level rise is unlikely to be a threat (inland counties)	*29,334,730*	*94.6%*
No sea level rise data available	*5,491*	*<0.1%*
Total area	**30,998,884**	**100%**


Table 41: Projected sea level rise by decade within Mississippi. Values are based on area-weighted averages of decadal projections for 1-degree grid cells that overlap this area based on NOAA's 2022 Sea Level Rise Report. 2060 corresponds to the SECAS goal: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.

SLR Scenario	2020 (ft)	2030 (ft)	2040 (ft)	2050 (ft)	2060 (ft)	2070 (ft)	2080 (ft)	2090 (ft)	2100 (ft)
Low	0.42	0.64	0.87	1.1	1.3	1.5	1.6	1.8	2
Intermediate-low	0.45	0.7	0.97	1.2	1.5	1.8	2	2.3	2.6
Intermediate	0.45	0.72	1	1.4	1.7	2.2	2.7	3.3	4
Intermediate-high	0.45	0.76	1.1	1.6	2.1	2.9	3.7	4.6	5.6
High	0.45	0.77	1.2	1.8	2.5	3.5	4.7	5.9	7.2

Urban Growth

The FUTURES urban growth model predicts the likelihood that an area will urbanize at every decade from 2020 to 2100. Developed areas from the 2021 National Landcover Database serve as the baseline for current urban areas. The model simulates landscape change based on trends in population growth, local development suitability factors, and an urban patch-growing algorithm. It considers environmental drivers like distance to floodplain, slope, and available infrastructure, and even socio-economic status. The probability of urbanization for each area reflects how many times it urbanized out of 50 model runs.

To explore maps for additional time periods, click here.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

64 128 257 miles



Probability of urbanization by 2060

- Urban in 2021
- Very high likelihood of urbanization (>50% probability)
- High likelihood of urbanization (25 - 50% probability)
- Moderate likelihood of urbanization (2 - 25% probability)
- Not likely to urbanize

Table 42: Extent of projected urbanization by decade within Mississippi. Values from FUTURES model projections for the contiguous United States developed by the Center for Geospatial Analytics, NC State University. 2060 corresponds to the SECAS goal: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.


Decade	Acres	Percent of Area
Urban in 2021	1,998,807	6.4%
2030 projected extent	2,054,182	6.6%
2040 projected extent	2,076,003	6.7%
2050 projected extent	2,090,126	6.7%
2060 projected extent	2,101,899	6.8%
2070 projected extent	2,109,279	6.8%
2080 projected extent	2,114,429	6.8%
2090 projected extent	2,117,532	6.8%
2100 projected extent	2,119,221	6.8%
Not projected to urbanize by 2100	*28,879,308*	*93.2%*
No urbanization data available	*355*	*<0.1%*
Total area	**30,998,884**	**100%**

6.4% of this area is already urban in 2021, and an additional 7.2% has at least a moderate probability of urbanizing by 2060.

By 2060, the size of the urban footprint is projected to increase **5.2%** over 2021 levels.

Wildfire Likelihood

Wildfire likelihood data originate from the Wildfire Risk to Communities project developed by the U.S. Forest Service. This layer depicts the probability of wildfire burning in a specific location in any given year. Annual burn probabilities in the United States range from 0-14%, but do not exceed 8% in the Southeast. Wildfire likelihood is based on fire behavior modeling across thousands of simulations of possible fire seasons. In each simulation, factors contributing to the probability of a fire occurring (such as weather, topography, and ignitions) vary based on patterns derived from observations in recent decades. Wildfire likelihood is not predictive and does not reflect any forecasted future weather or fire danger conditions. It also does not say anything about the intensity of fire if it occurs. To explore additional wildfire risk information, please see the Wildfire Risk to Communities website.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

257 miles
64 128



Wildfire likelihood (annual burn probability)

- High
- Moderate-high
- Moderate
- Low-moderate
- Low
- Not predicted to experience wildfire

Table 43: Area in each wildfire probability category within Mississippi. Values from the Wildire Risk To Communities project developed by the USDA Forest Service.

Wildfire likelihood (annual burn probability)	Acres	Percent of Area
Not predicted to experience wildfire (0% probability)	2,968,483	9.6%
Low (>0 - 0.01% probability)	1,356,803	4.4%
Low-moderate (>0.01 - 0.02154% probability)	881,903	2.8%
Low-moderate (>0.02154 - 0.04642% probability)	3,766,306	12.1%
Moderate (>0.04642 - 0.1% probability)	8,719,615	28.1%
Moderate (>0.1 - 0.21544% probability)	8,126,716	26.2%
Moderate (>0.21544 - 0.46416% probability)	3,377,660	10.9%
Moderate-high (>0.46416 - 1% probability)	1,656,151	5.3%
Moderate-high (>1 - 2.15443% probability)	140,649	0.5%
High (>2.15443 - 4.64159% probability)	0	0%
High (>4.64159% probability)	0	0%
No wildfire risk data available	*4,599*	*<0.1%*
Total area	**30,998,884**	**100%**