Southeast Conservation Blueprint Summary

for United States Virgin Islands

Created 10/21/2026

Table of Contents

About the Southeast Blueprint 3

Southeast Blueprint Priorities 4

Hubs and Corridors 6

Indicator Summary 8

More Information 49

The Southeast Conservation Blueprint 2026

Southeast Conservation Adaptation Strategy

[THIS PAGE INTENTIONALLY LEFT BLANK]

About the Southeast Blueprint

The Southeast Conservation Blueprint is the primary product of the Southeast Conservation Adaptation Strategy (SECAS). It is a living, spatial plan to achieve the SECAS vision of a connected network of lands and waters across the Southeast and Caribbean. The Blueprint is regularly updated to incorporate new data, partner input, and information about on-the-ground conditions.

The Blueprint identifies priority areas based on a suite of natural and cultural resource indicators representing terrestrial, freshwater, and marine ecosystems. A connectivity analysis identifies corridors that link coastal and inland areas and span climate gradients.

For more information:

- Visit the Blueprint webpage
- Review the Blueprint 2026 Development Process
- View and download the Blueprint data and make maps on the Blueprint page of the SECAS Atlas

We're here to help!


- Do you have a question about the Blueprint?
- Would you like help using the Blueprint to support a proposal or inform a decision?
- Do you have a suggestion on how to improve the Blueprint? The Blueprint and its inputs are regularly revised based on input from people like you.
- Do you have feedback on how to improve the Blueprint Explorer interface?

If you need help or have questions, contact Southeast Blueprint staff by reaching out to a member of the user support team.

We're here to support you. We really mean it. It's what we do!

This report was generated by the Southeast Conservation Blueprint Explorer, which was developed by Astute Spruce, LLC in partnership with the U.S. Fish and Wildlife Service under the Southeast Conservation Adaptation Strategy.

Southeast Blueprint Priorities



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Priorities for a connected network of lands and waters

- Highest priority
- High priority
- Medium priority
- Priority connections

Priority Categories

For a connected network of lands and waters

In total, Blueprint priorities and priority connections cover roughly 50% of the Southeast Blueprint geography.

Highest priority

Areas where conservation action would make the biggest impact, based on a suite of natural and cultural resource indicators. This class covers roughly 10% of the Southeast Blueprint geography.

High priority

Areas where conservation action would make a big impact, based on a suite of natural and cultural resource indicators. This class covers roughly 15% of the Southeast Blueprint geography.

Medium priority

Areas where conservation action would make an above-average impact, based on a suite of natural and cultural resource indicators. This class covers roughly 20% of the Southeast Blueprint geography.

Priority connections

Connections between priority areas that cover the shortest distance possible while routing through as much Blueprint priority as possible. This class covers roughly 5% of the Southeast Blueprint geography.


Table 1: Extent of each Blueprint priority category within United States Virgin Islands.

Priority Category	Acres	Percent of Area
Highest priority	25,091	10.0%
High priority	37,636	15.0%
Medium priority	56,271	22.4%
Priority connections	14,704	5.9%
Lower priority	117,208	46.7%
Total area	**250,911**	**100%**

Hubs and Corridors

The Blueprint uses a least-cost path connectivity analysis to identify corridors that link hubs across the shortest distance possible, while also routing through as much Blueprint priority as possible.

In the Caribbean, hubs are large patches (~500+ acres) of highest priority Blueprint areas and/or protected lands.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Hubs
Corridors

Table 2: Extent of hubs and corridors within United States Virgin Islands.

Type	Acres	Percent of Area
Hubs	34,992	13.9%
Corridors	57,173	22.8%
Not a hub or corridor	158,745	63.3%
Total area	**250,911**	**100%**

Indicator Summary

Table 3: Terrestrial indicators.

Indicator	Present
Caribbean greenways & trails	✓
Caribbean habitat patch size (small islands)	✓
Caribbean island habitat	✓
Caribbean karst habitat	✓
Caribbean landscape condition	✓
Caribbean low-urban historic landscapes	✓
Caribbean reforestation potential	✓
Caribbean urban park size	✓
U.S. Virgin Islands imperiled terrestrial species	✓

Table 4: Freshwater indicators.

Indicator	Present
Caribbean imperiled aquatic species	✓
Caribbean natural landcover in floodplains	✓
Caribbean network complexity	✓
Caribbean permeable surface	✓

Table 5: Coastal & marine indicators.

Indicator	Present
Caribbean beach habitat	✓
Caribbean coastal shoreline condition	✓
Caribbean fish hotspots	✓
Caribbean fish nursery habitat	✓
Caribbean mangrove migration	✓
Caribbean seagrass	✓
Caribbean shallow hardbottom & coral	✓

Terrestrial

Caribbean greenways & trails

This cultural resource indicator measures both the natural condition and connected length of greenways and trails in the U.S. Caribbean to characterize the quality of the recreational experience. Natural condition is based on the amount of impervious surface surrounding the path. Connected length captures how far a person can go without leaving a dedicated path, based on common distances for walking, running, and biking. This indicator originates from OpenStreetMap data and the National Oceanic and Atmospheric Administration's Coastal Change Analysis Program landcover.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km
- Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km
- Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km
- Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km
- Developed and connected for <1.9 km
- Sidewalk
- Not identified as a trail, sidewalk, or other path

Table 6: Indicator values for Caribbean greenways & trails within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Mostly natural and connected for 5 to <40 km or partly natural and connected for ≥40 km	283	0.1%	
	Mostly natural and connected for 1.9 to <5 km, partly natural and connected for 5 to <40 km, or developed and connected for ≥40 km	223	<0.1%	
	Mostly natural and connected for <1.9 km, partly natural and connected for 1.9 to <5 km, or developed and connected for 5 to <40 km	251	0.1%	↑ In good condition
	Partly natural and connected for <1.9 km or developed and connected for 1.9 to <5 km	245	<0.1%	↓ Not in good condition
	Developed and connected for <1.9 km	272	0.1%	
	Sidewalk	14	<0.1%	
↓ Low	Not identified as a trail, sidewalk, or other path	94,558	37.7%	
	Area not evaluated for this indicator	*155,065*	*61.8%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Caribbean habitat patch size (small islands)

This indicator represents the size of natural habitat patches on small islands in the U.S. Caribbean that are unfragmented by roads, urban development, or agriculture. Large areas of intact natural habitat are important for many wildlife species, including reptiles and amphibians, birds, and large mammals. It uses LANDFIRE landcover and road data, mimicking Esri's intact habitat cores approach from their green infrastructure data.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Large patch (>1,000 acres)
- Medium patch (>100-1,000 acres)
- Small patch (>10-100 acres)
- Very small patch (≤10 acres)
- Developed or agriculture

Table 7: Indicator values for Caribbean habitat patch size (small islands) within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Large patch (>1,000 acres)	20,316	8.1%	
	Medium patch (>100-1,000 acres)	23,963	9.6%	↑ In good condition
	Small patch (>10-100 acres)	9,066	3.6%	↓ Not in good condition
	Very small patch (≤10 acres)	6,151	2.5%	
↓ Low	Developed or agriculture	29,056	11.6%	
	Area not evaluated for this indicator	*162,358*	*64.7%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Caribbean island habitat

This indicator represents the importance of island habitat in the U.S. Caribbean for federally listed and other imperiled species based on the presence of imperiled and invasive animals. The isolation of islands often makes them ecologically unique and protects them from disturbance and mainland predators; however, these factors also make them more vulnerable to invasive species. This indicator uses species data from Island Conservation's Threatened Island Biodiversity Database, U.S. Fish and Wildlife Service critical habitat, and the Puerto Rico and U.S. Virgin Islands Gap Analysis Project.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Island area with critical habitat for a threatened or endangered species
- Island area with no invasive animals and 2+ imperiled species
- Island area with no invasive animals and 1 imperiled species
- Island area with no invasive animals
- Island area with invasive animals and 2+ imperiled species
- Island area with invasive animals and 1 imperiled species
- Island area with invasive animals
- Not an island

Table 8: Indicator values for Caribbean island habitat within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Island area with critical habitat for a threatened or endangered species	2,834	1.1%
	Island area with no invasive animals and 2+ imperiled species	92	<0.1%
	Island area with no invasive animals and 1 imperiled species	188	<0.1%
	Island area with no invasive animals	0	0%
	Island area with invasive animals and 2+ imperiled species	5,498	2.2%
	Island area with invasive animals and 1 imperiled species	18,781	7.5%
	Island area with invasive animals	61,160	24.4%
↓ Low	Not an island	162,358	64.7%
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Caribbean karst habitat

This indicator for the U.S. Caribbean represents natural karst areas with limited human alteration from activities such as urban development and intensive agriculture. Karst is a geologically unique landscape where the movement of water through easily dissolved bedrock, particularly limestone, produces distinctive features like caves, sinkholes, and underground rivers. Areas characterized by karst geology support many unique and endemic species, help recharge freshwater aquifers, and often contain significant cultural and historical sites. This indicator combines LANDFIRE land cover with karst datasets from the Puerto Rico Department of Natural and Environmental Resources, National Park Service, and U.S. Geological Survey.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Karst with natural landcover
- Karst with pasture, crops, or open space in developed area
- Karst with developed landcover
- Not identified as karst

Table 9: Indicator values for Caribbean karst habitat within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Karst with natural landcover	7,696	3.1%	↑ In good condition
	Karst with pasture, crops, or open space in developed area	4,700	1.9%	↓ Not in good condition
	Karst with developed landcover	1,583	0.6%	
↓ Low	Not identified as karst	71,879	28.6%	
	Area not evaluated for this indicator	*165,053*	*65.8%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Caribbean landscape condition

This indicator for the U.S. Caribbean represents natural areas with limited human alteration while also considering the naturalness of the surrounding landscape. Examples of human alteration include urban development and intense agricultural use. The degree of naturalness across the landscape is a key ecological condition for sustaining species and ecosystem services that are sensitive to habitat fragmentation at multiple scales. This indicator uses LANDFIRE landcover and ideas from the Florida Critical Lands and Waters Identification Project's approach for evaluating landscape integrity.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

30 miles
7.5 15



- Natural landscape
- Mostly natural landscape
- Partly natural landscape
- Altered landscape
- Heavily altered landscape

Table 10: Indicator values for Caribbean landscape condition within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Natural landscape	30,500	12.2%	
	Mostly natural landscape	35,259	14.1%	↑ In good condition
	Partly natural landscape	18,716	7.5%	↓ Not in good condition
	Altered landscape	3,074	1.2%	
↓ Low	Heavily altered landscape	982	0.4%	
	Area not evaluated for this indicator	*162,380*	*64.7%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Caribbean low-urban historic landscapes

This cultural resource indicator is an index of sites on the National Register of Historic Places and other historic sites surrounded by limited urban development in the U.S. Caribbean. It identifies significant historical places that remain connected to their context in the natural world. This indicator uses LANDFIRE landcover and historic places data from the Puerto Rico State Historic Preservation Office, OpenStreetMap, and the University of the Virgin Islands.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Historic place with nearby low-urban buffer
- Historic place with nearby high-urban buffer
- Not identified as a historic place

Table 11: Indicator values for Caribbean low-urban historic landscapes within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.

	Indicator Values	Acres	Percent of Area
↑ High	Historic place with nearby low-urban buffer	240	<0.1%
	Historic place with nearby high-urban buffer	409	0.2%
↓ Low	Not identified as a historic place	177,409	70.7%
	Area not evaluated for this indicator	*72,853*	*29.0%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Caribbean reforestation potential

This indicator prioritizes areas to increase tree cover in the U.S. Caribbean based on current land uses and potential benefits for local drinking water supplies. It includes opportunities to improve water quality and species habitat by transitioning sun-grown coffee production to shade-grown; enhancing the overstory of shade-grown coffee areas; and reforesting open space in developed areas, pasture, and agricultural lands. The highest scores represent coffee plantations in watersheds with reservoirs. This indicator uses LANDFIRE landcover and U.S. Geological Survey watershed boundaries.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Highest reforestation potential (coffee production in watershed with reservoir)
- Very high reforestation potential (other coffee production)
- High reforestation potential (open space in developed area, pasture, or crops in watershed with reservoir)
- Medium reforestation potential (other open space in developed area, pasture, or crops)
- Low reforestation potential (already natural or developed)

Table 12: Indicator values for Caribbean reforestation potential within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Highest reforestation potential (coffee production in watershed with reservoir)	0	0%
	Very high reforestation potential (other coffee production)	0	0%
	High reforestation potential (open space in developed area, pasture, or crops in watershed with reservoir)	0	0%
	Medium reforestation potential (other open space in developed area, pasture, or crops)	3,056	1.2%
↓ Low	Low reforestation potential (already natural or developed)	175,002	69.7%
	Area not evaluated for this indicator	*72,853*	*29.0%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

Caribbean urban park size

This cultural resource indicator measures the size of parks and beaches in the urban environment in the U.S. Caribbean. Protected natural areas in urban environments provide urban residents a nearby place to connect with nature, and offer refugia for some species. All beaches in this region are open to the public. This indicator uses several protected areas and beach datasets (e.g., the Protected Areas Database of the United States, OpenStreetMap) and U.S. Census Bureau urban areas.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- 75+ acre urban park
- 50 to <75 acre urban park
- 30 to <50 acre urban park
- 10 to <30 acre urban park
- 5 to <10 acre urban park
- <5 acre urban park
- Not identified as an urban park

Table 13: Indicator values for Caribbean urban park size within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	75+ acre urban park	23,357	9.3%
	50 to <75 acre urban park	177	<0.1%
	30 to <50 acre urban park	51	<0.1%
	10 to <30 acre urban park	256	0.1%
	5 to <10 acre urban park	164	<0.1%
	<5 acre urban park	501	0.2%
↓ Low	Not identified as an urban park	226,406	90.2%
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Terrestrial

U.S. Virgin Islands imperiled terrestrial species

This indicator identifies potential habitat to support terrestrial species across the U.S. Virgin Islands. It focuses on species of conservation interest like endangered species and Species of Greatest Conservation Need (SGCN), which are identified in each state or territory's Wildlife Action Plan as most in need of conservation action. The indicator evaluates overall species richness, but places special emphasis on three narrow endemics that are restricted to very small ranges, which are therefore more vulnerable to habitat loss or change. The data originates from WildMon and the U.S. Fish and Wildlife Service Environmental Conservation Online System (ECOS).



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Very high species richness (top 10%) or presence of species with very small ranges
- High species richness (upper 10-30%)
- Medium species richness (above average)
- Low species richness (below average)
- Developed or agriculture

Table 14: Indicator values for U.S. Virgin Islands imperiled terrestrial species within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Very high species richness (top 10%) or presence of species with very small ranges	8,942	3.6%
	High species richness (upper 10-30%)	18,013	7.2%
	Medium species richness (above average)	17,716	7.1%
	Low species richness (below average)	23,324	9.3%
↓ Low	Developed or agriculture	20,584	8.2%
	Area not evaluated for this indicator	*162,332*	*64.7%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater

Caribbean imperiled aquatic species

This indicator measures the number of aquatic animal Species of Greatest Conservation Need (SGCN) observed within each 12-digit HUC subwatershed in Puerto Rico and the U.S. Virgin Islands, including fish, shrimp, crabs, and crayfish. It captures patterns of rare and endemic aquatic species diversity. SGCN are identified in State Wildlife Action Plans as most in need of conservation action. This indicator originates from data compiled by the Southeast Aquatic Resources Partnership from university researchers. It applies to the floodplain predicted to be inundated by a 100-year flood (also known as the 1% annual chance flood), derived from the Federal Emergency Management Agency's National Flood Hazard Layer, and buffered flowlines representing other streams.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Number of aquatic animal Species of Greatest Conservation Need (RSGCN) observed

- 8+ species
- 7 species
- 6 species
- 5 species
- 4 species
- 3 species
- 2 species
- 1 species
- 0 species
- Not identified as a floodplain in surveyed watersheds

Table 15: Indicator values for Caribbean imperiled aquatic species within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Number of aquatic animal Species of Greatest Conservation Need (RSGCN) observed	Acres	Percent of Area
↑ High	8+ species	0	0%
	7 species	0	0%
	6 species	0	0%
	5 species	1,808	0.7%
	4 species	2,046	0.8%
	3 species	2,459	1.0%
	2 species	2,888	1.2%
	1 species	550	0.2%
	0 species	818	0.3%
↓ Low	Not identified as a floodplain in surveyed watersheds	53,866	21.5%
	Area not evaluated for this indicator	*186,477*	*74.3%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater

Caribbean natural landcover in floodplains

This indicator measures the amount of natural landcover in the estimated floodplain of rivers and streams within each catchment in the U.S. Caribbean. It assesses the stream channel and its surrounding riparian buffer, measuring the percent of unaltered habitat like forests, wetlands, or open water (rather than agriculture or development) in the floodplain. Intact vegetated buffers within the floodplain of rivers and streams provide aquatic habitat, improve water quality, reduce erosion and flooding, recharge groundwater, and more. This indicator originates from LANDFIRE land cover. It applies to the floodplain predicted to be inundated by a 100-year flood (also known as the 1% annual chance flood), derived from the Federal Emergency Management Agency's National Flood Hazard Layer, and buffered flowlines representing other streams.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

30 miles
7.5 15



Percent natural landcover within the estimated floodplain, by catchment

- >90% natural landcover
- >80-90% natural landcover
- >70-80% natural landcover
- >60-70% natural landcover
- ≤60% natural landcover
- Not identified as a floodplain

Table 16: Indicator values for Caribbean natural landcover in floodplains within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values: Percent natural landcover within the estimated floodplain, by catchment	Acres	Percent of Area	
↑ High	>90% natural landcover	3,710	1.5%	
	>80-90% natural landcover	2,869	1.1%	↑ In good condition
	>70-80% natural landcover	2,355	0.9%	↓ Not in good condition
	>60-70% natural landcover	1,693	0.7%	
	≤60% natural landcover	4,165	1.7%	
↓ Low	Not identified as a floodplain	69,295	27.6%	
	Area not evaluated for this indicator	*166,823*	*66.5%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater

Caribbean network complexity

This indicator depicts the number of connected stream size classes in a river network between dams or waterfalls in the U.S. Caribbean. River networks with a variety of connected stream classes help retain aquatic biodiversity in a changing climate by allowing species to access climate refugia and move between habitats. This indicator originates from the Southeast Aquatic Resources Partnership. It applies to the estimated floodplain, which spatially defines areas predicted to be inundated by a 100-year flood (also known as the 1% annual chance flood), based on the Federal Emergency Management Agency's National Flood Hazard Layer, and buffered flowlines representing other streams.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Number of connected stream size classes

- 4 size classes
- 3 size classes
- 2 size classes
- 1 size class
- Not identified as a floodplain

Table 17: Indicator values for Caribbean network complexity within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Number of connected stream size classes	Acres	Percent of Area
↑ High	4 size classes	0	0%
	3 size classes	0	0%
	2 size classes	1,458	0.6%
	1 size class	12,333	4.9%
↓ Low	Not identified as a floodplain	70,296	28.0%
	Area not evaluated for this indicator	*166,823*	*66.5%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Freshwater

Caribbean permeable surface

This indicator measures the average percent of non-impervious cover within each catchment in the U.S. Caribbean. High levels of impervious surface degrade water quality and alter freshwater flow, impacting both aquatic species communities and ecosystem services for people, like the availability of clean drinking water. It originates from the National Oceanic and Atmospheric Administration's Coastal Change Analysis Program landcover.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Percent of catchment or small island permeable

- >95% permeable (likely high water quality and supporting most sensitive aquatic species)
- >90-95% permeable (likely declining water quality and supporting most aquatic species)
- >70-90% permeable (likely degraded water quality and not supporting many aquatic species)
- ≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)

Table 18: Indicator values for Caribbean permeable surface within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.

	Indicator Values: Percent of catchment or small island permeable	Acres	Percent of Area	
↑ High	>95% permeable (likely high water quality and supporting most sensitive aquatic species)	2,886	1.2%	↑ In good condition
	>90-95% permeable (likely declining water quality and supporting most aquatic species)	10,172	4.1%	↓ Not in good condition
	>70-90% permeable (likely degraded water quality and not supporting many aquatic species)	68,638	27.4%	
↓ Low	≤70% permeable (likely degraded instream flow, water quality, and aquatic species communities)	7,094	2.8%	
	Area not evaluated for this indicator	*162,120*	*64.6%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Caribbean beach habitat

This indicator evaluates beach habitat for six species of birds and sea turtles that nest on beaches in the U.S. Caribbean (Wilson's plover, American oystercatcher, and hawksbill, leatherback, green, and loggerhead sea turtles). It includes beach locations, sea turtle nest observations, and predicted suitable habitat for birds and sea turtles. Beaches, especially those known to support beach-nesting species, are particularly important habitats due to their limited spatial extent and vulnerability to development and sea level rise. This indicator combines multiple datasets from the Gap Analysis Project, State of the World's Sea Turtles, OpenStreetMap, and more.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Beach with 1+ nesting species predicted or observed
- Other beach
- Not identified as a beach

Table 19: Indicator values for Caribbean beach habitat within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Beach with 1+ nesting species predicted or observed	853	0.3%	↑ In good condition
	Other beach	475	0.2%	↓ Not in good condition
↓ Low	Not identified as a beach	249,582	99.5%	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Caribbean coastal shoreline condition

This indicator assesses shoreline alteration based on the presence of hardened structures like seawalls, groins, and riprap at the dynamic interface between land and water along the U.S. Caribbean coast. By restricting the natural movement of sediment, shoreline armoring increases erosion, prevents the inland migration of coastal ecosystems in response to sea level rise, and degrades habitat for birds, sea turtles, fish, plants, and other species both on and offshore. This indicator originates from the National Oceanic and Atmospheric Administration's Continually Updated Shoreline Product.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Natural

Armored

Table 20: Indicator values for Caribbean coastal shoreline condition within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Natural	4,524	1.8%	↑ In good condition
↓ Low	Armored	305	0.1%	↓ Not in good condition
	Area not evaluated for this indicator	*246,082*	*98.1%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Caribbean fish hotspots

This indicator represents areas of high predicted fish density and diversity in the U.S. Caribbean based on the presence of mangroves, seagrass, and coral in close proximity to one another. Many marine and estuarine fish species use mangroves, seagrass, and coral during different life stages or activities. These habitats provide nursery areas, foraging opportunities, shelter, and protection from predators. This indicator draws from research in Puerto Rico that examines fish density and the number of fish species present at different distances from various habitat types (Pittman et al. 2007). It uses benthic habitat data from The Nature Conservancy and landcover from the National Oceanic and Atmospheric Administration's Coastal Change Analysis Program.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Level of predicted fish density and diversity

- Highest density/diversity (mangrove, coral, and dense seagrass all present within 300 m)
- Very high density/diversity (either mangrove and coral, mangrove and dense seagrass, or coral and dense seagrass present within 300 m)
- High density/diversity (mangrove, coral, and dense seagrass all present within 600 m)
- Medium density/diversity (either mangrove and coral, mangrove and dense seagrass, or coral and dense seagrass present within 600 m)
- Low density/diversity (no coral, mangrove, or dense seagrass present within 600 m of one other)

Table 21: Indicator values for Caribbean fish hotspots within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values: Level of predicted fish density and diversity	Acres	Percent of Area
↑ High	Highest density/diversity (mangrove, coral, and dense seagrass all present within 300 m)	1,201	0.5%
	Very high density/diversity (either mangrove and coral, mangrove and dense seagrass, or coral and dense seagrass present within 300 m)	21,145	8.4%
	High density/diversity (mangrove, coral, and dense seagrass all present within 600 m)	1,724	0.7%
	Medium density/diversity (either mangrove and coral, mangrove and dense seagrass, or coral and dense seagrass present within 600 m)	23,464	9.4%
↓ Low	Low density/diversity (no coral, mangrove, or dense seagrass present within 600 m of one other)	117,863	47.0%
	Area not evaluated for this indicator	*85,514*	*34.1%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Caribbean fish nursery habitat

This indicator represents nursery and spawning habitat or other concentration areas for fish in the U.S. Caribbean. It captures places like mangrove lagoons, bays, estuaries, and some coral reefs. These areas serve as important nursery habitat for many fish species including snook, tarpon, snapper, great barracuda, grunt, mojarra, mullet, jack, bonefish, and more. This data originates from the National Oceanic and Atmospheric Administration's Environmental Sensitivity Index.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Fish spawning, nursery, or other concentration area

Not identified as a fish spawning, nursery, or other concentration area

Table 22: Indicator values for Caribbean fish nursery habitat within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Fish spawning, nursery, or other concentration area	4,437	1.8%
↓ Low	Not identified as a fish spawning, nursery, or other concentration area	246,474	98.2%
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Caribbean mangrove migration

This indicator represents current and potential future locations of mangroves in the U.S. Caribbean based on landcover and barriers to mangrove migration like slope, elevation, and development. Mangroves serve as essential habitat for migratory birds and economically important fish while also providing ecosystem services like stabilizing shorelines, trapping sediment, and protecting coastal communities from storm surge. This indicator adapts methods from the U.S. Virgin Islands Climate Change Ecosystem-Based Adaptation guidance document and expands the analysis to the full U.S. Caribbean. It uses landcover from the National Oceanic and Atmospheric Administration's Coastal Change Analysis Program as well as slope, elevation, roads, and landcover from LANDFIRE.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Current mangrove
- Potential future mangrove <0.5 km from current mangrove
- Potential future mangrove 0.5-1 km from current mangrove
- Potential future mangrove >1 km from current mangrove
- Not likely to be current or future mangrove

Table 23: Indicator values for Caribbean mangrove migration within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Current mangrove	268	0.1%
	Potential future mangrove <0.5 km from current mangrove	571	0.2%
	Potential future mangrove 0.5-1 km from current mangrove	112	<0.1%
	Potential future mangrove >1 km from current mangrove	390	0.2%
↓ Low	Not likely to be current or future mangrove	84,517	33.7%
	Area not evaluated for this indicator	*165,053*	*65.8%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Caribbean seagrass

This indicator represents the presence and density of seagrass at various depths in the U.S. Caribbean. Seagrasses provide food and habitat for a range of marine and estuarine wildlife. They also produce oxygen, filter water, sequester carbon, control erosion, and buffer storms. Seagrasses serve as an important indicator of the overall health of coastal ecosystems because they are sensitive to water quality and require sufficiently clear water for sunlight to penetrate. This indicator uses benthic habitat data from The Nature Conservancy and bathymetry data from the National Oceanic and Atmospheric Administration.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Very shallow and dense seagrass
- Very shallow and sparse seagrass
- Shallow and dense seagrass
- Shallow and sparse seagrass
- Deep and dense seagrass
- Deep and sparse seagrass
- Not seagrass

Table 24: Indicator values for Caribbean seagrass within United States Virgin Islands. A good condition threshold is not yet defined for this indicator.


	Indicator Values	Acres	Percent of Area
↑ High	Very shallow and dense seagrass	3,581	1.4%
	Very shallow and sparse seagrass	4,365	1.7%
	Shallow and dense seagrass	1,878	0.7%
	Shallow and sparse seagrass	3,963	1.6%
	Deep and dense seagrass	951	0.4%
	Deep and sparse seagrass	2,395	1.0%
↓ Low	Not seagrass	147,334	58.7%
	Area not evaluated for this indicator	*86,445*	*34.5%*
	Total area	**250,911**	**100%**

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

Coastal & marine

Caribbean shallow hardbottom & coral

This indicator measures the presence of hardbottom habitat and coral in the U.S. Caribbean. It also predicts the ability of coral to survive the impacts of climate change based on reef locations, past and future thermal conditions, hurricane impacts, and coral larval connectivity. Hardbottom and coral serve as important habitat for many marine species and provide economic and cultural benefits to nearby coastal communities, such as supporting fisheries, filtering seawater, and buffering the impacts of storms. This indicator combines benthic habitat and coral climate refugia data from The Nature Conservancy.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



- Coral with above average climate resilience
- Coral with unknown climate resilience
- Coral with below average climate resilience
- Hardbottom with sparse algae
- Hardbottom with dense algae
- Not coral or hardbottom

Table 25: Indicator values for Caribbean shallow hardbottom & coral within United States Virgin Islands. Good condition thresholds reflect the range of indicator values that occur in healthy, functioning ecosystems.


	Indicator Values	Acres	Percent of Area	
↑ High	Coral with above average climate resilience	10,923	4.4%	
	Coral with unknown climate resilience	890	0.4%	
	Coral with below average climate resilience	6,447	2.6%	↑ In good condition
	Hardbottom with sparse algae	29,486	11.8%	↓ Not in good condition
	Hardbottom with dense algae	43,542	17.4%	
↓ Low	Not coral or hardbottom	74,108	29.5%	
	Area not evaluated for this indicator	*85,514*	*34.1%*	
	Total area	**250,911**	**100%**	

To learn more and explore the GIS data, view this indicator in the SECAS Atlas.

More Information

Protected Areas

Protected areas include a variety of public lands owned or managed by federal, state, and local agencies and nonprofit organizations, as well as some private conservation lands and conservation easements. Protected areas are derived from the Protected Areas Database of the United States (PAD-US v4.1) and include Fee, Designation, Easement, Marine, and Proclamation (Dept. of Defense lands only) boundaries.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

7.5 15 30 miles



Within a protected area

Not within a protected area

Table 26: Extent of protected areas within United States Virgin Islands.

Protected area status	Acres	Percent of Area
Not within a protected area	192,985	76.9%
Within a protected area	57,926	23.1%
Total area	**250,911**	**100%**

Protected areas at this location:


- St. Croix East End Marine Park (26,056 acres)
- Virgin Islands National Park (14,763 acres)
- Virgin Islands National Park (National Park Service; 13,154 acres)
- Buck Island Reef National Monument (5,979 acres)
- Virgin Islands Coral Reef National Monument (National Park Service; 4,393 acres)
- Virgin Islands Coral Reef National Monument (4,393 acres)
- St. Thomas East End Reserves (2,297 acres)
- St. James Marine Reserve and Wildlife Sanctuary (1,734 acres)
- Salt River Bay National Historic Park and Ecological Preserve (896 acres)
- Buck Island Reef National Monument (National Park Service; 878 acres)
- Cas Cay-Mangrove Lagoon Marine Reserve & Wildlife Sanctuary (729 acres)
- Sandy Point National Wildlife Refuge (US Fish and Wildlife Service; 518 acres)
- Sandy Point National Wildlife Refuge (518 acres)
- Salt River Bay National Historical Park and Ecological Preserve (National Park Service; 238 acres)
- Estate Clairmont (Land Trust; 236 acres)
- Savana Island (US Virgin Islands Government; 175 acres)
- Estate Thomas EF Experimental Area (USDA Forest Service; 140 acres)
- Estate Thomas Ef Experimental Area (140 acres)
- Estate Thomas Experimental Forest (140 acres)
- East End (US Virgin Islands Government; 136 acres)
- East Bay and Point Udall (US Virgin Islands Government; 134 acres)
- Butler Bay (Land Trust; 115 acres)
- Magen's Bay (US Virgin Islands Government; 111 acres)
- Outer Brass Island (US Virgin Islands Government; 108 acres)
- Southgate (Land Trust; 104 acres)
- ... and 44 more protected areas ...

Note: areas are listed based on name, ownership, and boundary information in the Protected Areas Database of the United States, which may include overlapping and duplicate areas. Only protected areas with at least 1 acre of overlap with this area are included in the list above.

Sea Level Rise

NOAA's sea level rise (SLR) inundation models represent areas likely to experience flooding at high tide based on each foot of SLR above current levels. Darker blue areas will experience flooding first, and at greater depth, compared to lighter blue areas. These models are not linked to a future timeframe; see the projections below. NOAA calculates the inundation footprint at "mean higher high water", or the average highest daily tide. The area covered in each SLR scenario includes areas projected to be inundated at lower levels. For example, the area inundated by 4 ft of SLR also includes areas inundated by 3 ft, 2 ft, 1 ft, and 0 ft of SLR (where 0 ft represents current levels).

To explore additional SLR information, please see NOAA's Sea Level Rise Viewer.



Basemap credits: © Mapbox © OpenStreetMap Improve this basemap

30 miles
7.5 15



Flooding extent by projected sea level rise (ft)

0 1 2 3 4 5 6 7 8 9 10

Table 27: Extent of flooding by projected average highest daily tide due to sea level rise within United States Virgin Islands. Values from the NOAA sea level rise inundation data.

Feet of sea level rise	Acres	Percent of Area
0 feet	153,619	61.2%
1 foot	154,229	61.5%
2 feet	154,647	61.6%
3 feet	155,365	61.9%
4 feet	156,105	62.2%
5 feet	156,690	62.4%
6 feet	157,419	62.7%
7 feet	157,967	63.0%
8 feet	158,452	63.2%
9 feet	158,880	63.3%
10 feet	159,299	63.5%
Not projected to be inundated by up to 10 feet	*80,501*	*32.1%*
No sea level rise data available	*11,111*	*4.4%*
Total area	**250,911**	**100%**

Table 28: Projected sea level rise by decade within United States Virgin Islands. Values are based on area-weighted averages of decadal projections for 1-degree grid cells that overlap this area based on NOAA's 2022 Sea Level Rise Report. 2060 corresponds to the SECAS goal: a 10% or greater improvement in the health, function, and connectivity of Southeastern ecosystems by 2060.

SLR Scenario	2020 (ft)	2030 (ft)	2040 (ft)	2050 (ft)	2060 (ft)	2070 (ft)	2080 (ft)	2090 (ft)	2100 (ft)
Low	0.23	0.36	0.49	0.62	0.75	0.85	0.95	1.1	1.2
Intermediate-low	0.26	0.39	0.56	0.75	0.95	1.2	1.4	1.6	1.8
Intermediate	0.26	0.43	0.62	0.89	1.2	1.5	2	2.6	3.4
Intermediate-high	0.26	0.46	0.72	1.1	1.6	2.3	3.2	4	5
High	0.26	0.46	0.82	1.3	2	3	4.2	5.4	6.8

Urban growth

Projected future urbanization data is not currently available for this area.

Wildfire Likelihood

Wildfire likelihood data is not currently available for this area.