

# Trend Analysis of Uncensored Major Ions

Dave Lorenz

December 4, 2015

## Abstract

This example illustrates the data manipulations for the seasonal Kendall analysis of uncensored data. Major ions are typically uncensored in natural waters and provide a useful example. This example also uses a common time frame for all of the trend tests. The common time frame facilitates comparing trends among the stations. Most often users will want to divide trend analyses into similar groups of analytes like major ions, nutrients and so forth because they will be analyzed in similar ways and will have common sampling time frames.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Summarize the Sample Data</b>	<b>3</b>
<b>3</b>	<b>Set up the Project</b>	<b>6</b>
<b>4</b>	<b>Flow Adjustment</b>	<b>8</b>
<b>5</b>	<b>Seasonal Kendall Trend Test</b>	<b>13</b>
<b>6</b>	<b>Trend Results</b>	<b>15</b>
<b>7</b>	<b>Further Remarks</b>	<b>18</b>

# 1 Introduction

The data used in this application are a small subset of the data used by Schertz and others (1991). The data are samples taken from water year 1969 (October, 1968) through water year 1989 (September, 1989). Nineteen stations were selected and only calcium and chloride were selected for the major ions. The data were modified by removing the remark columns associated with those constituents to make the analysis more straightforward.

```
> # Load the restrend and smwrBase packages and the data
> library(restrend)
> library(smwrBase)
> data(EstrendSub)
> head(EstrendSub)
```

	STAID	DATES	QI	QD	RN.organic	PN.organic	RAmmonia	PAmmonia	RKjeldahl	PKjeldahl
1	07227500	1968-10-01	7.6	NA			NA	NA		NA
2	07227500	1968-10-03	5.3	NA			NA	NA		NA
3	07227500	1968-10-16	532.0	NA			NA	NA		NA
4	07227500	1968-10-19	17.0	NA			NA	NA		NA
5	07227500	1968-11-01	17.0	NA			NA	NA		NA
6	07227500	1968-12-01	6.6	NA			NA	NA		NA

	RTotal.P	PTotal.P	RCopper	PCopper	RIron	PIron	Calcium	Chloride
1		NA		NA		NA	95	280
2		NA		NA		NA	NA	NA
3		NA		NA		NA	42	106
4		NA		NA		NA	121	435
5		NA		NA		NA	150	512
6		NA		NA		NA	138	510

## 2 Summarize the Sample Data

In general, it is desirable, but not necessary, to subset the data before proceeding with the analysis of a subset of the constituents. Before these data are subsetted, the `FLOW` column must be created. The flow data are in two columns `QI`, the flow at the time of the sample; and `QD`, the mean flow on the day of the sample. The `coalesce` function in the `smwrBase` package can be used to select the non-missing value for flow.


```
> # Compute FLOW, the coalesce function is in smwrBase
> EstrendSub <- transform(EstrendSub, FLOW=coalesce(QI, QD))
> # Create the subset
> Majors <- subset(EstrendSub, select=c("STAID", "DATES", "FLOW",
+   "Calcium", "Chloride"))
```

The `sampReport` function creates a simple PDF file that contains a report of the sample date ranges and graph of samples for each site. It can be used to help define the starting and ending date ranges for the trend tests as well as identifying sample gaps and other sampling issues.

```
> # Create the report
> sampReport(Majors, DATES="DATES", STAID="STAID", file="MajorIonSampling")
```

The call to `sampReport` returns the file name invisibly (`MajorIonSampling.pdf`). Because it is a full-size portrait PDF file, it is inserted here with compressed pages. The report gives the actual begin and end dates for sampling and the graph shows the sampling dates for each station. It is easy to see that 5 stations (07233500, 07299200, 07299540, 07336800, and 07346045) were not sampled for the entire retrieval period.

	STAID	FirstSamp	LastSamp	NumSamp
1	07227500	1968-10-01	1989-08-15	350
2	07228000	1968-11-21	1989-08-16	198
3	07233500	1968-11-21	1979-07-24	62
4	07297910	1968-10-01	1989-08-15	308
5	07299200	1968-10-01	1980-07-09	217
6	07299540	1968-10-01	1986-08-25	276
7	07300000	1968-10-01	1989-08-16	283
8	07308500	1968-10-01	1989-08-18	336
9	07316000	1968-10-01	1989-08-16	694
10	07331600	1968-10-01	1989-08-15	228
11	07336800	1968-10-07	1981-09-03	69
12	07336820	1968-10-01	1989-08-15	393
13	07342500	1968-10-01	1989-06-19	325
14	07343000	1968-10-01	1989-06-19	317
15	07343200	1968-10-01	1989-08-16	366
16	07343500	1968-10-01	1989-06-21	303
17	07344500	1968-10-01	1989-08-07	294
18	07346045	1968-10-03	1981-08-27	109
19	07346070	1968-10-01	1989-08-10	300



### 3 Set up the Project

The user must balance the need to include as many stations as possible and the targeted time frame for the trend estimation. For these data, 5 stations have incomplete record, but to include all of those stations, the analysis period would need to be much shorter, though water year 1978. This example will use the full retrieval period.

The `(setProj)` function sets up the trend estimation project. There are many arguments to `(setProj)`, see the documentation for details. The constituent names or response variable names are referred to as **Snames** in keeping with the names used in the original ESTREND.

After projects have been set up, the user can get a list of the projects by using `lsProj` or can specify a project to use with `useProj`. The function `useProj` must be used to continue working on a project after the user quits from the R session.

```
> # Set up the project
> setProj("majors", Majors, STAID="STAID", DATES="DATES",
+         Snames=c("Calcium", "Chloride"), FLOW="FLOW",
+         type="seasonal", Start="1968-10-01", End="1989-10-01")

[1] "majors"
```

The `(setProj)` function creates a folder in the users workspace with that name. That folder contains R data that are updated after each successful call to an analysis function in `restrend`. Table 1 describes the data created in this example's call to `(setProj)`. Any object of class "matrix" or "by" are indexed by station and sname.

**Table 1.** The data created by `(setProj)`.

Name	Class	Description
estrend.cl	list	A record of the calls to analysis functions.
estrend.cn	matrix	A description of the censoring. May be "none," "left," or "multiple."
estrend.cp	matrix	The percent of observations that are left-censored.
estrend.df	by	The dataset, contains STAID, DATES, FLOW, and the response variable.
estrend.in	list	Information about the project, such as the start and end dates and the names of columns in each dataset.
estrend.sl	by	Details from the seasonal selection process. Each is a list from the potential comparisons from 12, 6, 4, and 3 seasons per year definition. See Lorenz (2014) for details.
estrend.ss	matrix	The "best" seasonal definition from the analysis recorded in <code>estrend.sl</code> .
estrend.st	matrix	The status for each station and sname. Must be "OK" to continue with the trend analysis.

It is useful to verify which stations and snames will be analyzed and what the seasonal definitions are. The user need only enter the name of the R data object in the console. For these data, the seasonal definition is 0 in all cases where the status is not "OK."

```
> # Which are OK?
> estrend.st
```

	snames	
stations	Calcium	Chloride
07227500	"OK"	"OK"
07228000	"OK"	"OK"
07233500	"short record"	"short record"
07297910	"OK"	"OK"
07299200	"short record"	"short record"
07299540	"short record"	"short record"
07300000	"OK"	"OK"
07308500	"OK"	"OK"
07316000	"OK"	"OK"
07331600	"OK"	"OK"
07336800	"short record"	"short record"
07336820	"OK"	"OK"
07342500	"OK"	"OK"
07343000	"OK"	"OK"
07343200	"OK"	"OK"
07343500	"OK"	"OK"
07344500	"OK"	"OK"
07346045	"short record"	"short record"
07346070	"OK"	"OK"

```
> # What seasonal definition?
> estrend.ss
```

	snames	
stations	Calcium	Chloride
07227500	6	6
07228000	6	6
07233500	0	0
07297910	6	6
07299200	0	0
07299540	0	0
07300000	6	6
07308500	6	6
07316000	12	12
07331600	12	12
07336800	0	0
07336820	12	12
07342500	12	12
07343000	12	12
07343200	12	12
07343500	12	12
07344500	12	12
07346045	0	0
07346070	12	12

## 4 Flow Adjustment

Computing flow-adjusted concentrations (flow adjustment) is an optional step in seasonal Kendall trend analysis. It is only appropriate for uncensored or slightly censored data. If the data are censored, the censoring is ignored and the values are taken as the detection limit. Flow adjustment is performed using the `flowAdjust` function and can immediately follow the call to `setProj`.



By default, all stations and snames are flow adjusted by `flowAdjust`. But specific combinations can be separately adjusted, using a different span for the LOWESS procedure for example, or if no satisfactory fit can be found, selected combinations can be completely undone by the `undoFA` function. Note that no relation between flow and concentration is not necessarily an unsatisfactory fit.



The `flowAdjust` function creates a PDF report, and returns the name of the report. The report shows graphs of flow and concentration by station on each page. Up to 6 combinations are shown on a page. For any seasonal Kendall trend test with flow adjustment, the user should review all flow-concentration relation. Only 2 pages are shown in this example, the first illustrates an acceptable fit and the second a marginal fit. The user may choose to revise other flow adjustments or accept all flow adjustments, but those for station 07331600 were selected to demonstrate customized flow adjustment.

```
> # Do the flow adjustment accepting all defaults
> flowAdjust()
```

```
[1] "majors_fa.pdf"
```









The revised fit shows an improved, more smooth fit to the data.

```
> # Do the flow adjustment accepting all defaults  
> flowAdjust(Station="07331600", Snames=c("Calcium", "Chloride"), span=1)  
  
[1] "majors_fa_01.pdf"
```



## 5 Seasonal Kendall Trend Test

After the optional flow-adjustment, these data are ready for the seasonal Kendall trend test. The function **SKTrends** executes the trend test on all valid combinations of stations and snames. It can also execute the test on subsets if some changes need to be made. An important argument is **nseas**, which can be used to force all analyses to use the same seasonal definition. This is essential for the regional seasonal Kendall test and an important consideration for other regional assessments because it levels the playing field for determining significant trends.

The **SKTrends** function also creates a PDF file that contains the result of the analysis and a series graph on each page. See the documentation for **seriesPlot** for information about that graph. The file reports the results for each sname by station with the flow-adjusted results following the untransformed results. Most trends are very small for these data; only the reports for Calcium at 07228000 is shown.


```
> # Trend tests, accepting default seasons
> SKTrends()
```

```
[1] "majors_sk.pdf"
```

07228000 Calcium

Seasonal Kendall with correlation correction

data: log(Calcium) (21 years and 6 seasons)  
tau = 0.41513, p-value = 0.0003365  
alternative hypothesis: true slope is not equal to 0  
sample estimates:  
slope median.data median.time  
0.02154697 4.70048046 10.50000000



## 6 Trend Results

When completed, or to check on intermediate results, the estimated trends can be extracted using the `getTrends` function. By default, all stations and snames are extracted. The output dataset is explained in the documentation for `getTrends`. The user has the option to set a significance level to determine whether there is a significant trend, the default level is 0.05.

```
> # get the trends
> majors.tnd <- getTrends()
> print(majors.tnd)
```

	Station	Response		Type	NumYears	NumSeas	Nobs	RepValue	
1	07227500	Calcium	uncensored	seasonal	Kendall	21	6	123	100.00001
2	07227500	Calcium	flow-adjusted	seasonal	Kendall	21	6	123	100.00001
3	07227500	Chloride	uncensored	seasonal	Kendall	21	6	124	469.89363
4	07227500	Chloride	flow-adjusted	seasonal	Kendall	21	6	124	469.89363
5	07228000	Calcium	uncensored	seasonal	Kendall	21	6	111	110.00001
6	07228000	Calcium	flow-adjusted	seasonal	Kendall	21	6	111	110.00001
7	07228000	Chloride	uncensored	seasonal	Kendall	21	6	111	660.00008
8	07228000	Chloride	flow-adjusted	seasonal	Kendall	21	6	111	660.00008
9	07297910	Calcium	uncensored	seasonal	Kendall	21	6	122	530.00011
10	07297910	Calcium	flow-adjusted	seasonal	Kendall	21	6	122	530.00011
11	07297910	Chloride	uncensored	seasonal	Kendall	21	6	122	3000.00035
12	07297910	Chloride	flow-adjusted	seasonal	Kendall	21	6	122	3000.00035
13	07300000	Calcium	uncensored	seasonal	Kendall	21	6	123	480.00003
14	07300000	Calcium	flow-adjusted	seasonal	Kendall	21	6	123	480.00003
15	07300000	Chloride	uncensored	seasonal	Kendall	21	6	123	270.00002
16	07300000	Chloride	flow-adjusted	seasonal	Kendall	21	6	123	270.00002
17	07308500	Calcium	uncensored	seasonal	Kendall	21	6	119	340.00001
18	07308500	Calcium	flow-adjusted	seasonal	Kendall	21	6	119	340.00001
19	07308500	Chloride	uncensored	seasonal	Kendall	21	6	119	1837.99988
20	07308500	Chloride	flow-adjusted	seasonal	Kendall	21	6	119	1837.99988
21	07316000	Calcium	uncensored	seasonal	Kendall	21	12	182	200.00001
22	07316000	Calcium	flow-adjusted	seasonal	Kendall	21	12	182	200.00001
23	07316000	Chloride	uncensored	seasonal	Kendall	21	12	188	1000.00010
24	07316000	Chloride	flow-adjusted	seasonal	Kendall	21	12	188	1000.00010
25	07331600	Calcium	uncensored	seasonal	Kendall	21	12	200	88.49858
26	07331600	Calcium	flow-adjusted	seasonal	Kendall	21	12	200	88.49858
27	07331600	Chloride	uncensored	seasonal	Kendall	21	12	203	320.00001
28	07331600	Chloride	flow-adjusted	seasonal	Kendall	21	12	203	320.00001
29	07336820	Calcium	uncensored	seasonal	Kendall	21	12	185	65.00001
30	07336820	Calcium	flow-adjusted	seasonal	Kendall	21	12	185	65.00001
31	07336820	Chloride	uncensored	seasonal	Kendall	21	12	188	166.49328
32	07336820	Chloride	flow-adjusted	seasonal	Kendall	21	12	188	166.49328
33	07342500	Calcium	uncensored	seasonal	Kendall	21	12	183	40.00000
34	07342500	Calcium	flow-adjusted	seasonal	Kendall	21	12	183	40.00000
35	07342500	Chloride	uncensored	seasonal	Kendall	21	12	183	16.00000
36	07342500	Chloride	flow-adjusted	seasonal	Kendall	21	12	183	16.00000
37	07343000	Calcium	uncensored	seasonal	Kendall	21	12	170	79.49842
38	07343000	Calcium	flow-adjusted	seasonal	Kendall	21	12	170	79.49842

39	07343000	Chloride	uncensored	seasonal	Kendall	21	12	170	35.00000
40	07343000	Chloride	flow-adjusted	seasonal	Kendall	21	12	170	35.00000
41	07343200	Calcium	uncensored	seasonal	Kendall	21	12	184	55.99999
42	07343200	Calcium	flow-adjusted	seasonal	Kendall	21	12	184	55.99999
43	07343200	Chloride	uncensored	seasonal	Kendall	21	12	184	21.00000
44	07343200	Chloride	flow-adjusted	seasonal	Kendall	21	12	184	21.00000
45	07343500	Calcium	uncensored	seasonal	Kendall	21	12	179	14.00000
46	07343500	Calcium	flow-adjusted	seasonal	Kendall	21	12	179	14.00000
47	07343500	Chloride	uncensored	seasonal	Kendall	21	12	179	20.00000
48	07343500	Chloride	flow-adjusted	seasonal	Kendall	21	12	179	20.00000
49	07344500	Calcium	uncensored	seasonal	Kendall	21	12	186	16.00000
50	07344500	Calcium	flow-adjusted	seasonal	Kendall	21	12	186	16.00000
51	07344500	Chloride	uncensored	seasonal	Kendall	21	12	186	44.49719
52	07344500	Chloride	flow-adjusted	seasonal	Kendall	21	12	186	44.49719
53	07346070	Calcium	uncensored	seasonal	Kendall	21	12	183	7.00000
54	07346070	Calcium	flow-adjusted	seasonal	Kendall	21	12	183	7.00000
55	07346070	Chloride	uncensored	seasonal	Kendall	21	12	183	30.00000
56	07346070	Chloride	flow-adjusted	seasonal	Kendall	21	12	183	30.00000
		Trend	Trend.pct	P.value	Trend.dir				
1	0.00000000	0.000000000	0.8336014152	*					
2	-0.02836618	-0.028366178	0.9601243138	none					
3	12.79590098	2.723148421	0.0336792469	up					
4	12.64550036	2.691141044	0.0129020214	up					
5	2.39588639	2.178078333	0.0003365278	up					
6	1.28918026	1.171981945	0.0130635500	up					
7	17.25881435	2.614971565	0.0007134676	up					
8	11.59983333	1.757550299	0.0029419661	up					
9	1.38744659	0.261782321	0.8355770111	none					
10	3.99106379	0.753030747	0.2728573084	none					
11	30.77328679	1.025776108	0.8205802441	none					
12	81.97615333	2.732538128	0.0758367777	none					
13	0.00000000	0.000000000	0.8724372387	*					
14	-0.01634082	-0.003404338	0.9618166685	none					
15	2.26912466	0.840416463	0.0046260357	up					
16	2.25459597	0.835035468	0.0180431604	up					
17	-0.87151697	-0.256328516	0.6151251197	none					
18	-1.02469247	-0.301380133	0.5201364756	none					
19	8.02723876	0.436737719	0.5162076950	none					
20	-3.45285768	-0.187859516	0.8067187071	none					
21	0.69571531	0.347857631	0.5388302803	none					
22	2.11482920	1.057414530	0.0956373215	none					
23	9.27514624	0.927514535	0.3566403389	none					
24	18.17821565	1.817821390	0.0474613905	up					
25	0.62565134	0.706962055	0.2394109964	none					
26	0.38056581	0.430024789	0.3169723749	none					
27	2.57714511	0.805357813	0.5003621578	none					
28	0.69401673	0.216880221	0.7803473473	none					
29	0.84057835	1.293197225	0.0961654186	none					
30	0.66254212	1.019295374	0.1187556982	none					
31	4.10125215	2.463313924	0.0698596239	none					
32	3.75668736	2.256359751	0.0728764534	none					



33	-0.35416765	-0.885419081	0.2286658287	none
34	-0.30449254	-0.761231313	0.1911620051	none
35	-0.31416803	-1.963550197	0.1585604995	none
36	-0.16540169	-1.033760534	0.2915501893	none
37	0.20723170	0.260673977	0.6322352886	none
38	-0.18962446	-0.238526065	0.6531360745	none
39	-0.16173531	-0.462100829	0.7745313048	none
40	-0.57363605	-1.638959992	0.0628329962	none
41	-0.40639101	-0.725698359	0.2162646502	none
42	-0.56856580	-1.015296249	0.0369244702	down
43	-0.58466480	-2.784117838	0.0263182856	down
44	-0.53644449	-2.554497343	0.0054448307	down
45	-0.07162687	-0.511620472	0.5025991797	none
46	-0.05764936	-0.411781115	0.3829599321	none
47	-0.32744925	-1.637246215	0.2295798212	none
48	-0.32282807	-1.614140285	0.0948637649	none
49	-0.09071358	-0.566959859	0.2795946300	none
50	-0.12661440	-0.791340014	0.0314434171	down
51	-0.49735710	-1.117726880	0.3637532592	none
52	-0.94332415	-2.119963232	0.0002551555	down
53	-0.04453901	-0.636271629	0.0876174122	none
54	-0.05656804	-0.808114896	0.0062053259	down
55	-0.70519603	-2.350653317	0.0265707411	down
56	-0.71787733	-2.392924312	0.0005073280	down

## 7 Further Remarks

Because trend analysis is not necessarily a straightforward process, but requires user assessments at several points in the process, it is not necessarily a good idea to simply create scripts and run them without any user review and interaction. To overcome recording the steps in a script, the functions in `restrend` record all changes to the projects database in a list called `estrend.cl`. It can be viewed at any time simply by entering `estrend.cl` in the console window. It can be saved with the data to ensure that the trend analysis is reproducible.

```
> # get the history
> estrend.cl
```

```
[[1]]
setProj(project = "majors", data = Majors, STAID = "STAID", DATES = "DATES",
        Snames = c("Calcium", "Chloride"), FLOW = "FLOW", type = "seasonal",
        Start = "1968-10-01", End = "1989-10-01")
```

```
[[2]]
flowAdjust()
```

```
[[3]]
flowAdjust(Stations = "07331600", Snames = c("Calcium", "Chloride"),
           span = 1)
```

```
[[4]]
SKTrends()
```

## References

- [1] Lorenz, D.L., in preparation, `restrend`: an R package for ESTimate TRENDS: U.S. Geological Survey Open File Report, ? p.
- [2] Schertz, T.L., Alexander, R.B., and Ohe, D.J., 1991, The computer program ESTimate TREND (ESTREND), a system for the detection of trends in water-quality data: U.S. Geological Survey Water Resources Investigations Report 91-4040, 72 p.