

INTERIM USE LIMITATION AREA FOR THE CALIFORNIA TIGER SALAMANDER AND ALAMEDA WHIPSNAKE

Within this area, follow the indicated instruction(s) for the active ingredient(s) you intend to use. If multiple instructions apply to your pesticide use, you should follow all of the instructions.

PESTICIDE	INSTRUCTION
2,4-D	B
acephate	A
acrolein	B
aluminum phosphide	A and B
bensulide	A
beta-cyfluthrin	A
bifenthrin	A
brodifacoum	B
bromadiolone	B
bromethalin	B
chlorophacinone	A and B
chlorothalonil	A
chlorpyrifos	A
cyfluthrin	A
cyhalothrin (lambda)	A
cypermethrin	A
deltamethrin	A
difenacoum	B
difethialone	B
dimethoate	A
diphacinone	A and B
endosulfan	A
esfenvalerate	A
ethoprop	A
fenpropathrin	A
fipronil	A

PESTICIDE	INSTRUCTION
fluvalinate	A
imidacloprid	A
magnesium phosphide	B
malathion	A
mancozeb	A
maneb	A
metam sodium	A
methomyl	A
methoprene	A
methyl bromide	A
oryzalin	A
oxydemeton-methyl	A
oxyfluorfen	A
PCNB	A
phenothrin	A
phosmet	A
potassium nitrate	B
propargite	A
resmethrin	A
S-metolachlor	A
sodium nitrate	B
strychnine	A and B
tetramethrin	A
warfarin	B
zeta-cypermethrin	A
zinc phosphide	B

INSTRUCTION A –Do not apply the pesticide within 200 feet if applied by ground nor within 400 feet if applied by air, of fresh-water ponds (including natural or manmade, such as stock ponds), slow moving streams or pools within streams, vernal pools and other ephemeral or permanent water bodies which typically support inundation during winter rains and hold water for a minimum of 12 weeks in a year of average rainfall.

INSTRUCTION B - Do not apply the pesticide within 100 feet if applied by ground nor within 400 feet if applied by air, of areas described as *shrub communities* known as chaparral consisting of northern coastal sage scrub and coastal sage. Also do not apply the pesticide in open grasslands, oak savanna, and oak-bay woodland occurring within 500 feet of the shrub communities described above.