

INTERIM USE LIMITATION AREA FOR THE SAN FRANCISCO GARTER SNAKE AND TIDEWATER GOBY

Within this area, follow the indicated instruction(s) for the active ingredient(s) you intend to use. If multiple instructions apply to your pesticide use, you should follow all of the instructions.

| PESTICIDE | INSTRUCTION |
|----------------------|-------------|
| acephate | A |
| aluminum phosphide | A |
| bensulide | A and B |
| beta-cyfluthrin | A and B |
| bifenthrin | A and B |
| chlorothalonil | A and B |
| chlorpyrifos | A |
| cyfluthrin | A and B |
| cyhalothrin (lambda) | A and B |
| cypermethrin | A and B |
| deltamethrin | A and B |
| diazinon | B |
| dimethoate | A and B |
| endosulfan | A |
| esfenvalerate | A and B |
| ethoprop | A and B |
| fenpropathrin | A and B |
| fipronil | A and B |

| PESTICIDE | INSTRUCTION |
|---------------------|-------------|
| fluvalinate | A and B |
| imidacloprid | A and B |
| magnesium phosphide | A |
| methomyl | A and B |
| naled | A |
| oxydemeton-methyl | A and B |
| oxyfluorfen | A and B |
| PCNB | A and B |
| permethrin | A |
| phenothrin | A and B |
| potassium nitrate | A |
| propargite | A and B |
| resmethrin | A and B |
| sodium nitrate | A |
| strychnine | A |
| tetramethrin | A and B |
| trifluralin | A |
| zeta-cypermethrin | A and B |

INSTRUCTION A – Do not apply the pesticide within 200 feet if applied by ground nor within 400 feet if applied by air, of fresh-water ponds (including natural or manmade, such as stock ponds), slow moving streams or pools within streams, vernal pools and other ephemeral or permanent water bodies which typically support inundation during winter rains and hold water for a minimum of 12 weeks in a year of average rainfall.

INSTRUCTION B - Do not apply the pesticide within 100 feet if applied by ground nor within 400 feet if applied by air, of lagoons, estuaries and backwater marshes that are adjacent to the Pacific Ocean; and freshwater streams up-gradient of and tributary to brackish habitats.