
(a) Pre-specified (frozen code)
Names within areas (primary, H2)
Same, corrected code (Norwegian names read)
Lexicon only (N-lex)
Character model only (N-learn)
Placebo: names reassigned
Position only, within areas (H1)
Position only, flexible models (H1, descriptive)
Total incl. regional composition (secondary)
Flexible model refitted with names (secondary)
Countries held out
Time held out (floods from 2024) *
Language model, floods after its cut-off (H4)
Lexicon and character model, same floods
Same, flexible models refitted (secondary)
Language model incl. its composition (secondary)
R 0.15, to 0.36
to 0.15
to 0.14
R 0.17, to 0.33
(b) Exploratory checks
Second stage refitted in each bootstrap replicate
Flood events held out whole (frozen code)
Flood events held out whole
Flood traces counted as not flooded
Flooded share over land only
Partly flooded cells excluded
Every AOI weighted equally (frozen code)
European AOIs alone, own yardstick
Without the Americas and Oceania
Strict reading
Gazetteer cleaned of city sections and duplicates
Without AOIs lacking a hydrography layer
Without AOIs lacking a flood type
Language-model reading, primary split
Floodplain reading (riverine and wetland classes)
Cleaned lexicon (two-letter affixes and English suffixes removed)
GFM labels, existing predictions rescored (frozen code)
(c) Ranges, not intervals
13 assignments of location groups to folds
Names reassigned 200 times, real composition
Names reassigned 200 times, composition recomputed
−0.075
−0.050
−0.025
0.000
0.025
0.050
0.075
R (improvement from names / improvement from terrain)