


# CH 37: Communities and ecosystems

李承叡


生態學與演化生物學研究所

生命科學館 1129

## Chapter 37: Big Ideas



**Community Structure  
and Dynamics**



**Ecosystem Structure  
and Dynamics**

# COMMUNITY STRUCTURE AND DYNAMICS

## 37.1 A community includes all the organisms inhabiting a particular area

- Biological **community**: an assemblage of all the populations of organisms living close enough together for potential interaction.
- Ecologists define the boundaries of the community according to the research questions they want to investigate.

TABLE 37.2 |    INTERSPECIFIC INTERACTIONS

Interspecific Interaction	Effect on Species 1	Effect on Species 2	Example
Competition	—	—	Squirrels/ black bears
Mutualism	+	+	Plants/ mycorrhizae
Predation	+	—	Crocodiles/fish
Herbivory	+	—	Caterpillars/leaves
Parasites and pathogens	+	—	Heartworm/dogs; <i>Salmonella</i> / humans

## 37.2 Interspecific interactions are fundamental to community structure

- **Interspecific competition** occurs when two different species compete for the same limited resource.
- In **mutualism**, both populations benefit.
- In **predation**, one species (the predator) kills and eats another (the prey).
- In **herbivory**, an animal consumes plant parts or algae.
- In **parasitism**, the host plants or animals are victimized by parasites or pathogens.

## 37.4 Mutualism benefits both partners

- Reef-building corals and photosynthetic dinoflagellates provide a good example of how mutualists benefit from their relationship.
- Photosynthetic dinoflagellates
  - gain a secure shelter that provides access to light,
  - produce sugars by photosynthesis that provide at least half of the energy used by the coral animals
  - use the coral's waste products, including carbon dioxide ( $\text{CO}_2$ ) and ammonia ( $\text{NH}_3$ )
  - Coral reef bleaching
  - <http://www.bbc.com/zhongwen/trad/science-39550781>



## 37.5 EVOLUTION CONNECTION: Predation leads to diverse adaptations in prey species

- Predation benefits the predator but kills the prey.
- Prey adapt using protective strategies that include
  - camouflage (<https://www.youtube.com/watch?v=GFUiCsUSzyw>),
  - mechanical defenses, and
  - chemical defenses.






## 37.6 EVOLUTION CONNECTION: Herbivory leads to diverse adaptations in plants


- Plant defenses against herbivores include
  - spines and thorns and
  - chemical toxins, often the substances that we use medicinally or for other purposes.
- “Secondary metabolites” (次級代謝物)
- Insect or pathogen resistance
- But human select against them
- A trade-off between crop quality vs. resistance

# Solanaceae (茄科)

- Many plants are toxic, with alkaloids (生物鹼)
  - Solanine (龍葵鹼 in tomato & potato)
  - Nicotine (尼古丁 in tobacco), etc.
  - Many others...
- Herbivores and plants undergo **coevolution**, a series of reciprocal evolutionary adaptations in two species in which change in one species acts as a new selective force on another species, and the resulting adaptations of the second species in turn affect the selection of individuals in the first species.

Figure 37.6





About 3,290,000 results (0.39 seconds)

[問卦] 房間突然有杏仁味，我該注意什麼？ - Gossiping板- Disp BBS

<https://disp.cc/b/163-9ai1> ▼ Translate this page

Jan 3, 2016 - 100+ posts - 100+ authors

大家好，打給後，太囑後，馬陸馬陸馬薩魯我現在在外面租房子樓上房東在施工剛剛忽然聞到杏仁味，就是杏仁豆腐杏仁茶那種味道本來以為是外面傳進來的但是我像狗一樣四處聞後發現只有我的房間有味道有沒。

標題[問卦] 氰化物真的有杏仁味嘛- terievv板- Disp BBS 26 posts 17 Aug 2017

[閒聊] 大家敢喝杏仁茶嗎- WomenTalk板- Disp BBS 100+ posts 1 Jul 2017

[問卦] 市面上杏仁茶你聞過的味道 ... 30 posts 8 Jun 2013

More results from disp.cc

氰化物- 维基百科，自由的百科全书

<https://zh.wikipedia.org/zh-tw/氰化物> ▼ Translate this page

氰化物中毒一般都很迅速。临床上常用的抢救方法是用硫代硫酸钠溶液进行静脉注射，同时使那些尚有意识的病人吸入亚硝酸异戊酯进行血管扩张来克服缺氧。常见的氰化物中毒原因是误食含氰果仁，比如生桃仁等。中毒後會發出一種獨特的苦杏仁味。一些政府把氰化物列作管制性危險品，未經申請許可者不得擅自製造、運輸、儲存及 ...

外观与气味 · 存在与应用 · 毒性 · 人體代謝過程

【化學】柯南相關--毒物(蘇老師3) - 痞客邦PIXNET

[linus871217.pixnet.net/151818498](https://linus871217.pixnet.net/151818498) - 【化學】柯南相關--毒物%2... ▼ Translate this page

Jul 9, 2013 - 氰化物在中毒之後會有杏仁味，這個是正確的，但有趣的是，只有六成的人聞的到這個杏仁味剩下的人沒有可以聞到的基因，相信柯南是那六成裡面的人 柯南動畫190~191集，小蘭的學校辦話劇表演，卻有觀眾的可樂被放了氰酸鉀 191集中，新一穿著黑衣騎士的衣服出場，解開了案情原來兇手將氰酸鉀(其實應該是氰 ...

- Cyanide
  - Bitter almonds
  - Peach
  - Apple
- Story: *Heliconius* and passionflower
- <https://www.youtube.com/watch?v=1ZRMpqw7Ak0>
- Related topic: Mimicry

# Mimicry



**EVOLUTION 3e, Figure 12.20**

© 2013 Sinauer Associates, Inc.

Up: *Heliconius melpomene*  
Down: *Heliconius erato*

Figure 37.7

Parasites



## 37.7 Parasites and pathogens can affect community composition

- Non-native pathogens can have rapid and dramatic impacts.
- **Why?**
  - The American chestnut was devastated by the chestnut blight protist.
  - A fungus-like pathogen is currently causing sudden oak death on the West Coast.




## 37.8 Trophic structure is a key factor in community dynamics

- Every community has a **trophic structure**, a pattern of feeding relationships consisting of several different levels.
  - The sequence of food transfer up the trophic levels is known as a **food chain**.
  - This transfer of food moves chemical nutrients and energy from producers up through the trophic levels in a community.




Figure 37.8



Don't forget:  
decomposers

Figure 37.9-4

Food web









## 37.10 Species diversity includes relative abundance and species richness

- **Species diversity** is defined by two components:
  1. species richness, the number of species in a community, and
  2. relative abundance, the proportional representation of a species in a community.

Table 37.10

TABLE 37.10 | RELATIVE ABUNDANCE OF TREE SPECIES IN WOODLOTS A AND B

Species	Relative Abundance in Woodlot A (%)	Relative Abundance in Woodlot B (%)
	80	25
	10	25
	5	25
	5	25



## 37.11 SCIENTIFIC THINKING: Some species have a disproportionate impact on diversity


- Ecologist Robert Paine hypothesized that the species diversity of a community is directly related to the ability of predators to prevent any one species from monopolizing local resources.
- To test this hypothesis, Paine
  - manually removed a predatory sea star known as *Pisaster* from certain areas of the intertidal zone,
  - left comparable areas intact as controls, and
  - determined the species richness of these experimental and control areas



Figure 37.11a



Figure 37.11b



Data from R.T. Paine, Food web complexity and species diversity, *American Naturalist* 100:65–75 (1996).

## 37.11 SCIENTIFIC THINKING: Some species have a disproportionate impact on diversity

- What accounted for the dramatic change?
  - A mussel of the genus *Mytilus* proved to be a superior competitor for the available space, eliminating most other invertebrates and algae.
  - In the control areas, population growth of *Mytilus* was suppressed by *Pisaster*.
  - Thus, interspecific interactions can be an important factor in the species diversity of a community.




Figure 37.11c



## 37.11 SCIENTIFIC THINKING: Some species have a disproportionate impact on diversity

- Paine's experiment and others like it gave rise to the concept of a **keystone species**, defined as a species
  - whose impact on its community is larger than its biomass or abundance indicates and
  - that occupies a niche that holds the rest of its community in place.



# How wolves change rivers

- <https://www.youtube.com/watch?v=ysa5OBhXz-Q>
- <https://dq.yam.com/post.php?id=6795>
- <https://www.nature.com/news/rethinking-predators-legend-of-the-wolf-1.14841>
- <https://www.nytimes.com/2014/03/10/opinion/is-the-wolf-a-real-american-hero.html>

## 37.12 Disturbance is a prominent feature of most communities

- **Disturbances**

- are events that damage biological communities and
  - include storms, fires, floods, drought, and human activity.
- The disturbed area may be colonized by a variety of species, which are gradually replaced by a succession of other species, in a process called **ecological succession**.

## 37.12 Disturbance is a prominent feature of most communities


- **Primary succession** begins in a virtually lifeless area with no soil.






## 37.12 Disturbance is a prominent feature of most communities

- **Secondary succession** occurs when a disturbance destroys an existing community but leaves the soil intact.



## 37.13 CONNECTION: Invasive species can devastate communities

- **Invasive species**
  - are organisms that have been introduced into non-native habitats by human actions and
- The absence of natural enemies often allows rapid population growth of invasive species.



# ECOSYSTEM STRUCTURE AND DYNAMICS




## 37.14 Ecosystem ecology emphasizes energy flow and chemical cycling

- An **ecosystem** consists of
  - all the organisms in a community and
  - the abiotic environment with which the organisms interact.
- In an ecosystem,
  - **energy flow** moves *through* the components of an ecosystem and
  - **chemical cycling** is the transfer of materials *within* the ecosystem.

Figure 37.15

Global warming, coral reef bleaching



Data from R.H. Whittaker, *Communities and Ecosystems*, second edition, New York: Macmillan (1975).

Figure 37.16a




Figure 37.16b

# The 10% rule




Figure 37.19

# Biogeochemical cycles







Figure 37.21


# Biogeochemical cycles




## 37.22 CONNECTION: A rapid inflow of nutrients degrades aquatic ecosystems

- In aquatic ecosystems, primary production is limited by low nutrient levels of
  - phosphorus (P)
  - Nitrogen (N)
- Too much P and N ? Eutrophication
- Eutrophication of lakes, rivers, and coastal waters
  - depletes oxygen levels and
  - decreases species diversity.

Figure 37.22b



**Winter**



**Summer**

Nitrogen runoff from  
Midwestern farm fields

Color: Concentrations of  
phytoplankton