

Statistics for Biology and Health

Chapter 4 Estimation of Basic Quantities for Other Sampling Schemes

Qi Guo

July 20,2019



THE UNIVERSITY OF TEXAS AT DALLAS
School of Natural Sciences and Mathematics

1. Introduction
2. Estimation of the Survival Function for Left, Double, and Interval Censoring
3. Estimation of the Survival Function for Right-Truncated Data
4. Estimation of Survival in the Cohort Life Table

Introduction

- Last Chapter we focus on estimate of the right-censored and left-truncated data, which is most common.
- In this chapter, we examine about left censoring, censored individuals provide information indicating only that the event has occurred prior to entry into the study.
- Right-truncated data, which samples arise when one samples individuals from event records and, retrospectively determines the time to event.
- Estimation techniques for grouped data.

Estimation of the Survival Function for Left, Double, and Interval Censoring

Estimation of Left Censoring Data

- Examples of pure left censoring are rare. More common are samples which include both left and right censoring.
- And in this case we use a modified KM estimator suggested by Turnbull, is based on an iterative procedure which extends the notion of a self-consistent estimator.
- To construct the estimator we assume that there is a grid of time points $0 = t_0 < t_1 < t_2 < \dots < t_m$ at which subjects are observed.
- Let d_i be the number of deaths at time t_i , and c_i, r_i are the number of individuals left-censored and right-censored at time t_i .

Algorithm

- The only information the left-censored observations at t_i give us is that the event of interest has occurred at some $t_j \leq t_i$
- The self-consistent estimator estimates the probability that this event occurred at each possible t_j less than t_i based on an initial estimate of the survival function.
- Using this estimate, we compute an expected number of deaths at t_j , which is then used to update the estimate of the survival function and the procedure is repeated until the estimated survival function stabilizes.

Algorithm

- Step 0: Produce an initial estimate of the survival function at each t_j , $S_0(t_j)$. Note any legitimate estimate will work. Turnbull's suggestion is to use the KM estimate obtained by ignoring the left-censored observations.
- Step $(K + 1)1$: Using the current estimate of S , estimate $p_{ij} = p[t_{j-1} < x \leq t_j | x \leq t_j]$ by $\frac{S_K(t_{j-1}) - S_K(t_j)}{1 - S_K(t_j)}$ for $j \leq i$.
- Step $(K + 1)2$: Using the results of the previous step, estimate the number of events at time t_i by $\hat{d}_i = d_i + \sum_{j=i}^m c_j p_{ij}$.
- Step $(K + 1)3$: Compute the KM estimator based on the estimated right-censored data with \hat{d}_i events and r_i right-censored observations at t_i , ignoring the left-censored data. If this estimate, $S_{K+1}(t)$, is close to $S_K(t)$ for all t_i , stop the procedure; if not, go to step 1.

Estimation of the interval-censored data

- Sometimes data may be interval-censored. Here the only information we have for each individual is that their event time falls in an interval $(L_i, R_i], i = 1, \dots, n$, but the exact time is unknown.
- And we still have a iteration like the estimation of left-censored data in page 6.

Estimation of the Survival Function for Right-Truncated Data

Right-Truncated Data

- For right-truncated data, only individuals for which the event has occurred by a given date are included in the study. Right truncation arises commonly in the study of infectious diseases.
- Let T_i denote the chronological time at which the i th individual is infected and X_i the time between infection and the onset of disease.
- Sampling consists of observing (T_i, X_i) for patients over the period $(0 \text{ to } \tau)$. Note that only patients who have the disease prior to τ are included in the study.
- Estimation for this type of data proceeds by reversing the time axis. Let $R_i = \tau - X_i$. The R_i 's are now left-truncated in that only individuals with value of $T_i \leq R_i$ are included in the sample.

Estimation of Survival in the Cohort Life Table

Estimation of Survival in the Cohort Life Table

- A “cohort” is a group of individuals who have some common origin from which the event time will be calculated
- They are followed over time and their event time or censoring time is recorded to fall in one of $k + 1$ adjacent, nonoverlapping intervals, $(a_{j-1}, a_j]$, $j = 1, \dots, k + 1$.
- A traditional cohort life table presents the actual mortality experience of the cohort from the birth of each individual to the death of the last surviving member of the cohort.
- The basic construction of the cohort life table is a little complicated, and more details in columns in P152.