



DECART Summer School 2018:

Causal Inference Module

Matching


# Why match?

- We need to adjust for covariate  $X_i$  in observational study, even when the assumption of no unmeasured confounding holds.
- Easy solution: Use a parametric model for  $E(Y_i(A)|X_i)$ .
- But misspecified parametric model will lead to wrong causal estimates.
- Matching is a natural alternative solution with two benefits:
  - Reduces the dependence of estimates on parametric models.
  - Can simplify the analysis of causal effects.

# Why parametric model failed?

- Often lead to **large variation** in the estimates of interest.
- Why does this occur?

- Parametric model will extrapolate to regions with only treated or only control.
- Modeling assumption will affect these extrapolation.



# What does matching do?

- Allows for relatively nonparametric ways of estimating the casual effect.
- **Caution:** Matching doesn't justify a causal effect automatically.
- **Important Note:** Without strong design (such as clinical trial), no statistical modeling could completely make the move from correlation to causation persuasive.

# Assumptions

1. Consistency:

$$Y_i = Y_i(A_i)$$

2. Unmeasured Confounders Assumption:

$$A_i \perp (Y_i(0), Y_i(1)) | X_i$$

3. Positivity

$$0 < P(A_i = 1 | X_i) < 1$$

# (Simple) Exact Matching Without Replacement

- Let  $X_i$  take on a finite number of values,  $x$ .
- Let  $I_t = \{1, 2, \dots, N_t\}$  be the set of treated units.
- Exact matching without replacement:

For each treated unit,  $i \in I_t$

1. Find the set  $M_i^c$  of unmatched control units  $j$  such that  $X_i = X_j$ ,  
for  $j \in M_i^c$

2. Randomly select one of these control units to be the match,  
indicated  $j(i)$ .

- Let  $I_c = \{j(1), j(2), \dots, j(N_t)\}$  be the set of control units.
- The distribution of  $X_i$  will be exactly the same for the treated and matched control:

$$P(X_i = x | A_i = 1) = P(X_i = x | A_i = 0, I_c)$$

If the data is exactly matched, then an unbiased estimator for the average treatment effect for the treated (ATT) is:

$$\hat{\tau}_t^{match} = \frac{1}{N_t} \sum_{i:A_i=1} \hat{\tau}_i^{match} = \frac{1}{N_t} \sum_{i:A_i=1} (Y_i^{obs} - Y_{m_i^c}^{obs})$$

i.e.

$$E[\hat{\tau}_t^{match}] = \tau_{ATT} = E[Y_i(1)|A_i = 1] - E[Y_i(0)|A_i = 1]$$

Proof:

$$\begin{aligned} E[\hat{t}_t^{match}] &= E\left[\frac{1}{N_t} \sum_{i:A_i=1} (Y_i^{obs} - Y_{m_i^c}^{obs})\right] \\ &= E\left[\frac{1}{N_t} \sum_{i:A_i=1} (Y_i^{obs})\right] - E\left[\frac{1}{N_t} \sum_{i:A_i=1} E(Y_{m_i^c}^{obs})\right] \\ &= \int E[Y_i|A_i = 1, X_i = x]dP(X_i = x|A_i = 1) \\ &\quad - \int E[Y_i|A_i = 0, X_i = x]dP(X_i = x|A_i = 1) \\ &= \int E[Y_i(1)|A_i = 1, X_i = x]dP(X_i = x|A_i = 1) \\ &\quad - \int E[Y_i(0)|A_i = 1, X_i = x]dP(X_i = x|A_i = 1) \\ &= E[Y_i(1)|A_i = 1] - E[Y_i(0)|A_i = 1] \end{aligned}$$



# Weakening the identification assumptions

- Consistency, no unmeasured confounders, total expectation and exact matching property  
⇒ identifying the ATT.
- Can weaken no unmeasured confounders to conditional mean independence (CMI):
$$E[Y_i(0)|X_i = x, A_i = 1] = E[Y_i(0)|X_i = x, A_i = 0]$$
- Nice features of CMI:
  1. Only make assumptions about  $Y_i(0)$  not  $Y_i(1)$ .
  2. Only make assumptions on the means, not other aspects of distribution (variance, skewness, kurtosis, etc).

# Analyzing exactly matched data

- Simple difference in observed means:

$$\hat{t}_{ATT} = \frac{1}{N_T} \sum_{i \in I_t} Y_i - \frac{1}{N_c} \sum_{j \in I_c} Y_j$$

- In simple matching mentioned above (exact, 1-to-1, no replacement):

$$\hat{t}_{ATT} = \frac{1}{N_T} \sum_{i=1 \dots N_T} (Y_i - Y_{j(i)})$$

$$\widehat{var}(\hat{t}_{ATT}) = \frac{1}{N_T} \sum_{i=1 \dots N_T} (Y_i - Y_{j(i)} - \hat{t}_{ATT})^2$$

- In practice, such an exact matching scheme is rarely feasible
  - exact matching is typically impossible
  - the pool of potential matches is often too small to ignore the conflicts

# Inexact Matching without Replacement

- Match the  $i$ th treated unit with covariate values  $X_i$  to control unit  $m_i$ , that is, the control unit that solves
$$m_i^c = \operatorname{argmin}_{i' \in I_c} \|X_i - X_{i'}\|$$
- one control unit might be identified as match for more than one unit

> match all units simultaneously

$$\operatorname{argmin}_{m_1^c, \dots, m_{N_t}^c \in I_c} \sum_{i=1}^{N_t} \|X_i - X_{m_i^c}\| \text{ subject to } m_i \neq m_{i'}$$

> match units sequentially (“greedy” matching algorithm):  
the ordering matters

One option: match those difficult ones first: the rank of the estimated propensity scores (high  $\rightarrow$  low)

# Inexact Matching without Replacement

- When multiple matches (equally close) to one treated unit
  - > use the average of the outcomes for this set of tied matches as the control potential outcome for treated unit  $i$ ,  $\sum_{i' \in M_i^c} Y_{i'}(0) / M_i$ , with  $M_i$  be the cardinality of  $M_i^c$ 
    - reduced sampling variance of the resulting estimator
    - removing more units from the pool of possible control units available for subsequent matches
    - > some selection mechanism, e.g. random selection

# Distance metrics

- We need a distance metric to define distance/similarity on  $X_i$  and  $X_j$ , which might be high dimensional.
  - Lower value  $\rightarrow$  more similar values
  - Choice of distance metric will lead to different matches
- Possible choices of distance:
  - > Propensity score distance metric
  - > Euclidean distance metric
  - > Mahalanobis distance metric
  - > Caliper metric
  - > Hybrid metric

# Propensity score distance

- Propensity score:  $e(X_i) = P(A_i = 1|X_i)$
- Rubin et al have shown that propensity score matching has good properties **if covariates are roughly normal**.
- Propensity score distance:

Option 1:  $D_{ij} = |e(X_i) - e(X_j)|$

Option 2:  $D_{ij} = |\text{logit}(e(X_i)) - \text{logit}(e(X_j))|$

# Euclidean distance

- Suppose that  $X_i = (X_{i1}, \dots, X_{iK})$ .
- The **Euclidean distance** metric is

$$D_{ij} = \sqrt{\sum_{k=1}^K \frac{(X_{ik} - X_{jk})^2}{\hat{\sigma}_k^2}},$$

$$\text{where } \hat{\sigma}_k^2 = \frac{1}{N-1} \sum_{i=1}^N (X_{ik} - \bar{X}_k)^2.$$



# Mahalanobis distance

- Intuition: if  $X_{ik}$  and  $X_{ik}'$  are highly correlated, then their contribution to the distances should be lower.
  - Easy to get close on correlated covariates, then downweight it
  - Harder to get close on uncorrelated covariates, then upweight it
- The Mahalanobis distance is

$$D_{ij} = \sqrt{(X_i - X_j)^T \hat{\Sigma}^{-1} (X_i - X_j)},$$

where weight matrix  $\hat{\Sigma} = \frac{1}{N} \sum_{i=1}^N (X_i - \bar{X})(X_i - \bar{X})^T$ .

# Caliper

- To overcome shortcomings of erroneously choosing control, we will only select the control if its distance to case meet condition:

$$D_{ij} < \varepsilon$$

- Rubin (1985) suggested using a caliper size of a quarter of a standard deviation of the sample estimate propensity score ( $\varepsilon = \sigma_{ps}/4$ ).

# Hybrid metrics

Example:

Exact on race/gender, Mahalanobis on the rest.

# The general matching procedure

1. Choose a number of matches
2. Choose a distance metric
3. Find matches (drop non-matches)
4. Check balance
5. Repeat 1-4 until balance is acceptable
6. Calculate the effect of the treatment on the outcome in the matched dataset.

# The Card-Krueger Minimum Wage Data

- Card and Krueger (1995) were interested in evaluating the effect of raising the state minimum wage in New Jersey in 1993.
- data were collected on employment at fast-food restaurants in New Jersey and in the neighboring state of Pennsylvania.
- Covariates are measured on restaurant level prior to the raise;
- The outcome is employment after the raise (final empl).

## The Card-Krueger New Jersey and Pennsylvania Minimum Wage Data

	(N = 347)		(N <sub>t</sub> = 279) (treated)		(N <sub>c</sub> = 68) (controls)		Nor	Log Ratio
	Mean	(S.D.)	Mean	(S.D.)	Mean	(S.D.)	Dif	of STD
initial empl	17.84	(9.62)	20.17	(11.96)	17.27	(8.89)	−0.28	−0.30
burger king	0.42	(0.49)	0.43	(0.50)	0.42	(0.49)	−0.02	−0.01
kfc	0.19	(0.40)	0.13	(0.34)	0.21	(0.41)	0.20	0.17
roys	0.25	(0.43)	0.25	(0.44)	0.25	(0.43)	0.00	−0.00
wendys	0.14	(0.35)	0.19	(0.40)	0.13	(0.33)	−0.18	−0.18
initial wage	4.61	(0.34)	4.62	(0.35)	4.60	(0.34)	−0.05	−0.02
time until raise	17.96	(11.01)	19.05	(13.46)	17.69	(10.34)	−0.11	−0.26
pscore	0.80	(0.05)	0.79	(0.06)	0.81	(0.04)	0.28	−0.35
final empl	17.37	(8.39)	17.54	(7.73)	17.32	(8.55)		

# Excise on the Card-Krueger Data

- For this illustration, we focus on a small subset of 20 restaurants
- 5 from New Jersey and 15 from Pennsylvania
- We use only initial employment (initial empl) and restaurant chain (burger king or kfc) as pre-treatment variables
- Inexact match without replacement

## 20 Units from the Card-Krueger Dataset

Unit	State	chain	initial empl	final empl
$i$	$W_i$	$X_{i1}$	$X_{i2}$	$Y_i^{\text{obs}}$
1	NJ	BK	22.5	40.0
2	NJ	KFC	14.0	12.5
3	NJ	BK	37.5	20.0
4	NJ	KFC	9.0	3.5
5	NJ	KFC	8.0	5.5
6	PA	BK	10.5	15.0
7	PA	KFC	13.8	17.0
8	PA	KFC	8.5	10.5
9	PA	BK	25.5	18.5
10	PA	BK	17.0	12.5
11	PA	BK	20.0	19.5
12	PA	BK	13.5	21.0
13	PA	BK	19.0	11.0
14	PA	BK	12.0	17.0
15	PA	BK	32.5	22.5
16	PA	BK	16.0	20.0
17	PA	KFC	11.0	14.0
18	PA	KFC	4.5	6.5
19	PA	BK	12.5	31.5
20	PA	BK	8.0	8.0

Match Order = 1,2,3,4,5; Metric = $x_1^2 + x_2^2$				
$i$	$m_i^c$	$Y_i^{\text{obs}}$	$Y_{m_i^c}^{\text{obs}}$	$\hat{\tau}_i^{\text{match}}$
1	11	40.0	19.5	20.5
2	7	12.5	17	-4.5
3	15	20.0	22.5	-2.5
4	8	3.5	10.5	-7
5	20	5.5	8.0	-2.5

$\hat{\tau}_t^{\text{match}}$  +0.8

Match Order = 1,2,3,5,4; Metric = $x_1^2 + x_2^2$				
$i$	$m_i^c$	$Y_i^{\text{obs}}$	$Y_{m_i^c}^{\text{obs}}$	$\hat{\tau}_i^{\text{match}}$
1	11	40.0	19.5	20.5
2	7	12.5	17.0	-4.5
3	15	20.0	22.5	-2.5
5	8	5.5	10.5	-5
4	20	3.5	8.0	-4.5

$\hat{\tau}_t^{\text{match}}$  +0.8

Match Order = 1,2,3,4,5; Metric = $100 \cdot x_1^2 + x_2^2$				
$i$	$m_i^c$	$Y_i^{\text{obs}}$	$Y_{m_i^c}^{\text{obs}}$	$\hat{\tau}_i^{\text{match}}$
1	11	40.0	19.5	20.5
2	7	12.5	17.0	-4.5
3	15	20.0	22.5	-2.5
4	8	3.5	10.5	-7
5	17	5.5	14.0	-8.5

$\hat{\tau}_t^{\text{match}}$  -0.4



# The Bias of Matching Estimator

- The potential bias created by discrepancies between the pre-treatment covariates of the units within a matched pair.

$$\begin{aligned} E \left[ \hat{\tau}_i^{match} | A_i = 1, X_i, X_{m_i^c} \right] &= E \left[ Y_i^1 - Y_{m_i^c}^0 | X_i, X_{m_i^c} \right] \\ &= \mu_t(X_i) - \mu_c(X_{m_i^c}) = \tau(X_i) + \mu_c(X_i) - \mu_c(X_{m_i^c}) \end{aligned}$$

- The unit-level bias is  $B_i = \mu_c(X_i) - \mu_c(X_{m_i^c})$
- Bias adjustment:

$$\hat{\tau}_t^{adj} = \frac{1}{N_t} \sum_{i:A_i=1} (Y_i - Y_{m_i^c} - \hat{B}_i)$$

$\hat{B}_i$  can be estimated through linear model

# Bias Correction Using Linear Model

- If we assume linear models for the group specific means  $\mu_c(x) = \alpha_d + x\beta_d$  and  $\mu_t(x) = \tau + \alpha_d + x\beta_d$
- Then we can estimate the bias as  $\hat{B}_i = \hat{\mu}_c(X_i) - \hat{\mu}_c(X_{m_i^c}) = (X_i - X_{m_i^c})\hat{\beta}_d$
- Three simple regression based approaches can be considered to obtain  $\hat{B}_i$ :
  1. **Regression on the Matching Discrepancy** ( $D_i = X_i - X_{m_i^c}$ )  
 $Y_i^{obs} - Y_{m_i^c}^{obs} = \tau + D_i\beta_d + v_i$ :  $Y_i^{obs} - Y_{m_i^c}^{obs} \sim D_i \Rightarrow \hat{\beta}_d$
  2. **Control Regression on Covariates**  
 $Y_{m_i^c} = \alpha_c + X_{m_i^c}\beta_c + v_{ci}$ :  $Y_{m_i^c} \sim X_{m_i^c} \Rightarrow \hat{\beta}_c$
  3. **Pooled Regression on Covariates**  
 $Y_i = \alpha_p + \tau_p A_i + X_i\beta_p + v_i$ :  $Y_i \sim A_i + X_i \Rightarrow \hat{\beta}_p$

# Data Illustration of Bias Correction

**Matching Discrepancy for the 20 Units from the Card-Krueger Data (Match Order 1,2,3,4,5; Metric  $x_1^2 + x_2^2$ )**

$i$	$m_i$	$Y_i^{\text{obs}}$	$Y_{m_i^c}^{\text{obs}}$	$\hat{\tau}_i^{\text{match}}$	$X_{i,1}$	$X_{i,2}$	$X_{m_i^c,1}$	$X_{m_i^c,2}$	$D_{i,1}$	$D_{i,2}$
1	11	40.0	19.5	20.5	0	22.5	0	20.0	0	2.5
2	7	12.5	17.0	-4.5	1	14.0	1	13.8	0	0.2
3	15	20.0	22.5	-2.5	0	37.5	0	32.5	0	5.0
4	8	3.5	10.5	-7.0	1	9.0	1	8.5	0	0.5
5	20	5.5	8.0	-2.5	1	8.0	0	8.0	1	0

**Bias-Adjustment Regression Coefficients for the 20 Units from the Card-Krueger Data**

	Difference Regression (Approach #1)	Control Regression (Approach #2)	Pooled Regression (Approach #3)
Regression coefficients			
Intercept	-1.30	4.21	12.01
Treatment indicator	—	—	1.63
Restaurant chain	-1.20	2.65	-7.32
Initial employment	1.43	0.62	0.39

# Data Illustration of Bias Correction

## Regression on the Matching Discrepancy (Difference Regression):

First pair  $(i, m_i) = (1, 11)$ ,  $X_1 = (0, 22.5)$ ,  $X_{m_1} = (0, 20.0)$

Thus the adjusted control outcome:

$$\begin{aligned}\hat{Y}_1(0) &= Y_{m_1} + D_1 \hat{\beta}_d = 19.5 - 1.20 \times D_{1,1} + 1.43 \times D_{1,2} \\ &= 19.5 - 1.20 \times 0 + 1.43 \times 2.5 = 23.1\end{aligned}$$

The adjusted estimate of the unit-level treatment effect

$$\hat{\tau}_1^{adj} = Y_1(1) - \hat{Y}_1(0) = 40.0 - 23.1 = 16.9$$

Similarly, we can obtain the following full set of results:

$i$	$m_i$	$Y_i(1)$	$Y_{m_i^c}(0)$	$X_{i,1}$	$X_{i,2}$	$X_{m_i^c,1}$	$X_{m_i^c,2}$	$D_{i,1}$	$D_{i,2}$	$\hat{\beta}_d^T D_i$	$\hat{Y}_i(0)$
1	11	40.0	19.5	0	22.5	0	20.0	0	2.5	3.6	23.1
2	7	12.5	17.0	1	14.0	1	13.8	0	0.2	0.3	17.3
3	15	20.0	22.5	0	37.5	0	32.5	0	5.0	7.1	29.6
4	8	3.5	10.5	1	9.0	1	8.5	0	0.5	0.7	11.2
5	20	5.5	8.0	1	8.0	0	8.0	1	0	-1.2	6.8
$\hat{\tau}_t^{match} = +0.8$				$\hat{\tau}_t^{adj} = -1.3$							

# Data Illustration of Bias Correction

## Control Regression on Covariates:

First pair  $(i, m_i) = (1, 11)$ ,  $X_1 = (0, 22.5)$ ,  $X_{m_1} = (0, 20.0)$

Thus the adjusted control outcome:

$$\begin{aligned}\hat{Y}_1(0) &= Y_{m_1} + D_1 \hat{\beta}_c = 19.5 + 2.65 \times D_{1,1} + 0.62 \times D_{1,2} \\ &= 19.5 - 2.65 \times 0 + 0.62 \times 2.5 = 21.1\end{aligned}$$

The adjusted estimate of the unit-level treatment effect

$$\hat{\tau}_1^{adj} = Y_1(1) - \hat{Y}_1(0) = 40.0 - 21.1 = 18.9$$

Similarly, we can obtain the following full set of results:

$i$	$m_i$	$Y_i(1)$	$Y_{m_i^c}(0)$	$X_{i,1}$	$X_{i,2}$	$X_{m_i^c,1}$	$X_{m_i^c,2}$	$D_{i1}$	$D_{i2}^*$	$\hat{\beta}_c^T D_i$	$\hat{Y}_i(0)$
1	11	40.0	19.5	0	22.5	0	20.0	0	2.5	1.5	21.0
2	7	12.5	17.0	1	14.1	1	13.8	0	0.2	0.1	17.1
3	15	20.0	22.5	0	37.5	0	32.5	0	5.0	3.1	25.6
4	8	3.5	10.5	1	9.0	1	8.5	0	0.5	0.3	10.8
5	20	5.5	8.0	1	8.0	0	8.0	1	0	2.7	10.7
				$\hat{\tau}_t^{match} = +0.8$		$\hat{\tau}_t^{adj} = -0.7$					

# Data Illustration of Bias Correction

## Pooled Regression on Covariates:

First pair  $(i, m_i) = (1, 11)$ ,  $X_1 = (0, 22.5)$ ,  $X_{m_1} = (0, 20.0)$

Thus the adjustment for the 1<sup>st</sup> pair:

$$\begin{aligned}\hat{B}_1 &= -7.32 \times D_{1,1} + 0.39 \times D_{1,2} \\ &= -7.32 \times 0 + 0.39 \times 2.5 = 0.98\end{aligned}$$

The adjusted estimate of the unit-level treatment effect

$$\hat{\tau}_1^{adj} = Y_1(1) - Y_{m_1}(0) - \hat{B}_1 = 40.0 - 19.5 - 0.98 = 19.52$$

Similarly, we can obtain the following full set of results:

$i$	$m_i$	$Y_i(1)$	$Y_{m_i^c}(0)$	$X_{i,1}$	$X_{i,2}$	$X_{m_i^c,1}$	$X_{m_i^c,2}$	$D_{i1}$	$D_{i2}$	$\hat{\beta}_s^T D_i$	$\hat{Y}_i(0)$
1	11	40.0	19.5	0	22.5	0	20.0	0	2.5	1.0	20.5
2	7	12.5	17.0	1	14.0	1	13.8	0	0.2	0.1	17.1
3	15	20.0	22.5	0	37.5	0	32.5	0	5.0	1.9	24.4
4	8	3.5	10.5	1	9.0	1	8.5	0	0.5	0.2	10.7
5	20	5.5	8.0	1	8.0	0	8.0	1	0	-7.3	0.7
$\hat{\tau}_t^{match} = +0.8$				$\hat{\tau}_t^{adj} = +1.6$							

# Matching with Replacement

- The set of controls selected does not depend on the ordering of treated units.
- Let  $L(i)$  be the number of times each control unit id used as a match

$$L(i) = \sum_{j=i}^{N_t} 1_{j \in M_i^c}$$

When matching without replacement,  $L(i) \in \{0,1\}$  for all units.

$$\hat{\tau}_t^{repl} = \frac{1}{N_t} \sum_{i=1}^N (A_i \cdot Y_i^{obs} - (1 - A_i) \cdot L(i) \cdot Y_i^{obs})$$

# With or without replacement

- Matching with replacement: a single control unit could be matched repeatedly with multiple treated units.
- Pro:
  1. Better matches!
  2. Order of matching does not matter.
- Con:
  1. need more complicated inference.
  2. need to account for multiple appearances with weights.
  3. potentially higher uncertainty (using the same data multiple times=relying on less data)



# The Number of Matches

- Let  $\sigma_c^2$  and  $\sigma_t^2$  be the super-population variances of  $Y_i^0$  and  $Y_i^1$  conditional on the covariates.

- If we use M matches, the estimator then

$$\hat{\tau}_t^{match,M} = \frac{1}{N_t} \sum_{i=1}^{N_t} \left( Y_i^1 - \frac{1}{M} \sum_{j \in M_i^c} Y_j^0 \right)$$

- The sample variance is then

$$Var(\hat{\tau}_t^{match,M}) = \frac{1}{N_t} \left( \sigma_t^2 + \frac{\sigma_c^2}{M} \right)$$

- Assume equal variance  $\sigma_c^2 = \sigma_t^2$

$$\frac{Var(\hat{\tau}_t^{match,1}) - Var(\hat{\tau}_t^{match,M})}{Var(\hat{\tau}_t^{match,1})} = \frac{M-1}{2M}$$

M=2 reduces the sample variance by 25% relative to using a single match

# Assessing balance

- All matching methods seek to find the balance:

$$P(X_i = x|A_i = 1, \mathcal{S}) = P(X_i = x|A_i = 0, \mathcal{S})$$

- Choice of balance metric will determine which matching method does better.
- Options: estimation of matching performance
  1. Differences in mean/medians, standardized.
  2. QQ plot/K-S statistics for comparing the entire distribution.
- ...

# Estimand

- Matching easiest to justify for the average treatment effect for the treated (ATT).

$$\hat{\tau}_t = \frac{1}{N_t} \sum_{i:A_i=1} (Y_i^{obs} - Y_{m_i^c}^{obs})$$

- Can also justify the average treatment effect for the controls (ATU) by finding matched treated units for the controls.

$$\hat{\tau}_c = \frac{1}{N_c} \sum_{i:A_i=0} (Y_{m_i^t}^{obs} - Y_i^{obs})$$

- Combined the two to obtain the average treatment effect for the entire sample (ATE):

$$\hat{\tau} = \frac{N_c}{N_c + N_t} \hat{\tau}_c + \frac{N_t}{N_c + N_t} \hat{\tau}_t$$