
# Bond Angles Read-ahead

### Introduction

The chemical properties of a given molecule depend in a large part on the geometry of the molecule. One of the simplest molecules is methane, CH<sub>4</sub>, which is the main component of natural gas. Each of the four hydrogen atoms is bonded to a single, central, carbon atom, forming a tetrahedron, as seen in Figure 1 below. In this set of problems, we determine the bond angle between any pair of hydrogen atoms.








Figure 2: A natural source of methane.

### Instructions


After reading through *Bond Angles* context and questions below, you should complete the reflection assignment in Canvas. Note: *you will have a chance to talk further with your coach before answering the questions below in detail.* The point of this read-ahead and the reflection is to "prime the pump" for further conversations with your coaches.

## **Bond Angles**

Draw a perpendicular line from the carbon atom (C) to the plane containing three of the hydrogen atoms. Let Q represent the foot of this perpendicular line (see figures below), and let y represent the distance between the carbon atom and any of the hydrogen atoms. Let a represent the distance from Q to one of the hydrogen atoms, and let x represent the measure of the required bond angle.

## Questions

- 1. Consider the angle HQH in Figure 4 above. Find the measure of this angle, in degrees.
- Find an expression for the distance a (the length of the line segment HQ) in terms of the distance y and the angle x.
- 3. Now, examine the triangle formed by two hydrogen atoms and point Q, as shown in Figure 5 below. The altitude from point Q divides the triangle HQH into two congruent triangles. So, the vertex angle HQH is divided into two angles whose measures are each  $60^{\circ}$ . In terms of a, what is the length of the line segment  $\overline{\text{HT}}$ ?
- 4. Figure 6 above represents the triangle formed by the carbon atom and two hydrogen atoms. From this figure, we see that  $\sin\left(\frac{x}{2}\right) = \frac{a\sqrt{3}}{2y}$ . Use the double angle formula  $\sin^2\left(\theta\right) = \frac{1-\cos\left(2\theta\right)}{2}$  to



rewrite the left-hand side of this equation in terms of  $\cos(x)$ , then combine the resulting equation with the expression for a you found in problem 2 above, and solve for  $\cos(x)$ . Note that there are two solutions, one positive, one negative.

5. Note that only one of the two values for cos(x) you found in the previous part makes sense. Find x to two decimal places.

# Instructions, part deux

After reading and reflecting on these questions, complete the pre-read assignment on Canvas. This will give your coach some insight on your thinking in order to best help you before you are required to formally answer these questions.