Shape as a Measure of Weapon Standardisation: A Replication

Corey Christopherson, Thomas Winegarden, Richard Todd, Zack Garcia

Repo

Birch, T., & Martinon-Torres, M. (2018). Shape as a measure of weapon standardisation: From metric to geometric morphometric analysis of the Iron Age 'Havor' lance from Southern Scandinavia. Journal of Archaeological Science, 101, 34-51. doi: https://doi.org/10.1016/j.jas.2018.11.002

Contents lists available at ScienceDirect



Journal of Archaeological Science





journal homepage: www.elsevier.com/locate/jas

Shape as a measure of weapon standardisation: From metric to geometric morphometric analysis of the Iron Age 'Havor' lance from Southern Scandinavia

Thomas Birch^{a,*}, Marcos Martinón-Torres^b

- Investigated two approaches to quantify the standardization of manufacturing
- Compared standard measurements to geometric morphometric (GMM) shape analysis
- The authors found that GMM allowed for an analysis of overall shape instead individual dimensions






(max)

socket (min)

(blade width)

Claims Addressed



Is this the same underlying data?


- Pairwise correlations of lance features
- Correlation coefficients recorded in R code comments

Replicate ANOVA Analysis Finding

 There was found to be an association between overall centroid size and shape that could not be explained through site difference

```
# Pairwise correlations of lance features
cor(socket$socket_length,socket$sock_bdiam,use="pairwise.complete.obs") # 0.52
cor(socket$sock_bdiam,socket$sock_sdiam,use="pairwise.complete.obs") # 0.52
cor(lances_M$width,lances_M$thickness,use="pairwise.complete.obs") # 0.53
cor(blade$blade_length,blade$width,use="pairwise.complete.obs") # 0.58
cor(blade$blade_length,blade$thickness,use="pairwise.complete.obs") # 0.47
```

Highlights/Changes 1 - pairwise correlation



Highlights/Changes 2 - spatial ANOVA

```
## Homogeneity of Slopes Test
## Allometry Model
## Call:
## procD.allometry(f1 = shape ~ size, f2 = ~site, print.progress = FALSE,
      data = lances geomorph, method = "PredLine")
##
##
##
## Homogeneity of Slopes Test
                                     SS
                                             Rsa
                                                              Z Pr(>F)
## Common Allometry 74 0.76683
## Group Allometries 72 0.74137 0.025466 0.030754 1.2366 0.73767 0.234
## The null hypothesis of parallel slopes is supported
    based on a significance criterion of alpha = 0.05
## Based on the results of this test, the following ANOVA table is most
appropriate
##
## Type I (Sequential) Sums of Squares and Cross-products
## Randomized Residual Permutation Procedure Used
## 1000 Permutations
## ANOVA effect sizes and P-values based on empirical F distributions
##
                                                      Z Pr(>F)
## log(size) 1 0.05502 0.055024 0.06645 5.3098 1.87975 0.012 *
## site
             2 0.00619 0.003096 0.00748 0.2987 -0.83913 0.812
## Residuals 74 0.76683 0.010363 0.92607
## Total
            77 0.82805
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
```

Reflections

- Had to switch papers after the first check-in
- Exposure to new academic field
- Knitting on each machine early on was DIFFICULT
- Chunks of the original R code were missing
- Solid balance of work (R wrangling, writing, managing supplemental files)