

Appendix A - Abrolhos Marine Park Western Rock Lobster Report

Yamatji Shallow Bank, 2025 and 2026 pot surveys

Table of contents

Survey effort	1
Abundance	2
Size structure	6
Standardised size distributions	10
Modelling: catch inside vs outside the National Park Zone	13
Data notes	22

Survey effort

Table 1: Summary of western rock lobster pot sampling on Yamatji Shallow Bank in the 2025 and 2026 surveys, showing the number of pots deployed, the number of pots that fished successfully and were counted, and the number of lobsters measured in each management zone. NPZ = National Park Zone (no take), SPZ = Special Purpose Zone (open to commercial and recreational fishing). Pots that were lost, had a missing trap door, or were flagged in the field as not countable are deployed but not counted.

Year	Zone	Pots deployed	Pots counted	Lobsters measured
2025	National Park Zone	55	54	273
2025	Special Purpose Zone	93	91	483
2026	National Park Zone	55	53	124
2026	Special Purpose Zone	84	83	248

Lobster pot locations, Abrolhos Marine Park




Figure 1: Locations of lobster pots on Yamatji Shallow Bank, shown separately for the 2025 and 2026 surveys. Commonwealth marine park zone boundaries and 10 m depth contours are shown. ABMP = Abrolhos Marine Park (Commonwealth).

Abundance

Table 2: Western rock lobster catch rates by zone and survey year, showing the number of pots counted, the total number of lobsters caught, and the mean and standard error of the number of lobsters per pot for all lobsters and for legal sized lobsters. Pots that caught no lobsters are included as zeros. Catch is reported per pot rather than per pot-hour because set times were not recorded in 2026.

Year	Zone	Pots	Total lobsters	Mean per pot	SE	Mean legal per pot	SE legal
2025	National Park Zone	54	261	4.83	0.77	4.09	0.57
2025	Special Purpose Zone	91	471	5.18	0.71	3.13	0.43
2026	National Park Zone	53	124	2.34	0.45	1.72	0.35
2026	Special Purpose Zone	83	248	2.99	0.58	0.94	0.19




Figure 2: Western rock lobster catch rate (mean number of lobsters per pot, +/- standard error) by management zone and survey year, from pot sampling on Yamatji Shallow Bank. Sample sizes are the number of pots counted in each zone and year. Pots that caught no lobsters are included as zeros.

Western rock lobster abundance per pot




Figure 3: Spatial distribution of total western rock lobster abundance per pot, shown separately for the 2025 and 2026 surveys. Symbol size is scaled to the number of lobsters caught and crosses mark pots that caught no lobsters. Commonwealth marine park zone boundaries are shown.




Figure 4: Spatial distribution of legal sized western rock lobster abundance per pot, shown separately for the 2025 and 2026 surveys. Symbol size is scaled to the number of legal sized lobsters caught. Legal size is a carapace length greater than 76 mm. Commonwealth marine park zone boundaries are shown.

Size structure

Table 3: Summary of western rock lobster carapace length by zone and survey year, showing the number of lobsters measured, the mean, median, minimum and maximum carapace length, and the percentage of measured lobsters above the legal minimum size. The minimum legal carapace length for western rock lobster is 76 mm.

Year	Zone	n	Mean length (mm)	Median (mm)	Min	Max	% legal
2025	National Park Zone	273	92.7	93	24.0	129.0	85.0
2025	Special Purpose Zone	483	81.6	82	50.0	129.0	60.7
2026	National Park Zone	124	86.6	90	51.0	120.0	73.4
2026	Special Purpose Zone	248	73.0	72	50.3	125.5	31.5




Figure 5: Carapace length frequency distribution of western rock lobster by management zone (columns) and survey year (rows), from pot sampling on Yamatji Shallow Bank. Bars are counts of lobsters in 5 mm length bins. The dashed red line shows the 76 mm legal minimum size.




Figure 6: Carapace length frequency distribution of western rock lobster by management zone (columns), survey year and sex (rows), from pot sampling on Yamatji Shallow Bank. Bars are counts of lobsters in 5 mm length bins, with zone setting the colour and sex the shade, and males shown as the lighter bars. Lobsters whose sex was not recorded are excluded. The dashed red line shows the 76 mm legal minimum size.

Standardised size distributions




Figure 7: Standardised carapace length frequency distribution of western rock lobster by management zone (columns) and survey year (rows), from pot sampling on Yamatji Shallow Bank. Each panel is scaled to sum to 100% so that the shape of the size distribution can be compared independently of the number of lobsters caught. Sample sizes are given in each panel, as percentages from a small sample are considerably noisier. The dashed red line shows the 76 mm legal minimum size.




Figure 8: Standardised carapace length frequency distribution of western rock lobster by management zone (columns), survey year and sex (rows), from pot sampling on Yamatji Shallow Bank. Each panel is scaled to sum to 100%, with zone setting the colour and sex the shade, and males shown as the lighter bars. Sample sizes are given in each panel. Lobsters whose sex was not recorded are excluded. The dashed red line shows the 76 mm legal minimum size.




Figure 9: Standardised carapace length frequency distribution of male western rock lobster only, by survey year (columns) and management zone (rows), from pot sampling on Yamatji Shallow Bank. Each panel is scaled to sum to 100%, with the two zones stacked within each year so that their size distributions can be compared directly. Sample sizes are given in each panel. The dashed red line shows the 76 mm legal minimum size.




Figure 10: Legal and sublegal western rock lobster catch rate (mean number of lobsters per pot, +/- standard error) by management zone (bars) and survey year (columns), from pot sampling on Yamatji Shallow Bank. Legal size is a carapace length greater than 76 mm. Pots that caught no lobsters are included as zeros.

Modelling: catch inside vs outside the National Park Zone

Catch per pot was modelled with a Tweedie generalised linear model (log link), fitted in `glmmTMB`:

`catch ~ status * year + depth_z`

where **status** is inside or outside the National Park Zone, **year** is a factor, and **depth_z** is pot depth standardised to zero mean and unit variance. Predicted means and standard errors for each status-by-year combination were obtained with **emmeans**, holding depth at its mean.

Six responses were modelled with this specification: legal catch, sublegal catch, legal males, legal females, males of 100 mm or more, and all lobsters combined. These are not independent tests of each other. Legal male and legal female catch together make up the legal catch, apart from 3 legal animals whose sex was not recorded; the large males are a subset of the legal males (137 of 281); and the all lobsters response is the legal and sublegal responses added together.

Two limitations to read these results against

The random effect for site is not included. The intended specification was `catch ~ status * year + depth_z + (1 | site)`, where **site** is the string (line) of pots. Neither survey records a string identifier, and the 2026 pot numbering does not follow position along the line, so strings cannot be recovered reliably from the data. Because pots within a string are not independent, the standard errors and p-values below are anti-conservative. The predicted means are much less affected than their uncertainty, so the plots are indicative while the significance tests are provisional.

Status is spatially confounded with survey area. The National Park Zone is a single contiguous block that was fished on its own day in both years. Comparing inside with outside is therefore partly comparing the middle area with the northern and southern areas, and any habitat difference between those areas is not separable from the effect of protection.

Table 4: Predicted western rock lobster catch per pot from the Tweedie generalised linear model, by size class, zoning status and survey year, with standard errors and asymptotic 95% confidence limits. Predictions are made with depth held at its mean. NPZ = National Park Zone.

Size class	Status	Year	Per pot	SE	Lower CL	Upper CL
Legal	Outside NPZ	2025	3.10	0.37	2.45	3.91
Legal	Inside NPZ	2025	3.79	0.55	2.84	5.04
Legal	Outside NPZ	2026	0.95	0.18	0.66	1.37
Legal	Inside NPZ	2026	1.64	0.32	1.12	2.40
Sublegal	Outside NPZ	2025	1.97	0.35	1.38	2.80
Sublegal	Inside NPZ	2025	0.62	0.20	0.32	1.18
Sublegal	Outside NPZ	2026	2.03	0.38	1.41	2.92
Sublegal	Inside NPZ	2026	0.62	0.20	0.32	1.17
Legal male	Outside NPZ	2025	1.18	0.15	0.92	1.52
Legal male	Inside NPZ	2025	1.51	0.23	1.13	2.03

Size class	Status	Year	Per pot	SE	Lower CL	Upper CL
Legal male	Outside NPZ	2026	0.36	0.08	0.23	0.56
Legal male	Inside NPZ	2026	0.75	0.15	0.51	1.12
Legal female	Outside NPZ	2025	1.88	0.25	1.45	2.44
Legal female	Inside NPZ	2025	2.27	0.37	1.65	3.12
Legal female	Outside NPZ	2026	0.58	0.12	0.38	0.88
Legal female	Inside NPZ	2026	0.90	0.20	0.57	1.40
Male 100 mm+	Outside NPZ	2025	0.42	0.08	0.29	0.61
Male 100 mm+	Inside NPZ	2025	0.97	0.17	0.69	1.36
Male 100 mm+	Outside NPZ	2026	0.14	0.05	0.08	0.27
Male 100 mm+	Inside NPZ	2026	0.38	0.10	0.23	0.64
All lobsters	Outside NPZ	2025	5.19	0.65	4.06	6.63
All lobsters	Inside NPZ	2025	4.58	0.78	3.28	6.40
All lobsters	Outside NPZ	2026	3.04	0.47	2.25	4.10
All lobsters	Inside NPZ	2026	2.30	0.48	1.54	3.46




Figure 11: Predicted western rock lobster catch per pot inside and outside the National Park Zone across the size progression from sublegal to legal to large males, from the Tweedie generalised linear model. Points are predicted means with standard errors, with depth held at its mean. Note that each panel has its own y axis scale. Legal size is a carapace length greater than 76 mm. NPZ = National Park Zone.




Figure 12: Predicted western rock lobster catch per pot inside and outside the National Park Zone for all five size class models, including legal males and legal females separately, from the Tweedie generalised linear model. Points are predicted means with standard errors, with depth held at its mean. Note that each panel has its own y axis scale. NPZ = National Park Zone.




Figure 13: Predicted total western rock lobster catch per pot, legal and sublegal combined, inside and outside the National Park Zone, from the Tweedie generalised linear model. Points are predicted means with standard errors, with depth held at its mean. This response is the sum of the legal and sublegal responses shown above rather than an independent size class. NPZ = National Park Zone.




Figure 14: Predicted western rock lobster catch per pot inside and outside the National Park Zone by size class and survey year, from the Tweedie generalised linear model, shown as bars with standard errors. Depth is held at its mean and each panel has its own y axis scale. NPZ = National Park Zone.




Figure 15: Observed western rock lobster catch per pot by size class, zoning status and survey year, showing the spread of the raw counts behind the modelled means. Boxes show the median and interquartile range with whiskers to 1.5 times the interquartile range. All panels share a y axis so that size classes can be compared against each other. Pots that caught no lobsters are included as zeros. NPZ = National Park Zone.

Table 5: Coefficients of the Tweedie generalised linear model of western rock lobster catch per pot, on the log scale, with standard errors, z statistics and p values. Reference levels are outside the National Park Zone in 2025, so the status term is the effect of protection in 2025 and the interaction term is the change in that effect by 2026. NPZ = National Park Zone.

Size class	Term	Estimate	SE	z	p
Legal	(Intercept)	1.131	0.119	9.50	<0.001
Legal	statusInside NPZ	0.200	0.187	1.07	0.285
Legal	year2026	-1.185	0.222	-5.34	<0.001
Legal	depth_z	0.249	0.084	2.97	0.003

Size class	Term	Estimate	SE	z	p
Legal	statusInside NPZ:year2026	0.346	0.328	1.06	0.290
Sublegal	(Intercept)	0.677	0.179	3.78	<0.001
Sublegal	statusInside NPZ	-1.161	0.376	-3.09	0.002
Sublegal	year2026	0.030	0.257	0.12	0.908
Sublegal	depth_z	-0.255	0.134	-1.91	0.057
Sublegal	statusInside NPZ:year2026	-0.029	0.533	-0.05	0.956
Legal male	(Intercept)	0.168	0.128	1.32	0.187
Legal male	statusInside NPZ	0.245	0.196	1.25	0.210
Legal male	year2026	-1.184	0.257	-4.61	<0.001
Legal male	depth_z	0.218	0.085	2.55	0.011
Legal male	statusInside NPZ:year2026	0.485	0.359	1.35	0.176
Legal female	(Intercept)	0.634	0.132	4.80	<0.001
Legal female	statusInside NPZ	0.186	0.207	0.90	0.369
Legal female	year2026	-1.182	0.252	-4.69	<0.001
Legal female	depth_z	0.268	0.095	2.83	0.005
Legal female	statusInside NPZ:year2026	0.253	0.375	0.68	0.499
Male 100 mm+	(Intercept)	-0.863	0.192	-4.49	<0.001
Male 100 mm+	statusInside NPZ	0.834	0.256	3.26	0.001
Male 100 mm+	year2026	-1.072	0.378	-2.83	0.005
Male 100 mm+	depth_z	0.233	0.112	2.09	0.037
Male 100 mm+	statusInside NPZ:year2026	0.143	0.490	0.29	0.771
All lobsters	(Intercept)	1.647	0.125	13.17	<0.001
All lobsters	statusInside NPZ	-0.124	0.211	-0.59	0.557
All lobsters	year2026	-0.536	0.198	-2.71	0.007
All lobsters	depth_z	0.098	0.090	1.09	0.277
All lobsters	statusInside NPZ:year2026	-0.152	0.332	-0.46	0.647

Table 6: Likelihood ratio test of the status by year interaction for each size class, comparing the full model with an additive model in which the difference between zones is held constant across years. A significant result would indicate that the difference between zones changed between 2025 and 2026.

Size class	Chi squared	df	p
Legal	1.116	1	0.291
Sublegal	0.003	1	0.956
Legal male	1.834	1	0.176
Legal female	0.455	1	0.500
Male 100 mm+	0.085	1	0.770
All lobsters	0.210	1	0.647

Data notes

- Pots that were lost, had a missing trap door, or were flagged in the field as not countable are excluded from catch rates but retained in the pot table.
- Pots that caught no lobsters are carried through the analysis as zeros.
- Set times were not recorded in 2026, so soak time is available for 2025 only and catch is reported per pot rather than per pot-hour.
- 2026 includes six extra pots labelled F3, F3-2 and FIII. These are kept as distinct pots rather than merged with the numbered pots.
- Three 2026 pots lost GPS fix on retrieval; corrected positions were read from the field comments.
- One 2026 pot was logged at 0 m depth, which is not a possible seabed depth here. It is treated as missing and drops from the models.
- Non-lobster catch from 2026 is written to a separate bycatch file and is not included in these results.
- One 2025 measurement of 24 mm (National Park Zone, pot 2) sits well below the next smallest lobster at 50 mm and is likely a data entry error. It has been retained pending confirmation.