

Appendix C - South-west Corner Marine Park (western arm) Data Analysis Report

South-west Marine Parks Network

Table of contents

Data Analysis	1
Benthic ecosystem extent	2
Demersal fish assemblage	5

Data Analysis

Statistical methods and model selection results underpinning the natural values reported in Appendix A.

Benthic ecosystem extent

Table C 1.1: Habitat model selection results. Top generalised additive models (GAMs) for predicting the probability of occurrence of key habitats from full subset analyses. Differences between the lowest reported corrected Akaike Information Criterion (ΔAICc), AICc weight (ωAICc), variance explained (R^2) and effective degrees of freedom (EDF) are reported for model comparison. Model selection was based on the most parsimonious model (fewest variables - shown in bold) within two units of the lowest AICc. Benthic samples are pooled across survey years, so survey year was not offered as a candidate predictor.

Response	Model	ΔAICc	ωAICc	R^2	EDF
Macroalgae	detrended+aspect+Z	0	0.707	0.63419	6.95
	detrended+roughness+aspect+Z	1.762	0.293	0.63419	7.81
Sand	detrended+roughness+as- pect+Z	0	1	0.50357	8.84
	detrended+roughness+as- pect+Z	0	0.997	0.38707	8.10
Rock	detrended+roughness+as- pect+Z	0	1	0.18945	8.49
	detrended+roughness+as- pect+Z	0	1	0.39016	8.96
Sessile invertebrates	detrended+roughness+as- pect+Z	0	1	0.39016	8.96
	detrended+roughness+as- pect+Z	0	1	0.45613	6.97
Reef	detrended+roughness+as- pect+Z	0	1	0.45613	6.97




Figure C 1.1: Variable importance scores for predicting the probability of occurrence of main habitat categories. Response variables included in the most parsimonious top model are indicated (X), with the colour gradient representing positive (red), zero (white) and negative (blue) relationships. Aspect is a circular variable, so the direction of its relationship should be read from the response curves rather than the colour gradient.




Figure C 1.2: Plots of the most parsimonious model found to predict the probability of occurrence of key habitat types. Benthic samples from 2020, 2021, 2023, 2024 and 2025 are pooled, because the campaigns sample different areas and a survey-year term would read as a spatial difference rather than change through time. Individual panes display the predicted relationships between model covariates and the probability of occurrence of each habitat class. Solid lines represent predicted Generalized Additive Model (GAM) fits with other variables held constant at their mean value. Dashed lines represent upper and lower standard error bounds around the prediction, and points represent the observed data.

Demersal fish assemblage

Table C 2.1: Demersal fish model selection results. Top generalised additive models (GAMs) for predicting the species and abundance distribution of metrics of interest from full subset analyses. Differences between the lowest reported corrected Akaike Information Criterion (ΔAICc), AICc weight (ωAICc), variance explained (R^2) and effective degrees of freedom (EDF) are reported for model comparison. Model selection was based on the best model (fewest variables - shown in bold) within two units of the lowest AICc. *Large Reef Fish Index representing the biomass of all fish greater than 20 cm (B20) has been adapted here for stereo-BRUV data by removing sharks and rays.

Response	Model	ΔAICc	ωAICc	AICc	R^2	EDF
Species richness	aspect+reef+status	0	0.667	3529.78	0.36545	5.22
	reef+status	1.386	0.333	3531.16	0.36157	3.99
Total abundance	aspect+Z+year	0	0.726	6707.14	0.33165	7.79
Large Reef Fish Index*	reef+year+status	0	0.495	11486.86	0.12857	6.30
Reef Fish Thermal Index	aspect+reef+year+status	0.011	0.492	11486.87	0.12857	7.30
	aspect+Z+year+status	0	0.153	1723.47	0.04549	7.00
	Z+year+status	0.001	0.153	1723.47	0.04549	6.00
	roughness+aspect+Z+status	0.504	0.119	1723.98	0.03977	5.16
	roughness+Z+status	0.505	0.119	1723.98	0.03977	4.16
	aspect+Z+year	1.713	0.065	1725.19	0.04012	6.00
	Z+year	1.726	0.065	1725.20	0.04014	5.01




Figure C 2.1: Variable importance scores for predicting the probability of demersal fish metrics. Response variables included in the most parsimonious top model are indicated (X), with the colour gradient representing positive (red), zero (white) and negative (blue) relationships. Aspect is a circular variable, so the direction of its relationship should be read from the response curves rather than the colour gradient. *Large Reef Fish Index representing the biomass of all fish greater than 20 cm (B20) has been adapted here for stereo-BRUV data by removing sharks and rays.




Figure C 2.2: Plots of the best model found to predict the probability of occurrence of key fish groups. Individual panes display the predicted relationships between model covariates and each demersal fish metric. Solid lines represent predicted Generalized Additive Model (GAM) fits with other variables held constant at their mean value, with year held at 2020 and zone status held at areas open to fishing. These terms are held only where model selection retained them: year was not retained for Species richness and zone status was not retained for Total abundance and Reef Fish Thermal Index. Dashed lines represent upper and lower standard error bounds around the prediction, and points represent the observed data from all survey years and zones. *Large Reef Fish Index representing the biomass of all fish greater than 20 cm (B20) has been adapted here for stereo-BRUV data by removing sharks and rays.