

hasklib: Stochastic Process Derivations

Hubert Kasinski

August 6, 2026

Abstract

This document collects the stochastic-calculus results that the stochastic layer of **hasklib** relies on: from the probability-space setup, Brownian motion, Itô SDE, to Geometric Brownian Motion, Ornstein–Uhlenbeck, Arithmetic Brownian Motion, Cox–Ingersoll–Ross, and numerical schemes. Standard measure-theoretic probability is assumed; see Shreve (2004) or Williams (1991) for construction.

Contents

1	Foundations: Itô Calculus and the SDE Framework	2
1.1	Probability space and Brownian motion	2
1.2	Quadratic variation	3
1.3	The Itô stochastic differential equation	4
1.4	Itô's lemma	5
2	Geometric Brownian Motion	7
3	The Ornstein–Uhlenbeck Process	8
4	Arithmetic Brownian Motion	9
5	The Cox–Ingersoll–Ross Process	10
6	Numerical Schemes: Euler–Maruyama and Milstein	11

Chapter 1

Foundations: Itô Calculus and the SDE Framework

1.1 Probability space and Brownian motion

We work on a filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ satisfying the usual conditions. The filtration $\{\mathcal{F}_t\}$ represents the information available up to time t .

Definition 1.1 (Brownian motion).

A standard Brownian motion $\{W_t\}_{t \geq 0}$ is a process with

1. $W_0 = 0$ almost surely;
2. Independent increments: for $s < t < u$, $W_u - W_t$ is independent of $W_t - W_s$;
3. Gaussian increments: $W_t - W_s \sim \mathcal{N}(0, t - s)$;
4. Continuous sample paths.

Remark 1.1.

In code, property (3) is what `NormalRng::draw()` supplies: a step of length Δt uses $W_{t+\Delta t} - W_t = \sqrt{\Delta t} Z$ with $Z \sim \mathcal{N}(0, 1)$. This substitution is the single bridge between the continuous theory here and the discrete simulation code.

One might notice (2) and (4) pull in different directions. Independent increments mean each step of the path is a fresh draw, carrying no memory of the past — nothing about $\mathcal{F}_t$ constrains $W_{t+h} - W_t$. Continuity, on the other hand, means the path never jumps: W_{t+h} must stay close to W_t as $h \rightarrow 0$. It is not obvious a single process can do both at once, and the price of doing so turns out to be that the path is extremely rough and nowhere differentiable.

Fixing t and considering the difference quotient over a step $h > 0$. By the Gaussian-increment property, $W_{t+h} - W_t \sim \mathcal{N}(0, h)$, so the difference quotient

$$\frac{W_{t+h} - W_t}{h}$$

is itself Gaussian with mean 0 and variance

$$\text{Var}\left(\frac{W_{t+h} - W_t}{h}\right) = \frac{1}{h^2} \text{Var}(W_{t+h} - W_t) = \frac{h}{h^2} = \frac{1}{h},$$

using that dividing a random variable by a constant h scales its variance by $1/h^2$. Hence the standard deviation of the difference quotient is $1/\sqrt{h}$, which diverges as $h \downarrow 0$: the path moves

by order $\sqrt{h}$ over a step of order h , so no finite slope survives the limit. This informal argument is made rigorous by the Paley–Wiener–Zygmund theorem (Paley, Wiener & Zygmund 1933): with probability one, $t \mapsto W_t$ is nowhere differentiable. It also follows that the path has infinite first-order (total) variation on every interval, so integrals $\int f dW_s$ cannot be defined pathwise as Riemann–Stieltjes integrals. Its quadratic variation, however, is finite, and it is precisely this finite second-order variation that the Itô integral and Itô’s lemma are built on.

1.2 Quadratic variation

The property that distinguishes stochastic from ordinary calculus is that Brownian motion accumulates quadratic variation at a nonzero rate.

Proposition 1.1 (Quadratic variation of Brownian motion).

For a standard Brownian motion, the quadratic variation over $[0, T]$ satisfies

$$\langle W \rangle_T = \lim_{\|\Pi\| \rightarrow 0} \sum_k (W_{t_{k+1}} - W_{t_k})^2 = T \quad \text{in } L^2,$$

where Π is a partition of $[0, T]$. Informally this is written

$$(dW_t)^2 = dt.$$

Proof.

Write the partition as $\Pi = \{0 = t_0 < t_1 < \dots < t_n = T\}$ and abbreviate

$$\Delta t_k = t_{k+1} - t_k, \quad \Delta W_k = W_{t_{k+1}} - W_{t_k}, \quad S_\Pi = \sum_{k=0}^{n-1} (\Delta W_k)^2.$$

“Convergence in L^2 ” means the mean-square error vanishes: $\mathbb{E}[(S_\Pi - T)^2] \rightarrow 0$ as the mesh $\|\Pi\| := \max_k \Delta t_k \rightarrow 0$. We use only two facts about the increments, both directly from the definition of Brownian motion:

- Each $\Delta W_k \sim \mathcal{N}(0, \Delta t_k)$.
- Increments over disjoint intervals are independent.

We then calculate the mean of S_Π , i.e $\mathbb{E}[S_\Pi]$

Since $\mathbb{E}[\Delta W_k] = 0$, the second moment is the variance:

$$\mathbb{E}[(\Delta W_k)^2] = \text{Var}(\Delta W_k) = \Delta t_k.$$

Summing over k ,

$$\mathbb{E}[S_\Pi] = \sum_k \mathbb{E}[(\Delta W_k)^2] = \sum_k \Delta t_k = T.$$

This holds for every partition, not merely in the limit. Because the mean is exactly T , the mean-square error is the variance:

$$\mathbb{E}[(S_\Pi - T)^2] = \text{Var}(S_\Pi).$$

The whole statement then reduces to showing $\text{Var}(S_\Pi) \rightarrow 0$.

We then calculate the variance of a single term.

Normalising the increment as $\Delta W_k = \sqrt{\Delta t_k} Z$ with $Z \sim \mathcal{N}(0, 1)$, so $(\Delta W_k)^2 = \Delta t_k Z^2$ and

$$\text{Var}((\Delta W_k)^2) = (\Delta t_k)^2 \text{Var}(Z^2).$$

For a standard normal, $\mathbb{E}[Z^2] = 1$ and $\mathbb{E}[Z^4] = 3$, hence $\text{Var}(Z^2) = \mathbb{E}[Z^4] - (\mathbb{E}[Z^2])^2 = 3 - 1 = 2$. (The fourth moment is one Gaussian integration by parts: with the density ϕ satisfying $\phi'(z) = -z\phi(z)$, integrating $\int z^4\phi dz$ by parts gives $\mathbb{E}[Z^4] = 3\mathbb{E}[Z^2] = 3$.) Therefore,

$$\text{Var}((\Delta W_k)^2) = 2(\Delta t_k)^2.$$

Step 3 Variance of the sum, then let the mesh shrink.

The increments ΔW_k are independent, meaning the squared increments $(\Delta W_k)^2$ are independent too, and the variance of a sum of independent variables is the sum of the variances:

$$\text{Var}(S_\Pi) = \sum_k \text{Var}((\Delta W_k)^2) = 2 \sum_k (\Delta t_k)^2.$$

Bounding one factor of Δt_k in each term by the mesh:

$$\sum_k (\Delta t_k)^2 \leq \left(\max_k \Delta t_k \right) \sum_k \Delta t_k = \|\Pi\| T.$$

Combining,

$$\mathbb{E}[(S_\Pi - T)^2] = \text{Var}(S_\Pi) \leq 2 \|\Pi\| T \xrightarrow{\|\Pi\| \rightarrow 0} 0,$$

which is exactly the claimed L^2 convergence. $\square$

Remark 1.2.

The mean computation ($\mathbb{E}[S_\Pi] = T$) shows $(dW_t)^2 = dt$ is plausible; the variance computation shows it is usable. Note the per-step fluctuation is not negligible, $(\Delta W_k)^2$ has standard deviation $\sqrt{2}\Delta t_k$, the same order as its own mean Δt_k — so the cancellation is a genuine consequence of independence across steps, not something true term by term. Inside a sum (equivalently, an integral), the random $(\Delta W_k)^2$ may therefore be replaced by the deterministic Δt_k : this replacement is the entire content of the shorthand $(dW_t)^2 = dt$ used in the next section.

1.3 The Itô stochastic differential equation

Definition 1.2 (Itô SDE).

An Itô process X_t solves the stochastic differential equation

$$dX_t = a(t, X_t) dt + b(t, X_t) dW_t, \tag{1.1}$$

interpreted as the integral equation

$$X_t = X_0 + \int_0^t a(s, X_s) ds + \int_0^t b(s, X_s) dW_s,$$

where the second integral is an Itô integral. We call a the drift and b the diffusion coefficients.

Remark 1.3 (Correspondence with the code).

Equation (1.1) is exactly what the `StochasticProcess` base class encodes: `drift(t,x)` returns $a(t,x)$ and `diffusion(t,x)` returns $b(t,x)$. Every subsequent process (GBM, OU, CIR) is a choice of these two functions.

Assumption 1.1 (Existence and uniqueness).

We assume a and b satisfy the standard Lipschitz and linear-growth conditions guaranteeing a unique strong solution to (1.1); see Øksendal (2003), Thm. 5.2.1.

1.4 Itô's lemma

Itô's lemma is the chain rule for functions of an Itô process, and it is the engine behind every closed-form solution used in `hasklib` (the GBM exact sampler, the OU solution).

Theorem 1.1 (Itô's lemma, one dimension). *Let X_t solve (1.1) and let $\varphi(t, x) \in C^{1,2}$. Then*

$$d\varphi(t, X_t) = \left(\frac{\partial \varphi}{\partial t} + a \frac{\partial \varphi}{\partial x} + \frac{1}{2} b^2 \frac{\partial^2 \varphi}{\partial x^2} \right) dt + b \frac{\partial \varphi}{\partial x} dW_t, \quad (1.2)$$

Informal derivation.

This is the standard second-order Taylor argument. It is not a proof in the strict sense, as a rigorous version requires the construction of the Itô integral (Shreve 2004, Sec. 4.2.1), which is outside the scope of this document. The one idea is this: because dX carries a Brownian term of size $\sqrt{dt}$, a contribution that ordinary calculus discards as “second order” is in fact first order and must be kept.

Orders of magnitude. Over a step of length dt the increment dW_t has typical size $\sqrt{dt}$ (it is $\mathcal{N}(0, dt)$). Ranking the elementary products by their power of dt :

$$dt = O(dt), \quad dW_t = O(dt^{1/2}), \quad (dW_t)^2 = O(dt), \quad dt dW_t = O(dt^{3/2}), \quad (dt)^2 = O(dt^2).$$

Discard everything of order smaller than dt , and use proposition 1.1 to replace $(dW_t)^2$ by its in-sum limit dt . This gives the *Itô multiplication table*

$$(dW_t)^2 = dt, \quad dt dW_t = 0, \quad (dt)^2 = 0. \quad (1.3)$$

The last two entries are not literally zero; they are negligible next to dt and vanish in the mesh limit for exactly the reason the fluctuations did in section 1.2.

Taylor expansion. Expand $\varphi(t + dt, X_t + dX)$ about (t, X_t) to second order:

$$d\varphi = \frac{\partial \varphi}{\partial t} dt + \frac{\partial \varphi}{\partial x} dX + \frac{1}{2} \frac{\partial^2 \varphi}{\partial t^2} (dt)^2 + \frac{\partial^2 \varphi}{\partial t \partial x} dt dX + \frac{1}{2} \frac{\partial^2 \varphi}{\partial x^2} (dX)^2 + \dots$$

Substitute the SDE $dX = a dt + b dW_t$ and square:

$$(dX)^2 = a^2 (dt)^2 + 2ab dt dW_t + b^2 (dW_t)^2.$$

Apply (1.3). In $(dX)^2$ the first two terms die and only $b^2 (dW_t)^2 = b^2 dt$ survives:

$$(dX)^2 = b^2 dt.$$

The two remaining second-order terms of the expansion vanish for the same reason:

$$\frac{1}{2} \frac{\partial^2 \varphi}{\partial t^2} (dt)^2 = 0, \quad \frac{\partial^2 \varphi}{\partial t \partial x} dt dX = \frac{\partial^2 \varphi}{\partial t \partial x} (a (dt)^2 + b dt dW_t) = 0.$$

What remains is

$$d\varphi = \frac{\partial \varphi}{\partial t} dt + \frac{\partial \varphi}{\partial x} dX + \frac{1}{2} \frac{\partial^2 \varphi}{\partial x^2} b^2 dt.$$

Substitute $dX = a dt + b dW_t$ once more and collect the dt and dW_t parts:

$$d\varphi = \left(\frac{\partial \varphi}{\partial t} + a \frac{\partial \varphi}{\partial x} + \frac{1}{2} b^2 \frac{\partial^2 \varphi}{\partial x^2} \right) dt + b \frac{\partial \varphi}{\partial x} dW_t,$$

which is (1.2). □

Remark 1.4. Had X been a smooth (finite-variation) path, $(dX)^2$ would have been genuinely $O((dt)^2)$ and dropped out, leaving the ordinary chain rule $d\varphi = \frac{\partial\varphi}{\partial t} dt + \frac{\partial\varphi}{\partial x} dX$. The whole gap between stochastic and ordinary calculus is the single surviving term $\frac{1}{2}b^2\frac{\partial^2\varphi}{\partial x^2} dt$, and it survives only because $(dW_t)^2 = dt$ rather than 0 — i.e. because Brownian motion has nonzero quadratic variation. Every closed-form result in `hasklib` that “corrects a drift by half a variance” (the $-\frac{1}{2}\sigma^2$ in the GBM log-solution, the Itô term in the OU solution) is exactly this one term.

Remark 1.5. The single downstream consequence to keep in view: applying theorem 1.1 with $f(t, x) = \ln x$ to geometric Brownian motion linearises the SDE and yields the exact terminal law sampled by `GBM::sample_terminal`.

Chapter 2

Geometric Brownian Motion

Chapter 3

The Ornstein–Uhlenbeck Process

Chapter 4

Arithmetic Brownian Motion

Chapter 5

The Cox–Ingersoll–Ross Process

Chapter 6

Numerical Schemes: Euler–Maruyama and Milstein

Bibliography

- Øksendal, B. (2003), *Stochastic Differential Equations: An Introduction with Applications*, 6th edn, Springer, Berlin.
- Paley, R. E. A. C., Wiener, N. & Zygmund, A. (1933), ‘Notes on random functions’, *Mathematische Zeitschrift* **37**(1), 647–668.
- Shreve, S. E. (2004), *Stochastic Calculus for Finance II: Continuous-Time Models*, Springer Finance, Springer, New York.
- Williams, D. (1991), *Probability with Martingales*, Cambridge University Press, Cambridge.