

Arithmetic Brownian Motion: Exact Solution

hasklib mathematical companion

Hubert

July 20, 2026

Abstract

This note derives the exact terminal law of arithmetic Brownian motion (ABM), the process underlying Shreve (2004)’s classical treatment and, historically, Bachelier’s original 1900 model of a stock price. Unlike GBM and OU, neither Itô’s lemma nor an integrating factor is needed: both coefficients are already constant, so the SDE integrates directly.

1 The ABM SDE

Definition 1 (Arithmetic Brownian motion). X_t follows arithmetic Brownian motion with drift $\mu \in \mathbb{R}$ and volatility $\sigma > 0$ if it solves

$$dX_t = \mu dt + \sigma dW_t, \quad X_0 \in \mathbb{R}. \quad (1)$$

In the notation of `ito_foundations.tex` this is $a(t, x) = \mu$, $b(t, x) = \sigma$ — both constant, with no x -dependence at all.

Remark 1 (Correspondence with the code). `ABM::drift` returns the constant μ and `ABM::diffusion` returns the constant σ , independent of x in both cases. Since neither coefficient depends on x , the Lipschitz and linear-growth conditions of the existence–uniqueness assumption in `ito_foundations.tex` hold trivially — this is the simplest possible case of that assumption.

2 Exact solution

Both GBM and OU needed a substitution to remove an X_t -dependence from the coefficients before the SDE could be integrated: $\ln x$ for GBM, the integrating factor $e^{\kappa t}$ for OU. Here there is nothing to remove — μ and σ are already constants — so the general relation from `ito_foundations.tex`,

$$X_t = X_0 + \int_0^t a(s, X_s) ds + \int_0^t b(s, X_s) dW_s,$$

integrates immediately once $a = \mu$ and $b = \sigma$ are substituted.

Theorem 1 (Exact terminal law of ABM). *Let X_t solve (1). Then for any $T > 0$,*

$$X_T = X_0 + \mu T + \sigma W_T, \quad (2)$$

where $W_T \sim \mathcal{N}(0, T)$, so

$$X_T \sim \mathcal{N}(X_0 + \mu T, \sigma^2 T). \quad (3)$$

Proof. Substituting $a(s, X_s) = \mu$ and $b(s, X_s) = \sigma$,

$$X_T = X_0 + \int_0^T \mu \, ds + \int_0^T \sigma \, dW_s = X_0 + \mu T + \sigma \int_0^T dW_s.$$

The remaining integral is $\int_0^T dW_s = W_T - W_0 = W_T$, giving (2). Since $X_0 + \mu T$ is a fixed number and $W_T \sim \mathcal{N}(0, T)$, the affine transformation $X_0 + \mu T + \sigma W_T$ is itself Gaussian with mean $X_0 + \mu T$ and variance $\sigma^2 \text{Var}(W_T) = \sigma^2 T$, giving (3). $\square$

Remark 2. Equation (3) is sampled with a single normal draw: generate $Z \sim \mathcal{N}(0, 1)$ and set $X_T = X_0 + \mu T + \sigma \sqrt{T} Z$. As with `GBM::sample_terminal` and `OU::sample_terminal`, no time-stepping is needed — this is what `ABM::sample_terminal` implements.

3 Moments and the relation to OU

Theorem 1 already gives the moments directly, with no separate lemma needed:

$$\mathbb{E}[X_T] = X_0 + \mu T, \quad \text{Var}(X_T) = \sigma^2 T.$$

Unlike OU, the variance grows without bound as $T \rightarrow \infty$: there is no mean-reverting pull holding the process in, so uncertainty simply accumulates. Correspondingly, $\mathbb{E}[X_T]$ grows in a straight line rather than curving toward a long-run level — there is no θ here for it to curve toward.

Remark 3 (ABM as a degenerate OU process). Setting $\kappa = 0$ in the OU drift $\kappa(\theta - X_t)$ removes it entirely, leaving $dX_t = \sigma dW_t$ — driftless Brownian motion. ABM generalises this by adding back a constant drift μ that is *not* tied to any mean-reversion mechanism. Equivalently: ABM is what OU degenerates into once mean-reversion is switched off, plus an unconstrained drift term OU's $\kappa = 0$ case does not have on its own.

References

Shreve, S. E. (2004), *Stochastic Calculus for Finance II: Continuous-Time Models*, Springer Finance, Springer, New York.