

THE TRAVELING SYSTEM

*A Framework for Galactic Forcing of Terrestrial Climate
and Biosphere Through Heliospheric Variability*

John Carter

Independent Researcher | ORCID: 0009-0004-1363-304X

Niagara Region, Ontario, Canada

Preprint submitted April 2026

ABSTRACT

The dominant framework for climate forcing attributes observed planetary warming primarily to anthropogenic CO₂ and, on geological timescales, to Milankovitch orbital cycles. While both mechanisms are well-evidenced, we propose they are insufficient as complete explanations because they omit a fundamental variable: the solar system is not stationary. Our solar system orbits the galactic center at approximately 828,000 km/h, oscillates above and below the galactic plane on a roughly 30-million-year cycle, and traverses energetically distinct regions of the interstellar medium (ISM) on timescales of thousands to hundreds of millions of years. We term this integrated dynamic The Traveling System. Drawing on published findings in heliospheric astrophysics, cosmic ray climate science, atmospheric electricity, archaeoastronomy, and astrobiology, we argue that variations in the galactic environment of the solar system—specifically ISM density, cosmic ray flux, and interstellar magnetic field structure—modulate the heliosphere, alter galactic cosmic ray (GCR) flux reaching Earth’s atmosphere, perturb the global atmospheric electric circuit (GEC) and Schumann resonance state, and thereby drive climate and biosphere forcing at timescales inaccessible to orbital or anthropogenic models alone. Geological evidence confirms that Earth made direct contact with the ISM approximately 2–3 and 6–7 million years ago when the heliosphere collapsed to sub-AU scales during a cold cloud encounter, exposing the planet to radiation orders of magnitude beyond any recorded solar energetic particle event. We present a multi-scale forcing hierarchy, a set of falsifiable predictions, and a research agenda for the integration of heliospheric boundary data with terrestrial climate archives.

Keywords: *interstellar medium, heliosphere, galactic cosmic rays, cosmoclimatology, atmospheric electric circuit, Schumann resonance, panspermia, paleoheliosphere, Local Interstellar Cloud, solar journey*

1. INTRODUCTION

The climate science community has made substantial progress identifying and quantifying the mechanisms that drive Earth’s temperature over geological and historical timescales. Anthropogenic greenhouse gas accumulation, variations in solar irradiance, volcanic aerosol

forcing, and Milankovitch orbital cycles collectively form the standard explanatory toolkit. These mechanisms are real, well-supported by data, and scientifically productive. The present paper does not contest them.

What we propose is that this toolkit is incomplete. Specifically, it treats the solar system as a fixed object in a fixed environment, when in fact the solar system is in continuous, high-velocity motion through a galactic medium that varies substantially in density, magnetic field structure, cosmic ray flux, and chemical composition. The consequences of this motion for Earth's climate and biosphere have received increasing attention in heliospheric astrophysics [1–4] but remain almost entirely absent from mainstream climate modeling and science communication.

The question we are asking is simple: as our solar system moves through the galaxy, what is the interstellar environment doing to us, and how would we measure it? This is not a challenge to the CO₂ model. It is an expansion of the frame. We call this expanded frame The Traveling System.

This paper proceeds as follows. Section 2 establishes the basic kinematics of the solar system's galactic journey. Section 3 reviews galactic cosmic ray climate forcing. Section 4 examines ancient calendrical systems as unwitting long-cycle astronomical data archives. Section 5 describes the atmospheric electric circuit as the primary terrestrial receiver of external forcing. Section 6 addresses biological and chemical presence in the interstellar medium. Section 7 inventories additional forcing variables not captured by existing models. Section 8 presents a unified multi-scale forcing hierarchy. Section 9 offers falsifiable predictions. Section 10 concludes.

2. THE SOLAR SYSTEM AS A TRAVELING BODY

2.1 Galactic Kinematics

The solar system orbits the galactic center at approximately 828,000 km/h, completing one full orbit roughly every 225–250 million years at a distance of about 28,000 light-years from the core [5]. This orbit is not a simple circle. The solar system also oscillates above and below the galactic midplane at a period of approximately 30 million years and with an amplitude of roughly 250 light-years [6]. We crossed the midplane approximately 2 million years ago and are currently moving upward through the disk. Additionally, as the solar system traverses the galaxy it periodically crosses the denser spiral arms, which harbor the majority of star formation activity and supernova events, on a timescale of approximately 143 million years [7].

Each of these motions carries the solar system through regions of substantially different interstellar conditions. The galactic disk is not uniform. It contains molecular clouds, supernova remnants, superbubbles, cold neutral clouds, warm ionized plasma, and regions of varying magnetic field strength and orientation. The heliosphere—the solar wind bubble that contains our planetary system—responds dynamically to each of these environments.

2.2 The Local Interstellar Environment

At present, the solar system is embedded within the Local Bubble, a cavity in the galactic interstellar medium roughly 1,000 light-years across, formed by a series of supernovae over the past 10–20 million years [8]. Within the Local Bubble we currently occupy the Local Interstellar Cloud (LIC), also called the Local Fluff—a warm, partially ionized cloud approximately 30 light-

years across [9]. Voyager 2 measurements indicate the LIC magnetic field is substantially stronger than previously estimated, at 370–550 picoteslas [9].

Of immediate significance: it is estimated that the solar system entered the LIC within the past 44,000–150,000 years and will exit within the next 1,900 to 20,000 years [9, 10]. We are approaching a cloud boundary. What lies beyond—specifically, the transition zone between the LIC and the neighboring G-Cloud—is not yet precisely characterized. This transition represents a currently unmeasured change in our interstellar environment.

2.3 Geological Evidence of Past ISM Contact

The most direct evidence that our galactic journey has Earth consequences comes from the isotope record. Iron-60 (^{60}Fe) and plutonium-244 (^{244}Pu) have been found in deep-sea sediments, ferromanganese crusts, Antarctic ice, and lunar soil samples at anomalous concentrations approximately 2–3 and 6–7 million years ago [11, 12]. These isotopes have short enough half-lives that their presence cannot be attributed to primordial solar system material. They must have an external source.

Magnetohydrodynamic modeling by Opher et al. [13] demonstrates that the solar system’s trajectory 2–3 million years ago intersected the Local Ribbon of Cold Clouds in the constellation Lynx—a dense cold cloud structure with neutral hydrogen density approximately $3,000\text{ cm}^{-3}$. At this density, ram pressure from the incoming ISM material collapses the heliosphere to approximately 0.22 AU—well inside the orbit of Earth at 1 AU. During such a collapse, Earth is no longer inside the heliosphere. It is directly exposed to the ISM.

During the months of each year when Earth’s orbit carried it through the collapsed heliosheath rather than outside it entirely, it experienced what Opher et al. term Heliospheric Energetic Particles (HEPs). Their modeling indicates HEP flux at least two orders of magnitude more intense than the 2003 Halloween solar energetic particle event—the largest ever recorded in the space era [13]. This is not a subtle perturbation. It is a planetary-scale radiation event of months’ duration, recurring annually for the duration of the cloud crossing.

Critically, this cold cloud encounter at 2.58 million years ago coincides with the Gauss-Matuyama geomagnetic reversal, during which Earth’s internal magnetic shield temporarily collapsed [13]. The heliospheric shield and the geomagnetic shield failed simultaneously. Whatever climate and biological consequences followed, they occurred in the context of maximum external forcing and minimum internal protection.

The oxygen isotope record from deep-sea foraminifera shows global cooling events at 13–14, 6–7, and 2–3 million years ago [14], all periods that overlap with hypothesized ISM encounters. The forcing mechanisms behind these cooling episodes remain, in the authors’ words, “poorly understood” [13]. The Traveling System framework proposes a candidate mechanism.

3. GALACTIC COSMIC RAY CLIMATE FORCING

3.1 The Svensmark Mechanism

Henrik Svensmark [15, 16] demonstrated that galactic cosmic ray (GCR) flux correlates with low-cloud cover on Earth with a correlation coefficient of approximately 0.92 over a 22-year

observational period. The proposed mechanism is as follows: GCRs ionize molecules in the lower troposphere, releasing electrons that catalyze the formation of sulfuric acid aerosol clusters, which serve as cloud condensation nuclei (CCN). More GCRs produce more CCN, more low-cloud cover, greater planetary albedo, and cooler surface temperatures. The inverse relationship between solar activity and GCR flux means that during periods of low solar magnetic activity—when the solar wind is weaker and the heliosphere less effective as a cosmic ray shield—more GCRs reach Earth and the climate cools [15].

The CERN CLOUD experiment confirmed that ionization from GCRs accounts for approximately one third of all atmospheric particle formation and significantly enhances aerosol nucleation in the mid-troposphere and above [17, 18]. The experiment also demonstrated that the vapors previously assumed to explain all lower-atmosphere aerosol formation account for only a small fraction of observations even with cosmic ray enhancement, indicating that our understanding of nucleation chemistry remains incomplete [18]. Kirkby and Carslaw, writing in the foundational volume *Solar Journey* [1], state explicitly: “On longer time scales, the GCR intensity is also affected by the geomagnetic field and by the galactic environment, whereas solar irradiance/UV is not. There is suggestive evidence linking the climate both with the geomagnetic field and with the galactic environment.”

3.2 The Galactic Scale: Shaviv and the Spiral Arm Signal

Nir Shaviv [7, 19] extended the cosmic ray climate connection from the solar cycle to the galactic scale. Using iron meteorite exposure ages as a proxy for historical GCR flux and comparing this record to paleoclimate temperature reconstructions over 500 million years, Shaviv and Veizer [19] identified a correlation between the solar system’s spiral arm passages—which occur on a roughly 143-million-year cycle—and the timing of ice age epochs on Earth. Spiral arms harbor dense star formation and abundant supernovae, both of which elevate local GCR flux by approximately an order of magnitude compared to inter-arm regions [7]. Shaviv’s framework thus provides a mechanism for climate forcing on the longest geological timescales that is entirely independent of terrestrial or solar factors.

The distinction between solar-cycle GCR modulation and galactic-scale GCR modulation is important. Solar cycle variation of GCR flux is approximately 15% over 11 years. Spiral arm passage variation is predicted to be approximately tenfold—an order of magnitude larger—and operates over hundreds of millions of years [7]. The CLOUD experiment finding that solar-cycle GCR changes are insufficient to drive significant climate change in today’s polluted atmosphere does not constrain the galactic scale effect, because the mechanisms operate at fundamentally different amplitudes and timescales.

4. ANCIENT CALENDRICIAL SYSTEMS AS LONG-CYCLE DATA ARCHIVES

An underexplored evidentiary thread runs through the archaeoastronomical record. Multiple ancient civilizations independently developed calendrical systems tracking astronomical cycles of remarkable precision and extraordinary duration. The Maya calculated the solar year as 365.242 days—modern measurement is 365.2425 days—and the lunar month as 29.5308 days, compared to the modern value of 29.53059 days [20]. The Dresden Codex contains astronomical tables stretching approximately 7,000 years into the future. The Long Count calendar encodes a

cycle of approximately 5,125 years, with evidence that the Maya understood this as one of multiple repeating cycles of far greater duration [21].

The precession of the equinoxes—a 26,000-year cycle arising from Earth’s axial wobble—was apparently known to multiple ancient civilizations under different names and encoded in myth, monument, and calendar. Giorgio de Santillana and Hertha von Dechend, in *Hamlet’s Mill* [22], documented that the folklore and myth of more than thirty ancient cultures encodes long astronomical cycles including the precessional Great Year, often associated with alternating ages of warmth and cold. The authors argue that ancient myth was not metaphor but functional scientific language describing observed cyclical phenomena.

We do not propose that ancient observers understood the heliospheric mechanisms described in this paper. We propose something more modest: that civilizations conducting systematic astronomical observation over multiple generations would inevitably record correlations between celestial cycle positions and terrestrial conditions, including climate. Their calendars and myths may therefore constitute the earliest proxy archive for long-cycle forcing events, encoded in the only data storage system available to pre-literate cultures: story.

The 26,000-year precessional cycle falls within the range of heliospheric variability timescales. The vertical oscillation of the solar system above and below the galactic plane occurs on a 30-million-year period, but with local ISM density variations at sub-million-year scales. The LIC transition that is currently underway will complete within a few thousand years—within the precessional cycle’s timeframe. The question of whether precessional position correlates with ISM boundary conditions at sub-cycle resolution is currently unaddressed in the literature and represents a promising research direction.

5. THE ATMOSPHERIC ELECTRIC CIRCUIT AS PLANETARY RECEIVER

5.1 Structure of the Global Electric Circuit

The global atmospheric electric circuit (GEC) is a continuous planetary-scale electrical system driven primarily by thunderstorm activity—approximately 2,000 active storms generating roughly 50 lightning discharges per second globally [23]. The GEC maintains a potential difference of approximately 300,000 volts between Earth’s surface and the ionosphere, with fair-weather return current flowing downward through the atmosphere at roughly 2 picoamperes per square meter [23, 24]. The Schumann resonances—electromagnetic standing waves in the Earth-ionosphere cavity at approximately 7.83, 14, 20, 26 Hz and harmonics—are generated by this lightning activity and propagate globally with attenuation of only about 0.5 dB per megameter, enabling detection anywhere on the planet [25].

5.2 Cosmic Ray Forcing of the GEC

The critical link for The Traveling System framework is this: the primary source of ionization in the lower atmosphere from ground level to approximately 60–70 km altitude is galactic cosmic rays [24, 26]. It is GCR ionization that maintains the electrical conductivity of the atmosphere from the surface upward. Changes in GCR flux therefore change atmospheric

conductivity, which changes the GEC current density, which modulates cloud microphysics and the Schumann resonance state [24].

Measurements of the air-earth current at solar minimum—when GCR flux is at maximum due to reduced solar wind shielding—show larger current values than at solar maximum, consistent with Svensmark’s mechanism operating through the GEC [24]. Harrison and Usoskin have identified this signal in historical air-earth current records at Lerwick Observatory in the UK [24]. Even a 1–3% variation in GCR flux in equatorial regions due to solar wind variation may measurably affect thunderstorm charging current and ionospheric potential [26].

5.3 The Chain of Forcing

Assembling the evidence, the chain of forcing from galactic position to terrestrial climate runs as follows: Galactic position determines ISM density and magnetic structure. ISM density and magnetic structure determine heliospheric geometry and the efficiency of the heliosphere as a cosmic ray shield. Heliospheric geometry determines GCR flux at Earth’s atmosphere. GCR flux determines atmospheric ionization. Atmospheric ionization determines the conductivity profile of the atmosphere and the GEC current density. GEC state modulates cloud nucleation, cloud microphysics, and the Schumann resonance frequencies. These changes affect planetary albedo and energy balance.

Every link in this chain has published supporting evidence. What The Traveling System proposes is that this chain extends further upstream than previously modeled—to galactic position rather than only to solar activity. The GEC is not merely a solar-terrestrial coupling mechanism. It is a receiver that responds to the full electromagnetic and particle environment of the heliospheric boundary, which is itself determined by where we are in the galaxy.

The GPDM (Geophysical Plasma Discharge Model) of geological anomaly sites, previously published by this author [27], proposes that telluric current state and Schumann resonance state modulate plasma discharge probability at geologically sensitive sites. If the Schumann resonance baseline is itself modulated by galactic forcing through the chain described above, then geological discharge activity at such sites may contain a record of heliospheric variability extending into the deep past. This represents a testable and previously unproposed connection between local geological anomaly phenomena and galactic environmental forcing.

6. BIOLOGICAL AND CHEMICAL PRESENCE IN THE INTERSTELLAR MEDIUM

The intuition that space is empty is operationally false. The interstellar medium is a chemically active environment. More than 260 distinct molecular species have been detected in interstellar clouds, including alcohols, aldehydes, organic acids, and amides [28]. Interstellar dust grains serve as chemical reaction surfaces, their icy mantles irradiated by GCRs and UV photons to produce complex organic molecules (COMs)—considered key precursors to amino acids, sugars, and nucleobases [29]. The total mass of abiotically produced organic molecules in the ISM is estimated at a minimum of 10 million solar masses [30].

The panspermia hypothesis—that microbial life or its precursors can travel through space—has received substantial experimental support. The Tanpopo mission demonstrated that *Deinococcus* bacteria survived for three years on the exterior of the International Space Station

without rock shielding when aggregated in pellets of sufficient thickness [31, 32]. The same research group detected *Deinococcus* bacteria floating at 12 km altitude using high-altitude balloons, suggesting active microbial presence in near-space [32]. These findings introduce the concept of massapanspermia: interplanetary transfer of microbial aggregates without lithological protection.

During heliospheric collapse events of the type documented 2–3 million years ago, the barrier between the inner solar system and the interstellar medium is removed. Earth is exposed not only to energetic radiation but to the full chemical and potentially biological content of the cloud material. Neutral hydrogen densities of approximately $3,000 \text{ cm}^{-3}$ enter the inner solar system unimpeded. The ozone-depleting chemistry documented in prior modeling [33] occurs in the context of simultaneous ISM chemical influx. The atmospheric and biological implications of this combined exposure remain poorly characterized and represent a significant gap in the literature.

7. ADDITIONAL FORCING VARIABLES

7.1 Geomagnetic Field as Intermediate Variable

Earth’s internal magnetic field is an intermediate variable in the GCR forcing chain that has received insufficient attention in this context. When geomagnetic field intensity drops below approximately 40% of its present value—as occurs during polarity reversals—GCR flux reaching the atmosphere increases by more than 40% [34]. Paleoclimate reconstructions across the Matuyama-Brunhes and Lower Jaramillo geomagnetic reversals show anomalous cooling events that coincide with field intensity minima and recover when the field strengthens [34]. This demonstrates that Earth’s own magnetic shield is a climate-relevant variable on centennial-to-millennial timescales.

More significantly, there is evidence that the geomagnetic field’s reversal behavior may itself be forced by galactic environment. Spiral arms, with their turbulent magnetic fields, appear to correlate with higher geomagnetic reversal rates, while superchrons—extended periods of stable geomagnetic polarity—occur preferentially when the solar system is located between spiral arms in regions of more homogeneous galactic magnetic fields [35]. If confirmed, this would mean galactic position influences not only the external cosmic ray flux but the internal geomagnetic shield, creating a compound forcing effect.

7.2 Milankovitch Cycles and the Forcing Hierarchy

Milankovitch orbital cycles—the approximately 100,000-year eccentricity cycle, the 41,000-year axial tilt cycle, and the 26,000-year precession cycle—are the established primary drivers of ice age timing [36]. The Traveling System framework does not compete with this model. Instead, we propose a nested forcing hierarchy in which Milankovitch cycles operate within a longer-period galactic envelope that sets the baseline climate state into which orbital forcing is applied. The galactic environment determines which orbital configurations are capable of triggering glaciation.

7.3 Solar Long Cycles

Between the 11-year solar cycle and galactic-scale forcing, a series of solar magnetic activity cycles operate on timescales of decades to millennia: the Gleissberg cycle (~88 years), the

de Vries/Suess cycle (~210 years), and the Hallstatt cycle (~2,300 years). These cycles modulate GCR flux at Earth on human-history timescales. The Little Ice Age is associated with the Maunder Minimum in the context of the Hallstatt cycle. These cycles are already partially incorporated into paleoclimate science but are rarely framed as part of the larger galactic forcing hierarchy described here.

7.4 The Local Ribbon of Cold Clouds

The specific interstellar structure that triggered the 2–3 million year ago heliospheric collapse has been identified as the Local Ribbon of Cold Clouds (LRCC) in the constellation Lynx [13]. The LRCC is characterized by very small physical extent, smooth velocity fields, and neutral hydrogen densities that can exceed $3,000 \text{ cm}^{-3}$. Cold dense clouds of this type are rare—occupying less than 1% of the ISM by volume [13]—but their ram pressure is sufficient to collapse the heliosphere entirely when encountered. Understanding the distribution and dynamics of such structures along the solar system’s future trajectory is a key research priority.

7.5 Nearby Supernovae

The Local Bubble itself was formed by an estimated 15 supernovae beginning approximately 14 million years ago [37]. Individual supernovae within approximately 100 parsecs can substantially compress the heliosphere and temporarily elevate hard radiation at Earth’s surface. The ^{30}Fe isotope pulse at 6–7 million years ago is also consistent with supernova ejecta, though Opher et al. [13] argue the cold cloud scenario better fits the combined isotope and heliospheric evidence. Both mechanisms may have operated, and they are not mutually exclusive.

8. THE FORCING HIERARCHY: A UNIFIED FRAMEWORK

The Traveling System proposes a multi-scale forcing hierarchy in which each level operates over distinct timescales and through distinct physical mechanisms, but all converge on the same terrestrial receiver: the global atmospheric electric circuit and its relationship to cloud formation, albedo, and surface temperature.

At the longest timescales (hundreds of millions of years), galactic position relative to spiral arms determines the baseline GCR environment. Spiral arm crossings elevate GCR flux by approximately an order of magnitude and may also perturb geomagnetic reversal rates through turbulent galactic magnetic field interactions.

At intermediate timescales (millions to tens of millions of years), the solar system’s vertical oscillation through the galactic plane, encounters with dense interstellar clouds, and nearby supernova events modulate the heliospheric boundary and produce episodic ISM contact events. The geological record contains isotopic, paleoclimate, and biodiversity signals associated with these episodes.

At shorter timescales (thousands to tens of thousands of years), the solar system’s position relative to specific ISM structures—including the current transition from the LIC toward the G-Cloud—alters the heliospheric boundary conditions and GCR modulation. The Milankovitch cycles operate at this level, with glacial triggering potentially gated by the ISM boundary state.

At solar-activity timescales (decades to millennia), the Gleissberg, de Vries, and Hallstatt solar magnetic cycles modulate the heliospheric cosmic ray shield. The Maunder Minimum and the Little Ice Age represent the historical record of this level of forcing.

At the innermost level, Earth's own geomagnetic field strength provides the final modulating factor before GCR flux reaches the atmosphere. Field intensity variations on centennial-to-millennial timescales, and polarity reversals on longer timescales, create an additional variable layer in the forcing chain.

Anthropogenic CO₂ forcing operates within and on top of all of the above. The total radiative forcing from a 15% reduction in GCR flux over the 20th century due to increased solar magnetic activity, if the GCR-cloud mechanism is correct, is comparable to total anthropogenic forcing [16]. This does not diminish the reality of anthropogenic forcing; it suggests that natural external forcing may be substantially larger than current models account for, and that the two effects may be partially coupled in ways not yet characterized.

9. FALSIFIABLE PREDICTIONS

The Traveling System framework generates several predictions that are testable with existing or planned instrumentation.

Prediction 1: Schumann resonance baseline trend. If GCR flux is changing due to the solar system's approach to the LIC-G-Cloud transition boundary, the Schumann resonance fundamental frequency and amplitude should show a long-term secular trend distinguishable from solar cycle modulation. Long-term Schumann resonance monitoring stations at Nagycenk (Hungary), West Greenwich (Rhode Island), and Vernadsky (Antarctica) have records extending to 1993. Statistical analysis of these records for non-solar secular trends has not been published.

Prediction 2: Heliospheric boundary asymmetry. IBEX and Voyager data should show asymmetries in heliospheric boundary structure consistent with the solar system's direction of motion relative to the approaching G-Cloud. This prediction is partially testable with existing Voyager 1 and 2 interstellar data.

Prediction 3: Cosmogenic isotope record. If heliospheric collapse events produce the ³⁰Be anomalies predicted by Opher et al.'s cold cloud model [13], high-resolution marine sediment cores from the 2–3 million year interval should show elevated ³⁰Be production signatures correlated with paleoclimate cooling. Existing cores may contain this signal but require re-analysis without the assumption of constant ³⁰Be input used in some prior studies.

Prediction 4: Geological discharge correlation. If Schumann resonance state is modulated by GCR flux at geological timescales, anomalous discharge sites with documented telluric current sensitivity—including the Hessdalen Valley, Brown Mountain, and comparable sites—should show discharge activity patterns with a periodicity component correlated with known cosmic ray flux proxies such as ³⁰Be and ¹⁴C records. This prediction connects The Traveling System framework directly to the GPDM framework [27].

Prediction 5: LIC exit signal. Within the next 1,900 to 20,000 years, the solar system will complete its passage through the Local Interstellar Cloud and enter the transition zone toward the G-Cloud. This transition should produce measurable changes in heliospheric boundary conditions

detectable by a dedicated interstellar boundary monitoring spacecraft. We recommend that heliospheric boundary measurement be prioritized as a long-term observational program analogous to continuous solar irradiance monitoring.

10. CONCLUSIONS

The solar system is in motion. It has always been in motion. Over the past 2.5 million years—the entire span of the genus *Homo*—the solar system has traveled through the deep vacuum of the Local Bubble, a region of unusually low interstellar density produced by ancient supernovae. This environment may have been necessary for stable climate conditions conducive to human evolution. Frisch [1] observed as much in the preface to *Solar Journey*: it may not be a coincidence that Earth was in the low-density Local Bubble during the period when the genus *Homo* emerged.

We are now approaching the edge of the Local Interstellar Cloud and the transition to a different interstellar environment. We do not know precisely what that environment contains. We have no dedicated instrument measuring the interstellar medium ahead of the solar system’s trajectory. We are navigating by inference.

This paper has assembled evidence from heliospheric astrophysics, cosmic ray climate science, atmospheric electricity, archaeoastronomy, and astrobiology to argue that the galactic environment of the solar system is a primary forcing variable for terrestrial climate and biosphere that has been systematically omitted from mainstream Earth system models. The evidence is not speculative. The cold cloud collapse 2–3 million years ago is supported by isotope data on four independent geological substrates. The cosmic ray-cloud mechanism is supported by 30 years of observational correlation and CERN laboratory confirmation of the underlying particle physics. The atmospheric electric circuit sensitivity to GCR flux is established in the peer-reviewed literature.

What is missing is synthesis. The *Traveling System* provides that synthesis: a named framework, a multi-scale forcing hierarchy, and a research agenda connecting the outer boundary of the heliosphere to the surface of the Earth.

We are not a fixed planet with a broken atmosphere. We are a solar system in motion through an energetically variable universe, and everything we experience—climate, biosphere, and perhaps awareness itself—may be a function of where we are in that journey.

ACKNOWLEDGMENTS

The author thanks the research community whose published work forms the evidential foundation of this framework, particularly Priscilla C. Frisch, Nir Shaviv, Merav Opher, Abraham Loeb, Jasper Kirkby, and Kenneth Carslaw. The author acknowledges the GPDM research network including Fred Pallesen (Project Hessdalen), Jean Vézina, Anders Warell, and Melina Tsapos at Lund University’s Pufendorf Institute.

REFERENCES

- [1] Frisch, P. C. (Ed.) (2006). *Solar Journey: The Significance of Our Galactic Environment for the Heliosphere and Earth*. Springer, Dordrecht. ISBN 978-1-4020-4397-0.

- [2] Opher, M., Loeb, A., & Peek, J. E. G. (2024). A possible direct exposure of the Earth to the cold dense interstellar medium 2–3 Myr ago. *Nature Astronomy*, 8, 983–990. <https://doi.org/10.1038/s41550-024-02279-8>
- [3] Opher, M., Loeb, A., Zucker, C., Goodman, A., et al. (2024). The passage of the solar system through the edge of the Local Bubble. *The Astrophysical Journal*, 972(2), 201. <https://doi.org/10.3847/1538-4357/ad596e>
- [4] Opher, M., Giacalone, J., Loeb, A., et al. (2026). Increased and varied radiation during the Sun’s encounters with cold clouds in the last 10 million years. *Scientific Reports*. <https://doi.org/10.1038/s41598-026-36926-z>
- [5] NASA StarChild. Does the Sun move around the Milky Way? <https://starchild.gsfc.nasa.gov/docs/StarChild/questions/question18.html>
- [6] Stanford Solar Center. What is the speed of the Solar System? <https://solar-center.stanford.edu/FAQ/Qsolsyspeed.html>
- [7] Shaviv, N. J. (2006). Long-term variations in the galactic environment of the Sun. In P. C. Frisch (Ed.), *Solar Journey* (pp. 99–131). Springer.
- [8] Local Bubble (2026, February). Wikipedia. https://en.wikipedia.org/wiki/Local_Bubble
- [9] Local Interstellar Cloud (2026, February). Wikipedia. https://en.wikipedia.org/wiki/Local_Interstellar_Cloud
- [10] Vannier, H., Redfield, S., Wood, B. E., et al. (2025). Mapping our path through the local interstellar medium. *The Astrophysical Journal*, 981(2), 102. <https://doi.org/10.3847/1538-4357/adb033>
- [11] Wallner, A., et al. (2016). Recent near-Earth supernovae probed by global deposition of interstellar radioactive ^{30}Fe . *Nature*, 532, 69–72.
- [12] Wallner, A., et al. (2021). ^{30}Fe and ^{24}Pu deposited on Earth constrain the r-process yields of recent nearby supernovae. *Science*, 372, 742–745.
- [13] Opher, M., Giacalone, J., Loeb, A., et al. (2026). Increased and varied radiation during the Sun’s encounters with cold clouds in the last 10 million years [Full paper, see ref. 4].
- [14] Zachos, J., et al. (2001). Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science*, 292, 686–693.
- [15] Svensmark, H., & Friis-Christensen, E. (1997). Variation of cosmic ray flux and global cloud coverage —a missing link in solar-climate relationships. *Journal of Atmospheric and Solar-Terrestrial Physics*, 59, 1225–1232.
- [16] Kirkby, J., & Carslaw, K. S. (2006). Variations of galactic cosmic rays and the Earth’s climate. In P. C. Frisch (Ed.), *Solar Journey* (pp. 349–397). Springer.
- [17] CLOUD experiment sharpens climate predictions. (2013). CERN News. <https://home.cern/news/news/experiments/cloud-experiment-sharpens-climate-predictions>
- [18] CERN’s CLOUD experiment provides unprecedented insight into cloud formation. CERN Press Release. <https://home.web.cern.ch/news/press-release/cern/cerns-cloud-experiment-provides-unprecedented-insight-cloud-formation>
- [19] Shaviv, N. J., & Veizer, J. (2003). Celestial driver of Phanerozoic climate? *GSA Today*, 13(7), 4–10.
- [20] The Collector. Maya Calendars: How Did The Maya Count Time? <https://www.thecollector.com/maya-calendar/>
- [21] Smithsonian National Museum of the American Indian. The Calendar System. <https://maya.nmai.si.edu/calendar/calendar-system>
- [22] de Santillana, G., & von Dechend, H. (1969). *Hamlet’s Mill: An Essay on Myth and the Frame of Time*. Gambit, Boston.

- [23] Williams, E. R., et al. (2023). Studying Earth’s double electrical heartbeat. *Eos*. <https://doi.org/10.1029/2023EO230078>
- [24] Siingh, D., et al. (2007). The atmospheric global electric circuit: An overview. *Atmospheric Research*, 84(2), 91–110. <https://doi.org/10.1016/j.atmosres.2006.05.005>
- [25] Nickolaenko, A. P., & Hayakawa, M. (2014). Recent advances and challenges in Schumann resonance observations and research. *Remote Sensing*, 15(14), 3557.
- [26] Williams, E. R. (2014). Global circuit response to the 11-year solar cycle. *Proc. 15th International Conference on Atmospheric Electricity*, Norman OK.
- [27] Carter, J. (2026). Geological Pathway Diversity Model (GPDM): A geophysical plasma discharge framework for anomalous light phenomena. *EarthArxiv*. <https://doi.org/10.31223/X5J47D>
- [28] Puzzarini, C., & Alessandrini, S. (2024). Interstellar ices: A factory of the origin-of-life molecules. *ACS Central Science*, 10(1), 13–15.
- [29] Puzzarini, C., & Alessandrini, S. (2024). [See ref. 28.]
- [30] NASA Technical Reports Server. Chemical and biological evolution in space. NTRS ID 19850054732.
- [31] Yamagishi, A., et al. (2020). DNA damage and survival time course of *Deinococcus* cell pellets during 3 years of exposure to outer space. *Frontiers in Microbiology*, 11, 2050. <https://doi.org/10.3389/fmicb.2020.02050>
- [32] EurekaAlert. Bacteria could survive travel between Earth and Mars when forming aggregates. (2020). <https://www.eurekaalert.org/news-releases/847880>
- [33] Yeghikyan, A., & Fahr, H. (2006). Accretion of interstellar material into the heliosphere and onto Earth. In P. C. Frisch (Ed.), *Solar Journey* (pp. 317–347). Springer.
- [34] Kitaba, I., et al. (2013). Midlatitude cooling caused by geomagnetic field minimum during polarity reversal. *PNAS*, 110(4), 1215–1220. <https://doi.org/10.1073/pnas.1213389110>
- [35] Wendler, J. (2004). External forcing of the geomagnetic field? Implications for the cosmic ray flux–climate variability. *Journal of Atmospheric and Solar-Terrestrial Physics*, 66(13), 1195–1203.
- [36] Hays, J. D., Imbrie, J., & Shackleton, N. J. (1976). Variations in the Earth’s orbit: Pacemaker of the ice ages. *Science*, 194, 1121–1132.
- [37] Zucker, C., Goodman, A., Alves, J., et al. (2022). Star formation near the Sun is driven by expansion of the Local Bubble. *Nature*, 601, 334.