UINTAH · SPICE · gasCombustion

Central-upwind fluxes and a PIMPLE outer loop

Method and implementation notes for the SPICE component

Numerics follow *reactingPimpleCentralFoam* (Kraposhin et al. hybrid KT/PIMPLE solver)

In one paragraph. Detonations computed with ICE + gasCombustion reach the right wave speed and CJ state but show induction zones that are too thin and oscillations right behind the shock. Two causes were targeted. First, ICE builds face states from symmetric, centred averages that carry no upwind dissipation. Second, a single forward-Euler stage is not TVD for limited schemes. This work adds the Kurganov–Tadmor (KT) central-upwind flux construction of reactingPimpleCentralFoam: limited own/nei reconstructions, local wave speeds, α weights, the a_{Sf} dissipation term and the Mach/acoustic-Courant blending function κ. It also adds a PIMPLE-style outer loop that re-runs the whole ICE fluid chain at the latest iterate and combines iterates with a Crank–Nicolson θ rule. KT and the outer loop sit behind input flags that default to off; with them off, SPICE runs ICE's own tasks plus the consistent pressure-work formulation, which SPICE always uses.

Contents

1 Scope and the two switches **3**

2 Where the changes sit in an ICE timestep **3**

3 Limited own/nei face reconstruction **3**

4 Central-upwind weights and the blending function **5**
4.1 Local wave speeds and the α weights 5
4.2 The hybrid blending function κ 5

5 Split mass fluxes **6**

6 Transport: the CentralUpwind advector **6**

7 Face velocity, face pressure and the pressure update **7**
7.1 Face velocity (`faceVelocity`) 7
7.2 Face pressure (`facePressure`) 7
7.3 Pressure update (`ktPressureFlux`) 7

8 The PIMPLE outer loop **8**
8.1 The update rule 8
8.2 How it is realised in Uintah 8
8.3 Why explicit ICE is enough for detonations 9

9 Fidelity map **10**

10 Input reference **11**

11 Suggested checks **11**

12 Implementation lessons **12**

File index **12**

1 Scope and the two switches

The work touches explicit ICE with a single ICE material on a single level, reacting through the `gasCombustion` model. Two independent groups of options control it:

Switch	Effect
`<advection type="CentralUpwind">`	Turns on the KT machinery: weights, fluxes, the new advector, the KT face velocity and face pressure, and the KT term in the pressure update (Sections 3–7).
`<PIMPLE> nOuterCorrectors > 1`	Turns on the outer loop (Section 8). It works with any advection type but was designed to be paired with `CentralUpwind`.

The design decisions agreed before implementation were:

- **The GitHub solver is the specification for the numerics.** Earlier Uintah experiments were consulted only for framework plumbing (sub-scheduler data-warehouse handling, ghost widths).
- **Explicit ICE, not impICE.** pimpleCentralFoam is implicit for all-Mach efficiency, not for detonation stability. At the tutorial's `maxCo 0.1`, $\kappa = 1$ at the front and the scheme behaves like explicit KT (Section 8.3).
- **Energy stays in ICE's internal-energy form** with the existing `newPressWork 1` pressure work and kinetic-energy correction. No total-energy rewrite was made.

2 Where the changes sit in an ICE timestep

An explicit ICE step is a Lagrangian phase (pressure, face velocities, momentum and energy sources) followed by an advection (remap) phase and a conversion back to primitive variables. Figure 1 shows the task sequence with the new tasks marked, and the outer loop wrapped around it.

> **In the code** `SPICE/SPICE.cc`: `SPICE::scheduleTimeAdvance` calls `SPICE::scheduleFluidChain(sched, level, inOuterLoop)`, which holds the task span from the thermo properties through `advectAndAdvanceInTime`. The flag `inOuterLoop` replaces `scheduleComputeModelSources` by `schedulePimpleCopyModelSources`.

3 Limited own/nei face reconstruction

Every KT quantity starts from two reconstructions of a cell-centred field at a face: one from the low side (*own*) and one from the high side (*nei*). This matches OpenFOAM's `fvc::interpolate(q, own|nei, "reconstruct(q)")`. On a structured mesh the face f normal to direction d sits between cell P (low) and cell $N = P + \hat{e}_d$ (high):

$$q_{\text{own}} = q_P + \tfrac{1}{2}\,\psi(r_P)\,(q_N - q_P), \qquad r_P = \frac{q_P - q_{P-1}}{q_N - q_P}, \tag{1}$$

$$q_{\text{nei}} = q_N - \tfrac{1}{2}\,\psi(r_N)\,(q_N - q_P), \qquad r_N = \frac{q_{N+1} - q_N}{q_N - q_P}. \tag{2}$$

With a central cell gradient on a uniform mesh, OpenFOAM's gradient-based ratio $r = 2\,(\mathbf{d}\cdot\nabla q)_P/(q_N - q_P) - 1$ reduces to exactly r_P above, so the two constructions agree. The limiter $\psi(r)$ is zero for $r \le 0$ (a local extremum falls back to first order), and otherwise:




Figure 1: Explicit ICE timestep with the new and modified tasks. The main-graph pass is outer iteration $k = 0$. The model sources are evaluated once, from u^n, in the main graph only.

`limiter=`	$\psi(r)$, $r > 0$	note
`vanAlbada` (default)	$\dfrac{r(r+1)}{r^2+1}$	OpenFOAM definition, used by the reference tutorials
`vanLeer`	$\dfrac{2r}{1+r}$	
`minmod`	$\min(r, 1)$	most dissipative second-order choice
`MC`	$\min\left(2r, \frac{1+r}{2}, 2\right)$	least dissipative
`upwind`	0	first order everywhere

Vector fields are limited component by component. If $|q_N - q_P|$ is at round-off level, then $q_{own} = q_P$ and $q_{nei} = q_N$.

Boundaries. The four-cell stencil needs $P-1$ and $N+1$. Whether they exist is decided from the *level's* cell index range, including extra cells (`Level::findCellIndexRange`) and periodicity, never from the extent of a fetched array. When $P-1$ (or $N+1$) falls outside, that side is taken to first order ($\psi = 0$). The index substituted for the missing cell is P (or N), so no array is ever indexed out of range: C++ evaluates the arguments before the call even when their values go unused.

> **In the code** `SPICE/Core/CentralUpwind.h` (pure math, no Patch or DataWarehouse): `limiter()`, `ownNei()` for scalars and `Vector`, `upwindValue()`, `ktStencilFlags()`.




Figure 2: Face convention. In Uintah the face index equals the index of cell N.

> `SPICE/SPICE.cc`: the `Pm1Safe/Np1Safe` idiom at every call site.

4 Central-upwind weights and the blending function

4.1 Local wave speeds and the α weights

With u_{own}, u_{nei} the reconstructed face-normal velocities and c_{own}, c_{nei} the reconstructed sound speeds, the one-sided local speeds of propagation are

$$a^+ = \max\big(u_{own} + c_{own},\ u_{nei} + c_{nei},\ 0\big), \qquad a^- = \min\big(u_{own} - c_{own},\ u_{nei} - c_{nei},\ 0\big). \tag{3}$$

`fluxScheme`	α_{own}	a_{Sf}
`Kurganov` (default)	$\dfrac{a^+}{a^+ - a^-}$	$a^-\,\alpha_{own} \le 0$
`Tadmor`	$\frac{1}{2}$	$-\frac{1}{2}\max(a^+, -a^-)$

with $\alpha_{nei} = 1 - \alpha_{own}$. With these, the KT numerical flux of a conserved quantity U with physical flux F is

$$\mathcal{F}_f = \alpha_{own} F_{own} + \alpha_{nei} F_{nei} + a_{Sf}\,(U_{nei} - U_{own}). \tag{4}$$

Because $a_{Sf} \le 0$, the last term is a dissipation proportional to the jump between the two reconstructions. In smooth flow the two reconstructions nearly coincide and the term vanishes; at a shock the limiter opens the jump and the term switches on. Supersonic flow gives $a^- = 0$, so $\alpha_{own} = 1$ and $a_{Sf} = 0$: the flux becomes pure upwind, from the own side.

4.2 The hybrid blending function κ

reactingPimpleCentralFoam blends between the compressible KT flux and a pressure-based, upwind-limited flux with

$$\kappa = \min\left(\frac{\mathrm{Ma}_f}{\mathrm{Co}_{ac}},\ 1\right), \qquad \mathrm{Ma}_f = \frac{|\alpha_{own} u_{own} + \alpha_{nei} u_{nei}|}{\alpha_{own} c_{own} + \alpha_{nei} c_{nei}}, \qquad \mathrm{Co}_{ac} = \frac{\max(a^+, -a^-)\,\Delta t}{\Delta x_d} \tag{5}$$

(`kappaFunction MachToAcCourant`, the reference's *standardMachToAcCourantRatio*). The value `kappaFunction one` forces $\kappa = 1$. Near a detonation front $\mathrm{Ma} \sim 1$ and $\mathrm{Co}_{ac} \lesssim 0.5$, so $\kappa = 1$ (full KT). In a low-Mach deflagration $\kappa \ll 1$ and the scheme relaxes toward an upwind-limited flux close to baseline ICE.

> **Departure from the reference** The reference computes κ at the end of an outer iteration from the mass fluxes of that iteration, and uses it in the next. Here κ is computed from the reconstructed cell states at the start of each fluid-chain pass. Inside the outer loop this is equivalent to a lag of one iteration.

> **In the code** Task `SPICE::computeCentralWeights`, scheduled right after the equilibration pressure. Inputs: `rho_CC`, `speedSound_CC` (NewDW) and `vel_CC` (OldDW), all at ghost 2, plus `delT`. Outputs: `ktAlphaOwn_FC{X,Y,Z}`, `ktaSf_FC{X,Y,Z}` and `ktKappa_FC{X,Y,Z}`.

5 Split mass fluxes

Once ICE has produced the final face velocity u_f for the step (`uvel_FCME` after the face momentum exchange), the reference's split fluxes are formed:

$$\tilde{u}_{\text{own}} = \alpha_{\text{own}} u_f - a_{Sf}, \qquad \tilde{u}_{\text{nei}} = \alpha_{\text{nei}} u_f + a_{Sf}, \tag{6}$$

$$\rho^{\kappa}_{\text{own}} = \kappa\rho_{\text{own}} + (1-\kappa)\rho_{up}, \qquad \rho^{\kappa}_{\text{nei}} = \kappa\rho_{\text{nei}} + (1-\kappa)\rho_{up}, \tag{7}$$

$$\phi_{\text{own}} = \tilde{u}_{\text{own}}\,\rho^{\kappa}_{\text{own}}, \qquad \phi_{\text{nei}} = \tilde{u}_{\text{nei}}\,\rho^{\kappa}_{\text{nei}}, \tag{8}$$

where ρ_{up} is the own or nei density chosen by the sign of u_f, and $\tilde{u}_{\text{own}}$, $\tilde{u}_{\text{nei}}$ are the reference's `aphiv_own` and `aphiv_nei`. The κ blend of the densities is the reference's blend of `psi_own/nei` toward a single face value. Last comes the reference's `updateKappa` redistribution:

$$\phi_{\text{own}} \leftarrow \phi_{\text{own}} + (1-\kappa)\,\phi_{\text{nei}}, \qquad \phi_{\text{nei}} \leftarrow \kappa\,\phi_{\text{nei}}. \tag{9}$$

Two identities worth knowing. The redistribution leaves the total mass flux $\phi_{\text{own}} + \phi_{\text{nei}}$ unchanged. Writing $\rho_F = \kappa(\alpha_{\text{own}}\rho_{\text{own}} + \alpha_{\text{nei}}\rho_{\text{nei}}) + (1-\kappa)\rho_{up}$,

$$\phi_{\text{own}} + \phi_{\text{nei}} = \rho_F\,u_f + \kappa\,a_{Sf}\,(\rho_{\text{nei}} - \rho_{\text{own}}), \qquad \text{so} \qquad U_{KT} \equiv \frac{\phi_{\text{own}} + \phi_{\text{nei}}}{\rho_F} = u_f + \kappa\,a_{Sf}\,\frac{\rho_{\text{nei}} - \rho_{\text{own}}}{\rho_F}. \tag{10}$$

So the KT face volume flux is ICE's own face velocity plus an acoustic dissipation that is proportional to the density jump and switched by κ. The limits are:

- $\kappa = 1$: the full KT mass flux, $\alpha_{\text{own}}\rho_{\text{own}}u_f + \alpha_{\text{nei}}\rho_{\text{nei}}u_f + a_{Sf}(\rho_{\text{nei}} - \rho_{\text{own}})$.
- $\kappa = 0$: $\phi_{\text{own}} = \rho_{up}u_f$ and $\phi_{\text{nei}} = 0$, which is TVD upwind transport.

> **In the code** Task `SPICE::computeCentralFluxes`, scheduled after the face momentum exchange and before the pressure update. Outputs: `ktPhiOwn_FC*`, `ktPhiNei_FC*` and `ktUvol_FC*` ($= U_{KT}$).

6 Transport: the CentralUpwind advector

ICE advects per-cell extensive quantities Q_L (mass, momentum, internal energy, the `newPressWork` kinetic-energy carrier K_L, specific-volume × mass, and species mass), and adds the net transfer to Q_L. The new advector implements the reference's `fvmDiv(phi_own, q) + fvmDiv(phi_nei, q)` with vanAlbada-limited interpolation:

$$q = Q_L/m_L, \qquad \mathcal{F}_f = \phi_{\text{own}}\,q^{up}(\phi_{\text{own}}) + \phi_{\text{nei}}\,q^{up}(\phi_{\text{nei}}), \qquad q^{up}(\phi) = \begin{cases} q_{\text{own}} & \phi \geq 0 \\ q_{\text{nei}} & \phi < 0 \end{cases} \tag{11}$$

$$\Delta Q_c = \Delta t \sum_d A_d \left(\mathcal{F}_{f_d^-} - \mathcal{F}_{f_d^+} \right), \qquad Q_c^{adv} = Q_{L,c} + \Delta Q_c. \tag{12}$$

Mass is the case $q \equiv 1$. With $\kappa = 1$, the transfer of ρq reproduces the KT flux (4): the a_{Sf} parts of $\tilde{u}_{\text{own}}, \tilde{u}_{\text{nei}}$ carry $a_{Sf}\big((\rho q)_{\text{nei}} - (\rho q)_{\text{own}}\big)$.

For intensive fields (`vol_frac`, used by the pressure update) the advector uses a volume flux: $\Delta q_c = \frac{\Delta t}{V}\sum_d A_d\big(q_{f^-}u^{\star}_{f^-} - q_{f^+}u^{\star}_{f^+}\big)$, where q_f is the limited upwind value and $u^{\star}$ is whatever face velocity the caller passed in (Section 7).

Bullet-proofing. If the outgoing mass of a cell exceeds its Lagrangian mass, or the outgoing volume exceeds the cell volume, the advector requests a timestep recompute, exactly as the existing advectors do. AMR refluxing is not supported, and the advector throws if it is requested.

> **In the code** `SPICE/Advection/CentralUpwindAdvector.{h,cc}`, created by `SPICE::createAdvector` (handed to ICE through `ICE::setAdvectorFactory`) for type `CentralUpwind` with the `limiter` attribute. `SPICE::advectAndAdvanceInTime` hands it the six ϕ arrays through `setCentralFluxes()`. Face transfers are evaluated on demand per face, never cached in a patch-sized buffer: the face shared with a higher-index neighbour patch is not part of a patch's own face range (see Section 12).

7 Face velocity, face pressure and the pressure update

7.1 Face velocity (`faceVelocity`)

ICE's face velocity is a centred, density-weighted average of the two cell velocities (`term1`), plus a pressure-gradient term (`term2`) and gravity. With the flag on, only the average is replaced, by its KT analogue: the reference's `HbyA` interpolated own/nei and weighted by α,

$$\text{term1} = \alpha_{\text{own}}\, u_{\text{own}} + \alpha_{\text{nei}}\, u_{\text{nei}}. \tag{13}$$

The pressure and gravity terms are unchanged, and the flag-off expression is kept verbatim.

7.2 Face pressure (`facePressure`)

The reference builds the momentum pressure force as $\nabla p \approx \operatorname{div}\big((\alpha_{\text{own}} p_{\text{own}} + \alpha_{\text{nei}} p_{\text{nei}})\,\mathbf{S}_f\big)$. Correspondingly,

$$p_f = \alpha_{\text{own}}\, p_{\text{own}} + \alpha_{\text{nei}}\, p_{\text{nei}}. \tag{14}$$

This is an upwind-biased weighted average *without* a jump term. It is not a Riemann star pressure, which carries an impedance jump $Z_L Z_R (u_L - u_R)/(Z_L + Z_R)$ and adds dissipation a second time on top of the swept-volume advection.

Energy consistency. SPICE always forms the flow work from `press_FC` and cell-centred velocities, built term for term to match the momentum pressure source (the consistent pressure-work formulation; ICE keeps the original `delP_Dilatate` form). The KT p_f therefore feeds momentum and energy consistently, with no further change. The kinetic energy that the KT momentum transport dissipates is returned as heat by advecting total energy $E = I + K_L$ with the same KT advector and recovering $I = E_{adv} - |\mathbf{m}_{adv}|^2/(2\,m_{adv})$.

7.3 Pressure update (`ktPressureFlux`)

In explicit ICE the pressure increment is $\delta p_{\text{dil}} = -\big(\sum_m \Delta t\, \nabla\cdot(\theta_m u_f)\big) \,/\, \sum_m \kappa_m$, computed through the advector with the face velocity supplied by the caller. With the flag on, that velocity is U_{KT} from (10) instead of u_f. The dilatation then contains the extra term $\kappa\, a_{Sf}(\rho_{\text{nei}} - \rho_{\text{own}})/\rho_F$. This is the explicit counterpart of the convective part of the reference's pressure equation, $\operatorname{div}(\phi^d_{\text{own}} p) + \operatorname{div}(\phi^d_{\text{nei}} p)$: the KT mass-flux divergence with its acoustic upwinding.

> **Departure from the reference** The reference pressure equation also contains the implicit Laplacian $-\operatorname{lap}(D_{p,\text{own}}, p) - \operatorname{lap}(D_{p,\text{nei}}, p)$ and runs `nCorrectors` PISO sweeps. Explicit ICE has no global pressure solve. The closest analogue is the pressure term already in ICE's face velocity. See Section 8.3 for why this is acceptable for detonations.

8 The PIMPLE outer loop

8.1 The update rule

Let X be any conserved per-cell quantity: mass, momentum, specific volume $\times$ mass, species mass, or total energy. Write $\mathcal{C}(\cdot)$ for one pass of the fluid chain, which maps a starting state to its advected state. The main graph computes $X_0^{adv} = \mathcal{C}(X^n)$, and its conversion to primitives is iterate $X^{(0)}$. For $k = 1, \dots, N-1$ the chain is run again from the previous iterate, and the result is combined with the Crank–Nicolson rule:

$$\boxed{X^{(k)} = X^n + (1-\theta)\big(X_0^{adv} - X^n\big) + \theta\big(\mathcal{C}(X^{(k-1)}) - X^{(k-1)}\big)} \tag{15}$$

Each bracket is Δt times the fluid-chain tendency, evaluated at X^n and at the latest iterate respectively. Every quantity inside the chain is therefore re-evaluated at the latest iterate: the KT weights and κ, the face velocities, the pressure, the fluxes and the face states. This matches the reference, which calls `updateCentralWeights` and `updateKappa` in every outer iteration. If the iteration converges, the fixed point is the θ-scheme; $\theta = \frac{1}{2}$ is the tutorial's `ddt CrankNicolson 1`.

Two outer iterations are Heun's method. With $N = 2$ and $\theta = \frac{1}{2}$, write $\mathcal{C}(X) = X + \Delta t\, \mathcal{L}(X)$. Then $X^{(0)} = X^n + \Delta t \mathcal{L}(X^n)$ and

$$X^{(1)} = X^n + \tfrac{\Delta t}{2}\, \mathcal{L}(X^n) + \tfrac{\Delta t}{2}\, \mathcal{L}\big(X^{(0)}\big), \tag{16}$$

which is the two-stage strong-stability-preserving Runge–Kutta method. Unlike forward Euler, it keeps the TVD property of the limited spatial operator under a CFL restriction. This is the most direct reason to expect fewer post-shock oscillations. $\theta = 1$ gives a Picard iteration toward backward Euler.

Energy is blended as total energy. Blending momentum and internal energy separately would create or destroy heat through the nonlinearity of $\frac{1}{2}|\mathbf{m}|^2/M$. Instead, each of the four states $s \in \{n,\ adv_0,\ \mathcal{C},\ k-1\}$ is converted to $E_s = I_s + \frac{1}{2}|\mathbf{m}_s|^2/M_s$, E is blended with (15), and the internal energy is recovered as $I^{(k)} = E^{(k)} - \frac{1}{2}|\mathbf{m}^{(k)}|^2/M^{(k)}$. The internal energies at n and $k-1$ are built exactly as the Lagrangian phase builds them: $M\, e_s(T, Y)$ from the model that owns the caloric equation of state (gasCombustion), or Mc_vT otherwise.

Chemistry is evaluated once per step. As in the reference (`reaction->correct()` before `pimple.loop()`), the model sources (chemistry and diffusion) are computed from u^n in the main graph and copied unchanged into every outer iteration. Because both brackets of (15) contain the same frozen source, the source enters once, with weight one.

Residual. After each outer iteration the code reports $r_k = \max_c \max\big(|\rho^{(k)} - \rho^{(k-1)}|/\rho^{(k)},\ |T^{(k)} - T^{(k-1)}|/T^{(k)}\big)$, a global maximum. With `outerTolerance` > 0 the loop stops early once $r_k <$ `outerTolerance`. The default is a fixed count, as in the reference.

8.2 How it is realised in Uintah

1. **Sub-scheduler.** `d_pimpleSched` is created in `problemSetup` with the data-warehouse mapping ParentOld→0, ParentNew→1, Old→2, New→3 (coarse = fine on a single level). Its graph (Figure 1) is compiled once and recompiled whenever the main graph is.
2. **Driver task.** `pimpleOuterLoop` runs in the main graph right after conserved→primitive, *once per processor* (per-processor patch set, as with impICE's `implicitPressureSolve`), because `execute()` of the sub-graph is collective.

3. **Seeding.** The $k = 0$ primitives (`rho_CC`, `vel_CC`, `temp_CC`, `sp_vol_CC`, `press_CC`, `speedSound_CC`, `specific_heat`, species) are transferred from ParentNewDW into a primed sub-DW. Each iteration then rotates the sub-DWs so that the previous iterate becomes the sub-OldDW. `delT`, `timeStep` and `simTime` are put back in every iteration.
4. **Captured advected state.** Right after the main-graph advection, a small task copies `mass_adv`, `mom_adv`, `eng_adv`, `sp_vol_adv` and every species `var_adv` into `*_adv0` labels. `pimpleBlend` reads X_0^{adv} from them, which avoids requiring a label from ParentNewDW while modifying the same label in the sub-NewDW.
5. **Copy-back.** After the last iteration the primitives (including `mach`, `press_CC` and species) are copied into ParentNewDW over *extra cells too*, so the next step starts from the final iterate's boundary values as well as its interior values.
6. **Timestep control.** A recompute or abort request raised inside the sub-graph is forwarded to the main graph.

> **In the code** `SPICE/SPICE.cc` (each with its `schedule...` pair):
> `pimpleOuterLoop`, `capturePimpleAdv0`, `pimpleCopyModelSources`, `pimpleBlend`, `pimpleResidual`.
> `SPICE/Core/SPICELabel.*`: `mass_adv0`, `mom_adv0`, `eng_adv0`, `sp_vol_adv0`, `pimpleResidual`.
> Per-iteration residual print: `SCI_DEBUG="SPICE_PIMPLE:+"`.

8.3 Why explicit ICE is enough for detonations

pimpleCentralFoam is pressure-implicit so that it can take acoustic Courant numbers $\gg 1$ at low Mach, where $\kappa \to 0$ and the scheme becomes pressure-based. In the detonation tutorials, `maxCo 0.1` keeps $\mathrm{Co}_{ac} \approx 0.1$–$0.3$ at the front, so $\kappa = 1$ and the implicit Laplacian is a small correction. A large implicit step would push κ toward zero and smear the shock and the induction zone. The stability that matters here comes from the KT dissipation and the TVD time integration, both of which the explicit version has. An impICE variant would pay off only for runs that also need long, low-Mach phases, such as deflagration-to-detonation transition.

9 Fidelity map

reactingPimpleCentralFoam	**Uintah SPICE (this work)**
`reconstruct(rho, U, p, psi)`, own/nei, vanAlbada	`centralUpwind::ownNei`, default limiter vanAlbada
`ap`, `am`, `alpha_own`, `aSf` (Kurganov / Tadmor)	`computeCentralWeights`
`kappa` (standardMachToAcCourantRatio)	`computeCentralWeights`; κ-blend and redistribution in `computeCentralFluxes`
`aphiv_own/nei`, `phi_own/nei`, `updateKappa`	`computeCentralFluxes`
`fvmDiv(phi_own,·)+fvmDiv(phi_nei,·)` for U, Y, h	`CentralUpwindAdvector` (mass, momentum, $E = I + K_L$, specific volume, species)
`gradp` with α-weighted p_{own}, p_{nei}	`computePressFC` (`facePressure`)
`HbyA` interpolated own/nei with α	`computeVelFace` term1 (`faceVelocity`)
pEqn convective terms $\text{div}(\phi^d p)$	U_{KT} in the explicit pressure update (`ktPressureFlux`)
pEqn Laplacian, `nCorrectors`	not implemented (would need impICE)
`reaction->correct()` once per step	model sources once per step, frozen in the outer iterations
`pimple.loop()`, `nOuterCorrectors`	`pimpleOuterLoop` sub-scheduler
`ddt CrankNicolson 1`	θ rule (15), $\theta = 0.5$
`hEqn` (sensible enthalpy, KChange, KSource)	ICE internal energy with SPICE's consistent pressure work; total-energy blend in the loop
`maxCo 0.1`	`<cfl>` in the input (0.1 in the detonation input)

Departure from the references, collected

- κ is lagged by one outer iteration relative to the reference (Section 4).
- Transport inside one outer iteration is explicit, where OpenFOAM assembles implicit `fvm` matrices. The outer iterations supply the implicit coupling.
- There is no implicit pressure Laplacian and no PISO `nCorrectors`.
- Diffusion is frozen with chemistry across outer iterations. The reference re-solves diffusion inside `YEqn`/`hEqn`.
- Energy is carried as internal energy, not sensible enthalpy.

10 Input reference

Tag	Default	Meaning
`<CFD><SPICE><advection type="CentralUpwind" limiter="...">`		
`limiter`	`vanAlbada`	vanAlbada, vanLeer, minmod, MC, upwind
`<CFD><SPICE><CentralUpwind>` (optional; requires the advection type above)		
`fluxScheme`	`Kurganov`	or `Tadmor`
`kappaFunction`	`MachToAcCourant`	or `one` ($\kappa \equiv 1$)
`faceVelocity`	`true`	KT term1 in the face velocity
`facePressure`	`true`	α-weighted face pressure
`ktPressureFlux`	`true`	U_{KT} in the pressure update
`<CFD><SPICE><PIMPLE>` (optional)		
`nOuterCorrectors`	`1`	total passes per step; 1 = loop off; the reference uses 2
`theta`	`0.5`	θ in (15); $0 < \theta \leq 1$
`outerTolerance`	`0`	0 = fixed count; otherwise early exit on r_k
`nCorrectors`	`1`	parsed, reserved for an impICE variant

Enforced at start-up. SPICE runs on a single level (no AMR) with exactly one ICE material and no MPM materials: like reactingPimpleCentralFoam it is a single-fluid code, with species carried as transported scalars. The outer loop requires explicit ICE and no turbulence model. CentralUpwind requires explicit ICE.

Provided inputs. In `StandAlone/inputs/SPICE/`:

- `riemann_sm.ups, riemann_sm_KT.ups, riemann_sm_PIMPLE.ups, riemann_sm_KT_PIMPLE.ups`: the Sod shock tube of `inputs/ICE/riemann_sm.ups` with each combination of KT advection and the outer loop ($N = 2$, $\theta = 0.5$).
- `riemann_gc.ups`: the same shock tube with gasCombustion supplying the thermodynamic data (`idealAir.xml`, no chemistry), KT and PIMPLE. It exercises the outer loop's transported-scalar and model-source paths.

11 Suggested checks

1. **Oscillations.** Compare `nOuterCorrectors` 1 and 2 with everything else fixed. This isolates the effect of the time integration from that of the flux.
2. **Induction zone.** If it shortens or collapses, first set `facePressure false`. A dissipative face pressure applied as a cell-centred source is the known way to over-dissipate ICE's momentum equation. Then compare `ktPressureFlux true/false`.
3. **Blending.** Compare `kappaFunction one` with `MachToAcCourant`. The detonation front should already sit at $\kappa = 1$; inspect `ktKappa_FCX`.
4. **Limiter.** Compare minmod, vanAlbada and MC to bracket limiter sensitivity.
5. **Time rule.** Compare $\theta = 0.5$ with $\theta = 1$, and use `nOuterCorrectors 3` to check the convergence of the residual r_k with `SCI_DEBUG="ICE_PIMPLE:+"`.
6. **Regression.** Remove all new tags. The original code paths should reproduce the pre-change results.

12 Implementation lessons

Five defects surfaced during testing. Each is a general Uintah pitfall and is recorded here so it is not rediscovered.

Out-of-range stencil arguments

Passing `q[Pm1]` to a function that ignores it when the stencil is invalid still reads that memory, and at a domain edge with one extra cell this segfaulted. The fix is to substitute P or N before indexing (`Pm1Safe`/`Np1Safe`).

Patch-sized face buffers

`Patch::getSFC*HighIndex()` includes the upper face only when the patch has no neighbour there. A face buffer filled over a patch's own face range therefore lacks the face $c+1$ of the patch's last cell at an interior patch boundary. Reading it gave negative specific volume at every patch seam of the 12-rank flame run. Face transfers are now evaluated on demand from ghosted data.

Sub-scheduler driver scheduled per patch

A task that calls a sub-scheduler's collective `execute()` must be added on the per-processor patch set. On `level->eachPatch()` it ran once per patch, each time with only one patch seeded. This failed as soon as a rank owned more than one patch (4 ranks, 12 patches), and went unnoticed with one patch per rank.

Copy-back over interior cells only

The outer loop's final state has to be copied over extra cells too. Otherwise the next step's boundary values come from iteration $k = 0$, which matters for LODI and Neumann boundaries.

Limiter overflow (found in 2D)

The textbook vanAlbada and vanLeer forms, $r(r+1)/(r^2+1)$ and $2r/(1+r)$, evaluate $\infty/\infty = \mathrm{NaN}$ when r is huge. That happens when the jump across a face is tiny next to a finite upwind jump, which is routine for trace species on an oblique 2D front. A 2D channel detonation crashed with NaN temperatures after four steps. For $r > 1$ the code now uses the equivalent forms $(1+1/r)/(1+1/r^2)$ and $2/(1+1/r)$.

File index

`ICE/Core/CentralUpwind.h`	limiters, own/nei reconstruction, KT weights, κ (new)
`ICE/Advection/CentralUpwindAdvector.{h,cc}`	KT transport (new)
`ICE/Advection/Advector.h`	basket extended with the ϕ arrays
`ICE/Advection/AdvectionFactory.cc`, `sub.mk`	registration, `limiter` attribute
`ICE/ICE.{h,cc}`	flags, `scheduleFluidChain`, KT tasks, face velocity and pressure, pressure update, outer loop
`ICE/Core/ICELabel.{h,cc}`	`kt*_FC*`, `*_adv0`, `pimpleResidual` labels
`StandAlone/inputs/UPS_SPEC/ice_spec.xml`	new tags
`ICE/PIMPLE_KT.md`	short usage note