

Local Anesthetics DEFINITION

- Drugs which
 - produce a REVERSIBLE loss of sensation ...
 - in a localized part of the body.....
 - when applied directly onto nerve tissues or mucous membranes
- Local anesthetics are 'local' ONLY because of how they are administered! (*Selectivity*)

The first clinically used Local Anesthetic

Cocaine (ISA activity)

A natural alkaloid from Erythroxylum coca.

Prototype Drug ___ Lignocaine (Synthetic)

Properties Desirable in a Local Anesthetic

- Non-irritating
- Do not cause permanent damage to nerve structure
- Systemic toxicity should be low
- Effective through-Injectedor Applied locally
- Onset of action as short as possible
- DOA long enough to allow time for counter plated surgery

1. CLASSIFICATION ACCORDING TO CHEMICAL CHARACTERISTIC

• ESTERS

- > Cocaine
- > Procaine
- > Tetracaine
- > Benzocaine

• AMIDES

- > Lignocaine/Lidocaine
- Bupivacaine
- >Levobupivacaine
- > Mepivacaine
- > Prilocaine
- > Etidocaine
- > Ropivacaine

2. According to Duration of action

Short Duration of Action

Procaine

Medium Duration of Action

Cocaine, Lidocaine, Mepivacaine, Prilocaine

Long Duration of Action

Tetracaine, Bupivacaine, Etidocaine, Ropivacaine

CLASSIFICATION ACCORDING TO CLINCIAL USES

- SURFACE ANESTHESIA
- > Tetracaine
- > Cocaine
- > Benzocaine
- > Lignocaine
- INFILTRATION ANESTHESIA & FIELD BLOCK ANESTHESIA
 - > Procaine
 - > Lignocaine
 - > Bupivacaine

• NERVE BLOCK ANESTHESIA

- > Procaine
- > Tetracaine
- > Lignocaine
- Bupivacaine
- > Ropivacaine

*SPINAL ANESTHESIA

- > Tetracaine
- > Lignocaine
- Bupivacaine

* EPIDURAL ANESTHESIA

- > Lignocaine
- > Bupivacaine

ANESTHETIC USED IN OPHTHALMOLOGY


> Proparacaine



Most local anesthetics consist of 3 parts

- Lipophilic Aromatic group
- Intermediate chain
- Hydrophilic Amino group

LAs - Weak Bases (pKa:7.5-9)




LIPOPHILIC

HYDROPHILIC

Two types of linkages

give rise to 2 chemical classes of local anesthetics.

ESTER LINKAGE



C₁₃H₁₉CIN₂O₂ • HCI

PROCAINE


procaine (Novocaine)

tetracaine (Pontocaine)

benzocaine

cocaine

AMIDE LINKAGE



C₁₄H₂₂N₂O • HCl

LIDOCAINE

lidocaine (Xylocaine)

mepivacaine (Carbocaine)


bupivacaine (Marcaine)


etidocaine (Duranest)

ropivacaine (Naropin)



- Diffusion into the nerve fiber
- ✓ Blockade of sodium channels





Progressively increasing conc. of a LA applied to a nerve fiber produce blockade of more & more Na+ channels:

- The threshold for excitation increases
- Impulse conduction slows
- The rate of rise of AP declines
- The AP amplitude decreases
- Finally the ability to generate an AP is abolished

SUSCEPTIBILITY OF NERVE FIBER TO LAS DEPEND ON

- Potency
- Size of nerve fiber (small fibers blocked 1st)
- Effect of fiber diameter
- Rate of firing (rapidly firing fibers blocked 1st)
- Effect of fiber position in the nerve bundle (outer fibers blocked 1st, then core fibers)

ORDER OF BLOCKADE

- AUTONOMIC
- PAIN
- TEMPERATURE
- TOUCH
- DEEP PRESSURE
- MOTOR

Recovery in reverse order

PHARMACOKINETICS

Absorption

Dosage

Site of injection

(when used for major conduction blocks, the peak serum levels will vary as a function of the specific site of injection, with intercostal blocks among the highest, & sciatic & femoral among the lowest)

Lipid solubility (more lipid soluble – longer DOA)

PHARMACOKINETICS

Ph

Vascularity (highly vascular area – more rapid absorption – higher blood levels)

- Combination with vasoconstrictors (resultant reduction in blood flow reduces rate of systemic absorption & diminishes peak serum levels)
- Distribution
- Biotransformation & Excretion

Comparison of LA characteristics

	Relative lipid solubility	Relative potency	onset	рКа	Local duration	vasodilation	Plasm a protei n bindin g
procaine	1	1	slow	8.9	short	+++	5%
lidocaine	4	4	rapid	7.9	modera te	+++	55%
tetracain e	80	16	slow	8.5	long	+	75%
bupivaca ine	130	16	slow	8.1	long	+	90%

Plasma protein binding may be used as an indirect measure of tissue binding tendencies

ADVERSE EFFECTS

- CNS (1st stimulation, then depression)
- Local Neurotoxicity (cauda equina syndrome associated with continuous spinal anesthesia CSA)
- CVS (bupivacaine most cardiotoxic)
- ANS
- Motor Paralysis
- Hematological Effects
- Hypersensitivity reactions

Prevention of Toxicity

- Enquire about history of allergy.
- Caution in presence of liver/myocardial damage.
- Proper site (correct knowledge of nerve course).
- Minimal effective dose usage (avoid I/V adm).
- Wait after injection.
- Observe the face for any twitching, excitement, and pulse for tachycardia.
- Observe post op for allergic reactions.
- Avoid food intake at least 04 hrs prior to anesthesia to prevent vomiting.

SELECTIVE PHARMACOLOGICAL PROPERTIES OF SOME ESTER - type LA

Cocaine

- · Medical use limited to surface or topical anesthesia
- Avoid epinephrine because cocaine already has vasoconstrictor properties. (EXCEPTION!!!)
- A toxic action on heart may induce rapid and lethal cardiac failure.
- · A marked pyrexia is associated with cocaine overdose.

SELECTIVE PHARMACOLOGICAL PROPERTIES OF SOME ESTER - type LA

Benzocaine

- pKa ~ 3,
- Available in many preps for relief of pain and irritation
- for surface anesthesia (topical) only ... ointments, sprays, etc.
- · Used to produce anesthesia of mucous membranes
- Methemoglobinemia (Rare case side effect)

SELECTIVE PHARMACOLOGICAL PROPERTIES OF SOME AMIDE - type LA

- LIDOCAINE (Xylocaine) Most widely used LA
 - Effective by all routes.
 - Faster onset, more intense, longer lasting, than procaine.
 - Good alternative for those allergic to ester type
 - More potent than procaine but about equal toxicity
 - More sedative than others

SELECTIVE PHARMACOLOGICAL PROPERTIES OF SOME AMIDE - type LA

- Bupivacaine (Marcaine)
 - No topical effectiveness
 - Slower onset and one of the longer duration agents
 - Unique property of sensory and motor dissociation can provide sensory analgesia with minimal motor block
 - · has been popular drug for analgesia during labor
 - More cardiotoxic than other LA

SELECTIVE PHARMACOLOGICAL PROPERTIES OF SOME AMIDE - type LA

- Ropivacaine
 - Enantiomer of bupivacaine (5 stereoisomer)
 - No topical effectiveness
 - · Clinically ~ equivalent to bupivacaine
 - Similar sensory versus motor selectivity as bupivacaine with significantly less CV toxicity

CLINICAL APPLICATIONS

- SURFACE ANESTHESIA (Topical)
 - Ear, Nose, mouth, bronchial tree, nasopharynx, cornea, GIT and urinary tracts
 - Lidocaine, tetracaine, Benzocaine
 - EMLA cream (Eutectic Mixture of Local Anesthetics) lidocaine 2.5% + prilocaine 2.5% permits anesthetic penetration of keratinized layer of skin as deep as 5mm, producing localized numbness.

INFILTRATION ANESTHESIA

- Direct injection into tissues to reach nerve branches and terminals.
- Can be superficial as well as deep.
- Used in minor surgery.
- Immediate onset with variable duration.
- This type involve skin region as deep as intra abdominal tissue.

.Most LA's used

- NERVE BLOCK or FIELD BLOCK
 - Interruption of nerve conduction upon injection into the region of nerve plexus or trunk.
 - Used for surgery, dentistry, analgesia.
 - Less anesthetic needed than for infiltration
 - Given within specific nerve area such as brachial plexus, within intercostal nerves, abdominal nerves are targeted, cervical plexus when neck region is targeted.

.Most LA's used

SPINAL ANESTHESIA

- Injection into subarachnoid space below level of L2 vertebra to produce effect in spinal roots and spinal cord.
- Use hyperbaric or hypobaric solutions depending on area of blockade.
- Used for surgery to abdomen, pelvis or leg when can't use general anesthesia.
- Can be employed in pts of hepatic, renal & CVS diseases
 - Lidocaine, tetracaine

EPIDURAL AND CAUDAL ANESTHESIA

• Injection into epidural space usually at lumbar or sacral levels or near dura matter where nearly most nerves pass closely. Areas supplied by these nerves are targeted e.g.

.ligamentum flavum(post)

.spinal periosteum(laterally), dura(ant).

- Lower part of the body. Pelvic region
- For painless child birth.

- Unwanted effects similar to that of spinal (pain, hematoma, introduction of foreign particle, hypotension – Rx: raise foot-end of bed or give sympathomimetics, headache – Rx: small bore needle & blood patch, cauda equina syndrome, rarely respiratory paralysis)
 - Lidocaine, bupivacaine, ropivacaine

