



Patient Name : Mr.ASHUTOSH BAGGA

Age/Gender :31 Y 1 M 9 D /M

UHID/MR No : APJ1.0026161583 : DPNUOPV4770

Visit ID

Ref Doctor : Dr.SELF IP/OP NO

Collected : 05/Dec/2023 07:55AM Received : 06/Dec/2023 10:11AM Reported : 06/Dec/2023 02:23PM

Status : Final Report

Client Name : PUP 24X7\_CREDIT

Patient location : Panathur, Bangalore

## **DEPARTMENT OF SEROLOGY**

## SARS-COV-2 IGG ANTIBODY TEST

Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

Anti-SARS-CoV-2 lgG , SERUM	8.91	S/C	<1	ECLIA
-----------------------------	------	-----	----	-------

## **Comment:**

Anti-SARS-CoV-2 IgG Result(S/C)	Interpretation		
<1	Non-reactive for anti-SARS-CoV-2 IgG		
≥1	Reactive for anti-SARS-CoV-2 IgG		

- It is strictly advised to use IgG based assays only for conduct of serosurveys and not for the diagnosis of SARS-CoV-2
- Anti-SARS-CoV-2 IgG test is intended for use as an aid in identifying individuals with an adaptive immune response to SARS-CoV-2, indicating recent or prior infection.
- Anti-SARS-CoV-2 IgG test should not be used to diagnose acute SARS-CoV-2 infection. Negative results do no preclude acute SARS-CoV-2 infection. If acute infection is suspected, direct testing for SARS-CoV-2 is necessary.
- Heterophilic antibodies in serum samples may cause interference in immunoassays. These antibodies may be present in blood samples from individuals regularly exposed to animals or who have been treated with animal serum products.
- Results that are inconsistent with clinical observations indicate the need for additional testing.
- A non-reactive result can occur if the quantity of antibodies for the SARS-CoV-2 virus present in the specimen is below the detection limit of the assay, or the virus has undergone minor amino acid mutation(s) in the epitope recognized by the antibody detected by the test.
- The results obtained with this test should only be interpreted in conjunction with clinical findings, and the results from other laboratory tests and evaluations.
- This test should not be used for screening of donated blood for the purpose of preventing COVID-19 transmission.

\*\*\* End Of Report \*\*\*

DR.MIR SALMAN ALI M.B.B.S,MD Consultant Microbiologist

Page 1 of 1

