

# Understanding Analysis Solutions

Ulisse Mini



# Preface

Huge thanks to the [math discord](#) for answering my questions! I don't know how I'd manage without them ♡

If you don't find your exercise here check [linearalgebras.com](#) or (god forbid) [chegg](#).

You can see my progress [here](#).



# Contents

<b>1</b>	<b>The Real Numbers</b>	<b>1</b>
1.2	Some Preliminaries . . . . .	1
1.3	The Axiom of Completeness . . . . .	8
1.4	Consequences of Completeness . . . . .	13
1.5	Cardinality . . . . .	16
1.6	Cantor's theorem . . . . .	23
<b>2</b>	<b>Sequences and Series</b>	<b>27</b>
2.2	The Limit of a Sequence . . . . .	27
2.3	The Algebraic and Order Limit Theorems . . . . .	31
2.4	The Monotone Convergence Theorem and a First Look at Infinite Series . . . . .	38
2.5	Subsequences and the Bolzano–Weierstrass Theorem . . . . .	45
2.6	The Cauchy Criterion . . . . .	49
2.7	Properties of Infinite Series . . . . .	52
<b>3</b>	<b>Basic Topology of <math>\mathbf{R}</math></b>	<b>61</b>
3.2	Open and Closed Sets . . . . .	61
3.3	Compact Sets . . . . .	67
3.4	Perfect Sets and Connected Sets . . . . .	72
<b>4</b>	<b>Functional Limits</b>	<b>75</b>
4.2	Functional Limits . . . . .	75



# Chapter 1

## The Real Numbers

### 1.2 Some Preliminaries

#### Exercise 1.2.1

- (a) Prove that  $\sqrt{3}$  is irrational. Does a similar argument work to show  $\sqrt{6}$  is irrational?
- (b) Where does the proof break down if we try to prove  $\sqrt{4}$  is irrational?

#### Solution

- (a) Suppose for contradiction that  $p/q$  is a fraction in lowest terms, and that  $(p/q)^2 = 3$ . Then  $p^2 = 3q^2$  implying  $p$  is a multiple of 3 since 3 is not a perfect square. Therefore we can write  $p$  as  $3r$  for some  $r$ , substituting we get  $(3r)^2 = 3q^2$  and  $3r^2 = q^2$  implying  $q$  is also a multiple of 3 contradicting the assumption that  $p/q$  is in lowest terms. For  $\sqrt{6}$  the same argument applies, since 6 is not a perfect square.
- (b) 4 is a perfect square, meaning  $p^2 = 4q^2$  does not imply that  $p$  is a multiple of four as  $p$  could be 2.

#### Exercise 1.2.2

Show that there is no rational number satisfying  $2^r = 3$

#### Solution

Letting  $r = p/q$  we have  $2^{p/q} = 3$  implying  $2^p = 3^q$  which is impossible since 2 and 3 are coprime.

#### Exercise 1.2.3

Decide which of the following represent true statements about the nature of sets. For any that are false, provide a specific example where the statement in question does not hold.

- (a) If  $A_1 \supseteq A_2 \supseteq A_3 \supseteq A_4 \cdots$  are all sets containing an infinite number of elements, then the intersection  $\bigcap_{n=1}^{\infty} A_n$  is infinite as well.
- (b) If  $A_1 \supseteq A_2 \supseteq A_3 \supseteq A_4 \cdots$  are all finite, nonempty sets of real numbers, then the intersection  $\bigcap_{n=1}^{\infty} A_n$  is finite and nonempty.

- (c)  $A \cap (B \cup C) = (A \cap B) \cup C$ .
- (d)  $A \cap (B \cap C) = (A \cap B) \cap C$ .
- (e)  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

**Solution**

- (a) False, consider  $A_1 = \{1, 2, \dots\}$ ,  $A_2 = \{2, 3, \dots\}$ , ... has  $\bigcap_{n=1}^{\infty} A_n = \emptyset$ .
- (b) True.
- (c) False,  $A = \emptyset$  gives  $\emptyset = C$ .
- (d) True, intersection is associative.
- (e) True, draw a diagram.

**Exercise 1.2.4**

Produce an infinite collection of sets  $A_1, A_2, A_3, \dots$  with the property that every  $A_i$  has an infinite number of elements,  $A_i \cap A_j = \emptyset$  for all  $i \neq j$ , and  $\bigcup_{i=1}^{\infty} A_i = \mathbf{N}$

**Solution**

This question is asking us to partition  $\mathbf{N}$  into an infinite collection of sets. This is equivalent to asking us to unroll  $\mathbf{N}$  into a square, which we can do along the diagonal

1	3	6	10	15	...
2	5	9	14	...	
4	8	13	...		
7	12	...			
11	...				
⋮					

**Exercise 1.2.5 (De Morgan's Laws)**

Let  $A$  and  $B$  be subsets of  $\mathbf{R}$ .

- (a) If  $x \in (A \cap B)^c$ , explain why  $x \in A^c \cup B^c$ . This shows that  $(A \cap B)^c \subseteq A^c \cup B^c$
- (b) Prove the reverse inclusion  $(A \cap B)^c \supseteq A^c \cup B^c$ , and conclude that  $(A \cap B)^c = A^c \cup B^c$
- (c) Show  $(A \cup B)^c = A^c \cap B^c$  by demonstrating inclusion both ways.

**Solution**

- (a) If  $x \in (A \cap B)^c$  then  $x \notin A \cap B$  so  $x \notin A$  or  $x \notin B$  implying  $x \in A^c$  or  $x \in B^c$  which is the same as  $x \in A^c \cup B^c$ .
- (b) Let  $x \in A^c \cup B^c$  implying  $x \in A^c$  or  $x \in B^c$  meaning  $x \notin A$  or  $x \notin B$  implying  $x \notin A \cap B$  which is the same as  $x \in (A \cap B)^c$ .



- (c) First let  $x \in (A \cup B)^c$  implying  $x \notin A \cup B$  meaning  $x \notin A$  and  $x \notin B$  which is the same as  $x \in A^c$  and  $x \in B^c$  which is just  $x \in A^c \cap B^c$ . Second let  $x \in A^c \cap B^c$  implying  $x \in A^c$  and  $x \in B^c$  implying  $x \notin A$  and  $x \notin B$  meaning  $x \notin A \cup B$  which is just  $x \in (A \cup B)^c$ .

**Exercise 1.2.6**

- (a) Verify the triangle inequality in the special case where  $a$  and  $b$  have the same sign.
- (b) Find an efficient proof for all the cases at once by first demonstrating  $(a + b)^2 \leq (|a| + |b|)^2$
- (c) Prove  $|a - b| \leq |a - c| + |c - d| + |d - b|$  for all  $a, b, c$ , and  $d$ .
- (d) Prove  $||a| - |b|| \leq |a - b|$ . (The unremarkable identity  $a = a - b + b$  may be useful.)

**Solution**

- (a) We have equality  $|a + b| = |a| + |b|$  meaning  $|a + b| \leq |a| + |b|$  also holds.
- (b)  $(a + b)^2 \leq (|a| + |b|)^2$  reduces to  $2ab \leq 2|a||b|$  which is obviously true. and since squaring preserves inequality this implies  $|a + b| \leq |a| + |b|$ .
- (c) I would like to do this using the triangle inequality, I notice that  $(a - c) + (c - d) + (d - b) = a - b$ . Meaning I can use the triangle inequality for multiple terms

$$|a - b| = |(a - c) + (c - d) + (d - b)| \leq |a - c| + |c - d| + |d - b|$$

The general triangle inequality is proved by repeated application of the two variable inequality

$$|(a + b) + c| \leq |a + b| + |c| \leq |a| + |b| + |c|$$

- (d) I would like to cancel the subtraction inside  $||a| - |b||$  since then the inside will be positive, and the outer absolute value will vanish. Using the suggestion let  $a = (a - b) + b$

$$||a| - |b|| = ||(a - b) + b| - |b|| \stackrel{!}{\leq} ||a - b| + |b| - |b|| = |a - b|$$

However this is incorrect by itself, since  $|a| \leq |c|$  does not imply  $||a| - |b|| \leq ||c| - |b||$  (draw a picture, or use the counterexample  $a = 0, c = 1, b = 2$ ).

We can salvage this argument though, notice if  $|a| \geq |b|$  then  $|a| \leq |c|$  does imply  $||a| - |b|| \leq ||c| - |b||$ . And since we can swap  $a$  and  $b$  without changing anything, we can say without loss of generality assume  $|a| \geq |b|$  and then apply the previous argument.

**Exercise 1.2.7**

Given a function  $f$  and a subset  $A$  of its domain, let  $f(A)$  represent the range of  $f$  over the set  $A$ ; that is,  $f(A) = \{f(x) : x \in A\}$ .

- (a) Let  $f(x) = x^2$ . If  $A = [0, 2]$  (the closed interval  $\{x \in \mathbf{R} : 0 \leq x \leq 2\}$ ) and  $B = [1, 4]$ , find  $f(A)$  and  $f(B)$ . Does  $f(A \cap B) = f(A) \cap f(B)$  in this case? Does  $f(A \cup B) = f(A) \cup f(B)$ ?

- (b) Find two sets  $A$  and  $B$  for which  $f(A \cap B) \neq f(A) \cap f(B)$ .
- (c) Show that, for an arbitrary function  $g : \mathbf{R} \rightarrow \mathbf{R}$ , it is always true that  $g(A \cap B) \subseteq g(A) \cap g(B)$  for all sets  $A, B \subseteq \mathbf{R}$
- (d) Form and prove a conjecture about the relationship between  $g(A \cup B)$  and  $g(A) \cup g(B)$  for an arbitrary function  $g$

**Solution**

- (a)  $f(A) = [0, 4]$ ,  $f(B) = [1, 16]$ ,  $f(A \cap B) = [1, 4] = f(A) \cap f(B)$  and  $f(A \cup B) = [0, 16] = f(A) \cup f(B)$
- (b)  $A = \{-1\}$ ,  $B = \{1\}$  thus  $f(A \cap B) = \emptyset$  but  $f(A) \cap f(B) = \{1\}$
- (c) Suppose  $y \in g(A \cap B)$ , then  $\exists x \in A \cap B$  such that  $g(x) = y$ . But if  $x \in A \cap B$  then  $x \in A$  and  $x \in B$ , meaning  $y \in g(A)$  and  $y \in g(B)$  implying  $y \in g(A) \cap g(B)$  and thus  $g(A \cap B) \subseteq g(A) \cap g(B)$ .

Notice why it is possible to have  $x \in g(A) \cap g(B)$  but  $x \notin g(A \cap B)$ , this happens when something in  $A \setminus B$  and something in  $B \setminus A$  map to the same thing. If  $g$  is 1-1 this does not happen.

- (d) I conjecture that  $g(A \cup B) = g(A) \cup g(B)$ . To prove this we show inclusion both ways, First suppose  $y \in g(A \cup B)$ . then either  $y \in g(A)$  or  $y \in g(B)$ , implying  $y \in g(A) \cup g(B)$ . Now suppose  $y \in g(A) \cup g(B)$  meaning either  $y \in g(A)$  or  $y \in g(B)$  which is the same as  $y \in g(A \cup B)$  as above.

**Exercise 1.2.8**

Here are two important definitions related to a function  $f : A \rightarrow B$ . The function  $f$  is *one-to-one* (1 – 1) if  $a_1 \neq a_2$  in  $A$  implies that  $f(a_1) \neq f(a_2)$  in  $B$ . The function  $f$  is *onto* if, given any  $b \in B$ , it is possible to find an element  $a \in A$  for which  $f(a) = b$ . Give an example of each or state that the request is impossible:

- (a)  $f : \mathbf{N} \rightarrow \mathbf{N}$  that is 1 – 1 but not onto.
- (b)  $f : \mathbf{N} \rightarrow \mathbf{N}$  that is onto but not 1 – 1.
- (c)  $f : \mathbf{N} \rightarrow \mathbf{Z}$  that is 1 – 1 and onto.

**Solution**

- (a) Let  $f(n) = n + 1$  does not have a solution to  $f(a) = 1$
- (b) Let  $f(1) = 1$  and  $f(n) = n - 1$  for  $n > 1$
- (c) Let  $f(n) = n/2$  for even  $n$ , and  $f(n) = -(n + 1)/2$  for odd  $n$ .

**Exercise 1.2.9**

Given a function  $f : D \rightarrow \mathbf{R}$  and a subset  $B \subseteq \mathbf{R}$ , let  $f^{-1}(B)$  be the set of all points from the domain  $D$  that get mapped into  $B$ ; that is,  $f^{-1}(B) = \{x \in D : f(x) \in B\}$ . This set is called the *preimage* of  $B$ .

- (a) Let  $f(x) = x^2$ . If  $A$  is the closed interval  $[0, 4]$  and  $B$  is the closed interval  $[-1, 1]$ , find  $f^{-1}(A)$  and  $f^{-1}(B)$ . Does  $f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$  in this case? Does  $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$ ?
- (b) The good behavior of preimages demonstrated in (a) is completely general. Show that for an arbitrary function  $g : \mathbf{R} \rightarrow \mathbf{R}$ , it is always true that  $g^{-1}(A \cap B) = g^{-1}(A) \cap g^{-1}(B)$  and  $g^{-1}(A \cup B) = g^{-1}(A) \cup g^{-1}(B)$  for all sets  $A, B \subseteq \mathbf{R}$ .

**Solution**

- (a)  $f^{-1}(A) = [-2, 2]$ ,  $f^{-1}(B) = [-1, 1]$ ,  $f^{-1}(A \cap B) = [-1, 1] = f^{-1}(A) \cap f^{-1}(B)$ ,  $f^{-1}(A \cup B) = [-2, 2] = f^{-1}(A) \cup f^{-1}(B)$
- (b) First let  $x \in g^{-1}(A \cap B)$  meaning  $g(x) \in A \cap B$  implying  $g(x) \in A$  and  $g(x) \in B$  which is the same as  $x \in g^{-1}(A)$  and  $x \in g^{-1}(B)$  meaning  $x \in g^{-1}(A) \cap g^{-1}(B)$ .
- Second let  $x \in g^{-1}(A) \cap g^{-1}(B)$ , this is the same as  $x \in g^{-1}(A)$  and  $x \in g^{-1}(B)$  which is the same as  $g(x) \in A$  and  $g(x) \in B$  implying  $g(x) \in A \cap B$  and thus  $x \in g^{-1}(A \cap B)$ .

**Exercise 1.2.10**

Decide which of the following are true statements. Provide a short justification for those that are valid and a counterexample for those that are not:

- (a) Two real numbers satisfy  $a < b$  if and only if  $a < b + \epsilon$  for every  $\epsilon > 0$ .
- (b) Two real numbers satisfy  $a < b$  if  $a < b + \epsilon$  for every  $\epsilon > 0$ .
- (c) Two real numbers satisfy  $a \leq b$  if and only if  $a < b + \epsilon$  for every  $\epsilon > 0$ .

**Solution**

- (a) False, if  $a = b$  then  $a < b + \epsilon$  for all  $\epsilon > 0$  but  $a \not< b$
- (b) False, see above
- (c) True, suppose  $a < b + \epsilon$  for all  $\epsilon > 0$ . We want to show this implies  $a \leq b$ . We either have  $a \leq b$  or  $a > b$ , but  $a > b$  is impossible since the gap implies there exists an  $\epsilon$  small enough such that  $a > b + \epsilon$ . Now suppose  $a \leq b$ , obviously  $a < b + \epsilon$  for all  $\epsilon > 0$ .

**Exercise 1.2.11**

Form the logical negation of each claim. One trivial way to do this is to simply add “It is not the case that...” in front of each assertion. To make this interesting, fashion the negation into a positive statement that avoids using the word “not” altogether. In each case, make an intuitive guess as to whether the claim or its negation is the true statement.

- (a) For all real numbers satisfying  $a < b$ , there exists an  $n \in \mathbf{N}$  such that  $a + 1/n < b$
- (b) There exists a real number  $x > 0$  such that  $x < 1/n$  for all  $n \in \mathbf{N}$ .
- (c) Between every two distinct real numbers there is a rational number.

**Solution**

- (a) There exist real numbers satisfying  $a < b$  where  $a + 1/n \geq b$  for all  $n \in \mathbf{N}$  (false).

- (b) For every real number  $x > 0$  there exists an  $n \in \mathbf{N}$  such that  $x < 1/n$  (true).
- (c) There exist two real numbers  $a < b$  such that if  $r < b$  then  $r < a$  for all  $r \in \mathbf{Q}$  (false).

**Exercise 1.2.12**

Let  $y_1 = 6$ , and for each  $n \in \mathbf{N}$  define  $y_{n+1} = (2y_n - 6)/3$

- (a) Use induction to prove that the sequence satisfies  $y_n > -6$  for all  $n \in \mathbf{N}$ .
- (b) Use another induction argument to show the sequence  $(y_1, y_2, y_3, \dots)$  is decreasing.

**Solution**

- (a) Suppose  $y_n > -6$ , then  $y_{n+1} = (2y_n - 6)/3$  implying  $y_n = (3y_{n+1} + 6)/2 > -6$  implying  $y_{n+1} > -6$  by basic algebra.
- (b) Suppose  $y_{n+1} < y_n$  this implies  $2y_{n+1} < 2y_n$  implying  $2y_{n+1} - 6 < 2y_n - 6$  and finally  $(2y_{n+1} - 6)/3 < (2y_n - 6)/3$  which shows  $y_{n+2} < y_{n+1}$ .

**Exercise 1.2.13**

For this exercise, assume Exercise 1.2.5 has been successfully completed.

- (a) Show how induction can be used to conclude that

$$(A_1 \cup A_2 \cup \dots \cup A_n)^c = A_1^c \cap A_2^c \cap \dots \cap A_n^c$$

for any finite  $n \in \mathbf{N}$

- (b) It is tempting to appeal to induction to conclude

$$\left( \bigcup_{i=1}^{\infty} A_i \right)^c = \bigcap_{i=1}^{\infty} A_i^c$$

but induction does not apply here. Induction is used to prove that a particular statement holds for every value of  $n \in \mathbf{N}$ , but this does not imply the validity of the infinite case. To illustrate this point, find an example of a collection of sets  $B_1, B_2, B_3, \dots$  where  $\bigcap_{i=1}^n B_i \neq \emptyset$  is true for every  $n \in \mathbf{N}$ , but  $\bigcap_{i=1}^{\infty} B_i \neq \emptyset$  fails.

- (c) Nevertheless, the infinite version of De Morgan's Law stated in (b) is a valid statement. Provide a proof that does not use induction.

**Solution**

- (a) 1.2.5 Is our base case, Assume  $(A_1 \cup \dots \cup A_n)^c = A_1^c \cap \dots \cap A_n^c$ . We want to show the  $n + 1$  case. Using associativity we have

$$\begin{aligned} ((A_1 \cup \dots \cup A_n) \cup A_{n+1})^c &= (A_1 \cup \dots \cup A_n)^c \cap A_{n+1}^c \\ &= (A_1^c \cap \dots \cap A_n^c) \cap A_{n+1}^c \\ &= A_1^c \cap \dots \cap A_n^c \cap A_{n+1}^c \end{aligned}$$

- (b)  $B_1 = \{1, 2, \dots\}, B_2 = \{2, 3, \dots\}, \dots$

- (c) First suppose  $x \in (\bigcap_{i=1}^{\infty} A_i)^c$ , then  $x \notin \bigcap_{i=1}^{\infty} A_i$  meaning  $x \notin A_i$  for some  $i$ , which is the same as  $x \in A_i^c$  for some  $i$ , meaning  $x \in \bigcup_{i=1}^{\infty} A_i^c$ . This shows

$$\left( \bigcap_{i=1}^{\infty} A_i \right)^c \subseteq \bigcup_{i=1}^{\infty} A_i^c$$

Now suppose  $x \in \bigcup_{i=1}^{\infty} A_i^c$  meaning  $x \notin A_i$  for some  $i$ , which is the same as  $x \notin \bigcap_{i=1}^{\infty} A_i$  implying  $x \notin (\bigcap_{i=1}^{\infty} A_i)^c$ . This shows inclusion the other way and completes the proof.

## 1.3 The Axiom of Completeness

### Exercise 1.3.1

- (a) Write a formal definition in the style of Definition 1.3.2 for the *infimum* or *greatest lower bound* of a set.
- (b) Now, state and prove a version of Lemma 1.3.8 for greatest lower bounds.

### Solution

- (a) We have  $i = \inf A$  if and only if
  - (i) Lower bound,  $a \geq i$  for all  $a \in A$
  - (ii) Greatest lower bound, If  $b$  is a lower bound on  $A$  then  $b \leq i$
- (b) Theorem: Suppose  $i$  is a lower bound for  $A$ , it is the greatest lower bound if and only if for all  $\epsilon > 0$ , there exists an  $a \in A$  such that  $i + \epsilon < a$ .  
 Proof:  
 ( $\implies$ ) Suppose  $i = \inf A$ , then any  $i + \epsilon$  cannot be a lower bound since  $i$  is defined as the greatest lower bound, and  $i + \epsilon > i$ .  
 ( $\impliedby$ ) Suppose  $j > i$  is a lower bound on  $A$ , then set  $\epsilon = j - i$  and we have that  $i + \epsilon = j$  is not a lower bound since there exists  $a \in A$  such that  $j > a$ . Therefore  $i$  is the greatest lower bound.

### Exercise 1.3.2

Give an example of each of the following, or state that the request is impossible.

- (a) A set  $B$  with  $\inf B \geq \sup B$ .
- (b) A finite set that contains its infimum but not its supremum.
- (c) A bounded subset of  $\mathbf{Q}$  that contains its supremum but not its infimum.

### Solution

- (a) Let  $B = \{0\}$  we have  $\inf B = 0$  and  $\sup B = 0$  thus  $\inf B \leq \sup B$ .
- (b) Impossible, finite sets must contain their infimum and supremum.
- (c) Let  $B = \{r \in \mathbf{Q} \mid 1 < r \leq 2\}$  we have  $\inf B = 1 \notin B$  and  $\sup B = 2 \in B$ .

### Exercise 1.3.3

- (a) Let  $A$  be nonempty and bounded below, and define  $B = \{b \in \mathbf{R} : b \text{ is a lower bound for } A\}$ . Show that  $\sup B = \inf A$ .
- (b) Use (a) to explain why there is no need to assert that greatest lower bounds exist as part of the Axiom of Completeness.

### Solution

- (a) By definition  $\sup B$  is the greatest lower bound for  $A$ , meaning it equals  $\inf A$ .

- (b) (a) Proves the greatest lower bound exists using the least upper bound.

**Exercise 1.3.4**

Let  $A_1, A_2, A_3, \dots$  be a collection of nonempty sets, each of which is bounded above.

- (a) Find a formula for  $\sup(A_1 \cup A_2)$ . Extend this to  $\sup(\bigcup_{k=1}^n A_k)$ .  
 (b) Consider  $\sup(\bigcup_{k=1}^{\infty} A_k)$ . Does the formula in (a) extend to the infinite case?

**Solution**

- (a)  $\sup(\bigcup_{k=1}^n A_k) = \sup\{\sup A_k \mid k = 1, \dots, n\}$   
 (b) Yes. Let  $S = \{\sup A_k \mid k = 1, \dots, \infty\}$  and  $s = \sup S$ .  $s$  is obviously an upper bound for  $\bigcup_{k=1}^{\infty} A_k$ . to see it is the least upper bound suppose  $s' < s$ , then by definition there exists a  $k$  such that  $\sup A_k > s'$  implying  $s'$  is not an upper bound for  $A_k$ . Therefore  $s$  is the least upper bound.

**Exercise 1.3.5**

As in Example 1.3.7, let  $A \subseteq \mathbf{R}$  be nonempty and bounded above, and let  $c \in \mathbf{R}$ . This time define the set  $cA = \{ca : a \in A\}$ .

- (a) If  $c \geq 0$ , show that  $\sup(cA) = c \sup A$ .  
 (b) Postulate a similar type of statement for  $\sup(cA)$  for the case  $c < 0$ .

**Solution**

- (a) Let  $s = c \sup A$ . Suppose  $ca > s$ , then  $a > \sup A$  which is impossible, meaning  $s$  is an upper bound on  $cA$ . Now suppose  $s'$  is an upper bound on  $cA$  and  $s' < s$ . Then  $s'/c < s/c$  and  $s'/c < \sup A$  meaning  $s'/c$  cannot bound  $A$ , so there exists  $a \in A$  such that  $s'/c > a$  meaning  $s' > ca$  thus  $s'$  cannot be an upper bound on  $cA$ , and so  $s = c \sup A$  is the least upper bound.  
 (b)  $\sup(cA) = c \inf(A)$  for  $c < 0$

**Exercise 1.3.6**

Given sets  $A$  and  $B$ , define  $A + B = \{a + b : a \in A \text{ and } b \in B\}$ . Follow these steps to prove that if  $A$  and  $B$  are nonempty and bounded above then  $\sup(A + B) = \sup A + \sup B$

- (a) Let  $s = \sup A$  and  $t = \sup B$ . Show  $s + t$  is an upper bound for  $A + B$ .  
 (b) Now let  $u$  be an arbitrary upper bound for  $A + B$ , and temporarily fix  $a \in A$ . Show  $t \leq u - a$ .  
 (c) Finally, show  $\sup(A + B) = s + t$ .  
 (d) Construct another proof of this same fact using Lemma 1.3.8.

**Solution**

- (a) We have  $a \leq s$  and  $b \leq t$ , adding the equations gives  $a + b \leq s + t$ .

- (b)  $t \leq u - a$  should be true since  $u - a$  is an upper bound on  $b$ , meaning it is greater than or equal to the least upper bound  $t$ . Formally  $a + b \leq u$  implies  $b \leq u - a$  and since  $t$  is the least upper bound on  $b$  we have  $t \leq u - a$ .

- (c) From (a) we know  $s + t$  is an upper bound, so we must only show it is the least upper bound.

Let  $u = \sup(A + B)$ , from (a) we have  $t \leq u - a$  and  $s \leq u - b$  adding and rearranging gives  $a + b \leq 2u - s - t$ . since  $2u - s - t$  is an upper bound on  $A + B$  it is less than the least upper bound, so  $u \leq 2u - s - t$  implying  $s + t \leq u$ . and since  $u$  is the least upper bound  $s + t$  must equal  $u$ .

Stepping back, the key to this proof is that  $a + b \leq s, \forall a, b$  implying  $\sup(A + B) \leq s$  can be used to transition from all  $a + b$  to a single value  $\sup(A + B)$ , avoiding the  $\epsilon$ -hackery I would otherwise use.

- (d) Showing  $s + t - \epsilon$  is not an upper bound for any  $\epsilon > 0$  proves it is the least upper bound by Lemma 1.3.8. Rearranging gives  $(s - \epsilon/2) + (t - \epsilon/2)$  we know there exists  $a > (s - \epsilon/2)$  and  $b > (t - \epsilon/2)$  therefore  $a + b > s + t - \epsilon$  meaning  $s + t$  cannot be made smaller, and thus is the least upper bound.

### Exercise 1.3.7

Prove that if  $a$  is an upper bound for  $A$ , and if  $a$  is also an element of  $A$ , then it must be that  $a = \sup A$ .

### Solution

$a$  is the least upper bound since any smaller bound  $a' < a$  would not bound  $a$ .

### Exercise 1.3.8

Compute, without proofs, the suprema and infima (if they exist) of the following sets:

- (a)  $\{m/n : m, n \in \mathbf{N} \text{ with } m < n\}$ .
- (b)  $\{(-1)^m/n : m, n \in \mathbf{N}\}$ .
- (c)  $\{n/(3n + 1) : n \in \mathbf{N}\}$
- (d)  $\{m/(m + n) : m, n \in \mathbf{N}\}$

### Solution

- (a)  $\sup = 1, \inf = 0$
- (b)  $\sup = 1, \inf = -1$
- (c)  $\sup = 1/3, \inf = 1/4$
- (d)  $\sup = 1, \inf = 0$

### Exercise 1.3.9

- (a) If  $\sup A < \sup B$ , show that there exists an element  $b \in B$  that is an upper bound for  $A$ .



- (b) Give an example to show that this is not always the case if we only assume  $\sup A \leq \sup B$

**Solution**

- (a) By Lemma 1.3.8 we know there exists a  $b$  such that  $(\sup B) - \epsilon < b$  for any  $\epsilon > 0$ . We set  $\epsilon$  to be small enough that  $\sup A < (\sup B) - \epsilon$  meaning  $\sup A < b$  for some  $b$ , and thus  $b$  is an upper bound on  $A$ .
- (b)  $A = \{x \mid x \leq 1\}$ ,  $B = \{x \mid x < 1\}$  no  $b \in B$  is an upper bound since  $1 \in A$  and  $1 > b$ .

**Exercise 1.3.10 (Cut Property)**

The Cut Property of the real numbers is the following:

If  $A$  and  $B$  are nonempty, disjoint sets with  $A \cup B = \mathbf{R}$  and  $a < b$  for all  $a \in A$  and  $b \in B$ , then there exists  $c \in \mathbf{R}$  such that  $x \leq c$  whenever  $x \in A$  and  $x \geq c$  whenever  $x \in B$ .

- (a) Use the Axiom of Completeness to prove the Cut Property.
- (b) Show that the implication goes the other way; that is, assume  $\mathbf{R}$  possesses the Cut Property and let  $E$  be a nonempty set that is bounded above. Prove  $\sup E$  exists.
- (c) The punchline of parts (a) and (b) is that the Cut Property could be used in place of the Axiom of Completeness as the fundamental axiom that distinguishes the real numbers from the rational numbers. To drive this point home, give a concrete example showing that the Cut Property is not a valid statement when  $\mathbf{R}$  is replaced by  $\mathbf{Q}$ .

**Solution**

- (a) If  $c = \sup A = \inf B$  then  $a \leq c \leq b$  is obvious. So we must only prove  $\sup A = \inf B$ . If  $\sup A < \inf B$  then we can find  $c$  between  $A$  and  $B$  implying  $A \cup B \neq \mathbf{R}$ . If  $\sup A > \inf B$  then we can find  $a$  such that  $a > b$  by subtracting  $\epsilon > 0$  and using the least upper/lower bound facts, similarly to Lemma 1.3.8. Thus  $\sup A$  must equal  $\inf B$  since we have shown both alternatives are impossible.
- (b) Let  $B = \{x \mid e < x, \forall e \in E\}$  and let  $A = B^c$ . Clearly  $a < b$  so the cut property applies. We have  $a \leq c \leq b$  and must show the two conditions for  $c = \sup E$
- (i) Since  $E \subseteq A$ ,  $a \leq c$  implies  $e \leq c$  thus  $c$  is an upper bound.
- (ii)  $c \leq b$  implies  $c$  is the smallest upper bound.

Note: Using (a) here would be wrong, it assumes the axiom of completeness so we would be making a circular argument.

- (c)  $A = \{r \in \mathbf{Q} \mid r^2 < 2\}$ ,  $B = A^c$  does not satisfy the cut property in  $\mathbf{Q}$  since  $\sqrt{2} \notin \mathbf{Q}$

**Exercise 1.3.11**

Decide if the following statements about suprema and infima are true or false. Give a short proof for those that are true. For any that are false, supply an example where the claim in question does not appear to hold.

- (a) If  $A$  and  $B$  are nonempty, bounded, and satisfy  $A \subseteq B$ , then  $\sup A \leq \sup B$ .

- (b) If  $\sup A < \inf B$  for sets  $A$  and  $B$ , then there exists a  $c \in \mathbf{R}$  satisfying  $a < c < b$  for all  $a \in A$  and  $b \in B$ .
- (c) If there exists a  $c \in \mathbf{R}$  satisfying  $a < c < b$  for all  $a \in A$  and  $b \in B$ , then  $\sup A < \inf B$ .

**Solution**

- (a) True. We know  $a \leq \sup A$  and  $a \leq \sup B$  since  $A \subseteq B$ . since  $\sup A$  is the least upper bound on  $A$  we have  $\sup A \leq \sup B$ .
- (b) True. Let  $c = (\sup A + \inf B)/2$ ,  $c > \sup A$  implies  $a < c$  and  $c < \inf B$  implies  $c < b$  giving  $a < c < b$  as desired.
- (c) False. consider  $A = \{x \mid x < 1\}$ ,  $B = \{x \mid x > 1\}$ ,  $a < 1 < b$  but  $\sup A \not< \inf B$  since  $1 \not< 1$ .

## 1.4 Consequences of Completeness

### Exercise 1.4.1

Recall that  $\mathbf{I}$  stands for the set of irrational numbers.

- (a) Show that if  $a, b \in \mathbf{Q}$ , then  $ab$  and  $a + b$  are elements of  $\mathbf{Q}$  as well.
- (b) Show that if  $a \in \mathbf{Q}$  and  $t \in \mathbf{I}$ , then  $a + t \in \mathbf{I}$  and  $at \in \mathbf{I}$  as long as  $a \neq 0$ .
- (c) Part (a) can be summarized by saying that  $\mathbf{Q}$  is closed under addition and multiplication. Is  $\mathbf{I}$  closed under addition and multiplication? Given two irrational numbers  $s$  and  $t$ , what can we say about  $s + t$  and  $st$ ?

### Solution

- (a) Trivial.
- (b) Suppose  $a + t \in \mathbf{Q}$ , then by (a)  $(a + t) - a = t \in \mathbf{Q}$  contradicting  $t \in \mathbf{I}$ .
- (c)  $\mathbf{I}$  is not closed under addition or multiplication. consider  $(1 - \sqrt{2}) \in \mathbf{I}$  by (b), and  $\sqrt{2} \in \mathbf{I}$ . the sum  $(1 - \sqrt{2}) + \sqrt{2} = 1 \in \mathbf{Q} \notin \mathbf{I}$ . Also  $\sqrt{2} \cdot \sqrt{2} = 2 \in \mathbf{Q} \notin \mathbf{I}$ .

### Exercise 1.4.2

Let  $A \subseteq \mathbf{R}$  be nonempty and bounded above, and let  $s \in \mathbf{R}$  have the property that for all  $n \in \mathbf{N}$ ,  $s + \frac{1}{n}$  is an upper bound for  $A$  and  $s - \frac{1}{n}$  is not an upper bound for  $A$ . Show  $s = \sup A$ .

### Solution

This is basically a rephrasing of Lemma 1.3.8 using the archimedean property. The most straightforward approach is to argue by contradiction:

- (i) If  $s < \sup A$  then there exists an  $n$  such that  $s + 1/n < \sup A$  contradicting  $\sup A$  being the least upper bound.
- (ii) If  $s > \sup A$  then there exists an  $n$  such that  $s - 1/n > \sup A$  where  $s - 1/n$  is not an upper bound, contradicting  $\sup A$  being an upper bound.

Thus  $s = \sup A$  is the only remaining possibility.

### Exercise 1.4.3

Prove that  $\bigcap_{n=1}^{\infty} (0, 1/n) = \emptyset$ . Notice that this demonstrates that the intervals in the Nested Interval Property must be closed for the conclusion of the theorem to hold.

### Solution

Suppose  $x \in \bigcap_{n=1}^{\infty} (0, 1/n)$ , then we have  $0 < x < 1/n$  for all  $n$ , which is impossible by the archimedean property. In other words we can always set  $n$  large enough that  $x$  lies outside the interval.

### Exercise 1.4.4

Let  $a < b$  be real numbers and consider the set  $T = \mathbf{Q} \cap [a, b]$ . Show  $\sup T = b$

**Solution**

We must show the two conditions for a least upper bound

- (i) Clearly  $t \leq b$  for all  $t \in T$
- (ii) Let  $a < b' < b$ .  $b'$  Cannot be an upper bound for  $T$  since the density theorem tells us we can find  $r \in \mathbf{Q} \cap [a, b]$  such that  $b' < r < b$ .

**Exercise 1.4.5**

Using Exercise 1.4.1, supply a proof that  $\mathbf{I}$  is dense in  $\mathbf{R}$  by considering the real numbers  $a - \sqrt{2}$  and  $b - \sqrt{2}$ . In other words show for every two real numbers  $a < b$  there exists an irrational number  $t$  with  $a < t < b$ .

**Solution**

The density theorem lets us find a rational number  $r$  with  $a - \sqrt{2} < r < b - \sqrt{2}$ , adding  $\sqrt{2}$  to both sides gives  $a < r + \sqrt{2} < b$ . From 1.4.1 we know  $r + \sqrt{2}$  is irrational, so setting  $t = r + \sqrt{2}$  gives  $a < t < b$  as desired.

**Exercise 1.4.6**

Recall that a set  $B$  is dense in  $\mathbf{R}$  if an element of  $B$  can be found between any two real numbers  $a < b$ . Which of the following sets are dense in  $\mathbf{R}$ ? Take  $p \in \mathbf{Z}$  and  $q \in \mathbf{N}$  in every case.

- (a) The set of all rational numbers  $p/q$  with  $q \leq 10$ .
- (b) The set of all rational numbers  $p/q$  with  $q$  a power of 2.
- (c) The set of all rational numbers  $p/q$  with  $10|p| \geq q$ .

**Solution**

- (a) Dense.
- (b) Dense.
- (c) Not dense since we cannot make  $|p|/q$  smaller than  $1/10$ .

**Exercise 1.4.7**

Finish the proof of Theorem 1.4.5 by showing that the assumption  $\alpha^2 > 2$  leads to a contradiction of the fact that  $\alpha = \sup T$

**Solution**

Recall  $T = \{t \in \mathbf{R} \mid t^2 < 2\}$  and  $\alpha = \sup T$ . suppose  $\alpha^2 > 2$ , we will show there exists an  $n \in \mathbf{N}$  such that  $(\alpha - 1/n)^2 > 2$  contradicting the assumption that  $\alpha$  is the least upper bound.

We expand  $(\alpha - 1/n)^2$  to find  $n$  such that  $(\alpha^2 - 1/n) > 2$

$$2 < (\alpha - 1/n)^2 = \alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} < \alpha^2 + \frac{1 - 2\alpha}{n}$$

Then

$$2 < \alpha^2 + \frac{1 - 2\alpha}{n} \implies n(2 - \alpha^2) < 1 - 2\alpha$$

Since  $2 - \alpha^2 < 0$  dividing reverses the inequality gives us

$$n > \frac{1 - 2\alpha}{2 - \alpha^2}$$

This contradicts  $\alpha^2 > 2$  since we have shown  $n$  can be picked such that  $(\alpha^2 - 1/n) > 2$  meaning  $\alpha$  is not the least upper bound.

### Exercise 1.4.8

Give an example of each or state that the request is impossible. When a request is impossible, provide a compelling argument for why this is the case.

- (a) Two sets  $A$  and  $B$  with  $A \cap B = \emptyset$ ,  $\sup A = \sup B$ ,  $\sup A \notin A$  and  $\sup B \notin B$ .
- (b) A sequence of nested open intervals  $J_1 \supseteq J_2 \supseteq J_3 \supseteq \cdots$  with  $\bigcap_{n=1}^{\infty} J_n$  nonempty but containing only a finite number of elements.
- (c) A sequence of nested unbounded closed intervals  $L_1 \supseteq L_2 \supseteq L_3 \supseteq \cdots$  with  $\bigcap_{n=1}^{\infty} L_n = \emptyset$ . (An unbounded closed interval has the form  $[a, \infty) = \{x \in \mathbb{R} : x \geq a\}$ .)
- (d) A sequence of closed bounded (not necessarily nested) intervals  $I_1, I_2, I_3, \dots$  with the property that  $\bigcap_{n=1}^N I_n \neq \emptyset$  for all  $N \in \mathbb{N}$ , but  $\bigcap_{n=1}^{\infty} I_n = \emptyset$ .

### Solution

- (a)  $A = \mathbf{Q} \cap (0, 1)$ ,  $B = \mathbf{I} \cap (0, 1)$ .  $A \cap B = \emptyset$ ,  $\sup A = \sup B = 1$  and  $1 \notin A$ ,  $1 \notin B$ .
- (b) Impossible.  $\bigcap_{n=1}^{\infty} J_n$  is the same as asking what happens to  $J_n$  as  $n$  goes to  $\infty$ . since every  $J_n$  is nonempty,  $\bigcap_{n=1}^{\infty} J_n = J_{\infty}$  must have an uncountably infinite number of elements.
- (c)  $L_n = [n, \infty)$  has  $\bigcap_{n=1}^{\infty} L_n = \emptyset$
- (d) Impossible. Let  $J_n = \bigcap_{k=1}^n I_k$  and observe the following
  - (i) Since  $\bigcap_{n=1}^N I_n \neq \emptyset$  we have  $J_n \neq \emptyset$ .
  - (ii)  $J_n$  being the intersection of closed intervals makes it a closed interval.
  - (iii)  $J_{n+1} \subseteq J_n$  since  $I_{n+1} \cap J_n \subseteq J_n$
  - (iv)  $\bigcap_{n=1}^{\infty} J_n = \bigcap_{n=1}^{\infty} (\bigcap_{k=1}^n I_k) = \bigcap_{n=1}^{\infty} I_n$

By (i), (ii) and (iii) the Nested Interval Property tells us  $\bigcap_{n=1}^{\infty} J_n \neq \emptyset$ . Therefore by (iv)  $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$ .

## 1.5 Cardinality

### Exercise 1.5.1

Finish the following proof for Theorem 1.5.7. Assume  $B$  is a countable set. Thus, there exists  $f : \mathbf{N} \rightarrow B$ , which is 1-1 and onto. Let  $A \subseteq B$  be an infinite subset of  $B$ . We must show that  $A$  is countable.

Let  $n_1 = \min\{n \in \mathbf{N} : f(n) \in A\}$ . As a start to a definition of  $g : \mathbf{N} \rightarrow A$  set  $g(1) = f(n_1)$ . Show how to inductively continue this process to produce a 1-1 function  $g$  from  $\mathbf{N}$  onto  $A$ .

### Solution

Let  $n_k = \min\{n \in \mathbf{N} \mid f(n) \in A, n \notin \{n_1, n_2, \dots, n_{k-1}\}\}$  and  $g(k) = f(n_k)$ . since  $g : \mathbf{N} \rightarrow A$  is 1-1 and onto,  $A$  is countable.

### Exercise 1.5.2

Review the proof of Theorem 1.5.6, part (ii) showing that  $\mathbf{R}$  is uncountable, and then find the flaw in the following erroneous proof that  $\mathbf{Q}$  is uncountable:

Assume, for contradiction, that  $\mathbf{Q}$  is countable. Thus we can write  $\mathbf{Q} = \{r_1, r_2, r_3, \dots\}$  and, as before, construct a nested sequence of closed intervals with  $r_n \notin I_n$ . Our construction implies  $\bigcap_{n=1}^{\infty} I_n = \emptyset$  while NIP implies  $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$ . This contradiction implies  $\mathbf{Q}$  must therefore be uncountable.

### Solution

The nested interval property is not true for  $\mathbf{Q}$ . Consider  $I_n$  being rational bounds for  $\sqrt{2}$  with  $n$  decimal places, then  $\bigcap_{n=1}^{\infty} I_n = \emptyset$  since  $\sqrt{2} \notin \mathbf{Q}$ .

### Exercise 1.5.3

- (a) Prove if  $A_1, \dots, A_m$  are countable sets then  $A_1 \cup \dots \cup A_m$  is countable.
- (b) Explain why induction *cannot* be used to prove that if each  $A_n$  is countable, then  $\bigcup_{n=1}^{\infty} A_n$  is countable.
- (c) Show how arranging  $\mathbf{N}$  into the two-dimensional array

$$\begin{array}{cccccc} 1 & 3 & 6 & 10 & 15 & \dots \\ 2 & 5 & 9 & 14 & \dots & \\ 4 & 8 & 13 & \dots & & \\ 7 & 12 & \dots & & & \\ 11 & \dots & & & & \\ \vdots & & & & & \end{array}$$

leads to a proof for the infinite case.

### Solution

- (a) Let  $B, C$  be disjoint countable sets. We use the same trick as with the integers and list them as

$$B \cup C = \{b_1, c_1, b_2, c_2, \dots\}$$

Meaning  $B \cup C$  is countable, and  $A_1 \cup A_2$  is also countable since we can let  $B = A_1$  and  $C = A_2 \setminus A_1$ .

Now induction: suppose  $A_1 \cup \dots \cup A_n$  is countable,  $(A_1 \cup \dots \cup A_n) \cup A_{n+1}$  is the union of two countable sets which by above is countable.

- (b) Induction shows something for each  $n \in \mathbf{N}$ , it does not apply in the infinite case.
- (c) Rearranging  $\mathbf{N}$  as in (c) gives us disjoint sets  $C_n$  such that  $\bigcup_{n=1}^{\infty} C_n = \mathbf{N}$ . Let  $B_n$  be disjoint, constructed as  $B_1 = A_1, B_2 = A_1 \setminus B_1, \dots$  we want to do something like

$$f(\mathbf{N}) = f\left(\bigcup_{n=1}^{\infty} C_n\right) = \bigcup_{n=1}^{\infty} f_n(C_n) = \bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n$$

Let  $f_n : C_n \rightarrow B_n$  be bijective since  $B_n$  is countable, define  $f : \mathbf{N} \rightarrow \bigcup_{n=1}^{\infty} B_n$  as

$$f(n) = \begin{cases} f_1(n) & \text{if } n \in C_1 \\ f_2(n) & \text{if } n \in C_2 \\ \vdots & \end{cases}$$

- (i) Since each  $C_n$  is disjoint and each  $f_n$  is 1-1,  $f(n_1) = f(n_2)$  implies  $n_1 = n_2$  meaning  $f$  is 1-1.
- (ii) Since any  $b \in \bigcup_{n=1}^{\infty} B_n$  has  $b \in B_n$  for some  $n$ , we know  $b = f_n(x)$  has a solution since  $f_n$  is onto. Letting  $x = f_n^{-1}(b)$  we have  $f(x) = f_n(x) = b$  since  $f_n^{-1}(b) \in C_n$  meaning  $f$  is onto.

By (i) and (ii)  $f$  is bijective and so  $\bigcup_{n=1}^{\infty} B_n$  is countable. And since

$$\bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n$$

We have that  $\bigcup_{n=1}^{\infty} A_n$  is countable, completing the proof.

#### Exercise 1.5.4

- (a) Show  $(a, b) \sim \mathbf{R}$  for any interval  $(a, b)$ .
- (b) Show that an unbounded interval like  $(a, \infty) = \{x : x > a\}$  has the same cardinality as  $\mathbf{R}$  as well.
- (c) Using open intervals makes it more convenient to produce the required 1-1, onto functions, but it is not really necessary. Show that  $[0, 1) \sim (0, 1)$  by exhibiting a 1-1 onto function between the two sets.

#### Solution

- (a) We will start by finding  $f : (-1, 1) \rightarrow \mathbf{R}$  and then transform it to  $(a, b)$ . Example 1.5.4 gives a suitable  $f$

$$f(x) = \frac{x}{x^2 - 1}$$

The book says to use calculus to show  $f$  is bijective, first we will examine the derivative

$$f'(x) = \frac{x^2 - 1 - 2x^2}{(x^2 - 1)^2} = -\frac{x^2 + 1}{(x^2 - 1)^2}$$

The denominator and numerator are positive, so  $f'(x) < 0$  for all  $x \in (0, 1)$ . This means no two inputs will be mapped to the same output, meaning  $f$  is one to one (a rigorous proof is beyond our current ability)

To show that  $f$  is onto, we examine the limits

$$\begin{aligned}\lim_{x \rightarrow 1^-} \frac{x}{x^2 - 1} &= -\infty \\ \lim_{x \rightarrow -1^+} \frac{x}{x^2 - 1} &= +\infty\end{aligned}$$

Then use the intermediate value theorem to conclude  $f$  is onto.

Now we shift  $f$  to the interval  $(a, b)$

$$g(x) = f\left(\frac{2x - 1}{b - a} - a\right)$$

Proving  $g(x)$  is also bijective is a straightforward application of the chain rule.

- (b) We want a bijective  $h(x)$  such that  $h(x) : (a, \infty) \rightarrow (-1, 1)$  because then we could compose them to get a new bijective function  $f(h(x)) : (a, \infty) \rightarrow \mathbf{R}$ .

Let

$$h(x) = \frac{2}{x - a + 1} - 1$$

We have  $h : (a, \infty) \rightarrow (1, -1)$  since  $h(a) = 1$  and  $\lim_{x \rightarrow \infty} h(x) = -1$ .

Meaning that  $f(h(x)) : (a, \infty) \rightarrow \mathbf{R}$  is our bijective map.

- (c) With countable sets adding a single element doesn't change cardinality since we can just shift by one to get a bijective map. we'll use a similar technique here to essentially outrun our problems. Define  $f : [0, 1) \rightarrow (0, 1)$  as

$$f(x) = \begin{cases} 1/2 & \text{if } x = 0 \\ 1/4 & \text{if } x = 1/2 \\ 1/8 & \text{if } x = 1/4 \\ \vdots & \\ x & \text{otherwise} \end{cases}$$

Now we prove  $f$  is bijective by showing  $y = f(x)$  has exactly one solution for all  $y \in (0, 1)$ .

If  $y = 1/2^n$  then the only solution is  $y = f(1/2^{n-1})$  (or  $x = 0$  in the special case  $n = 1$ ),  
If  $y \neq 1/2^n$  then the only solution is  $y = f(y)$ .



**Exercise 1.5.5**

- (a) Why is  $A \sim A$  for every set  $A$ ?
- (b) Given sets  $A$  and  $B$ , explain why  $A \sim B$  is equivalent to asserting  $B \sim A$ .
- (c) For three sets  $A, B$ , and  $C$ , show that  $A \sim B$  and  $B \sim C$  implies  $A \sim C$ . These three properties are what is meant by saying that  $\sim$  is an *equivalence relation*.

**Solution**

- (a) The identity function  $f(x) = x$  is a bijection
- (b) If  $f : A \rightarrow B$  is bijective then  $f^{-1} : B \rightarrow A$  is bijective.
- (c) Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$ , since  $g \circ f : A \rightarrow C$  is bijective we have  $A \sim C$ .

**Exercise 1.5.6**

- (a) Give an example of a countable collection of disjoint open intervals.
- (b) Give an example of an uncountable collection of disjoint open intervals, or argue that no such collection exists.

**Solution**

- (a)  $I_1 = (0, 1)$ ,  $I_2 = (1, 2)$  and in general  $I_n = (n - 1, n)$
- (b) Let  $A$  denote this set. Intuitively no such collection should exist since each  $I_n$  has nonzero length.

The key here is to try and show  $A \sim \mathbf{Q}$  instead of directly showing  $A \sim \mathbf{N}$ .

For any nonempty interval  $I_n$  the density theorem tells us there exists an  $r \in \mathbf{Q}$  such that  $r \in I_n$ . Assigning each  $I \in A$  a rational number  $r \in I$  proves  $I \subseteq \mathbf{Q}$  and thus  $I$  is countable.

**Exercise 1.5.7**

Consider the open interval  $(0, 1)$ , and let  $S$  be the set of points in the open unit square; that is,  $S = \{(x, y) : 0 < x, y < 1\}$ .

- (a) Find a 1-1 function that maps  $(0, 1)$  into, but not necessarily onto,  $S$ . (This is easy.)
- (b) Use the fact that every real number has a decimal expansion to produce a 1-1 function that maps  $S$  into  $(0, 1)$ . Discuss whether the formulated function is onto. (Keep in mind that any terminating decimal expansion such as .235 represents the same real number as .234999...)

The Schröder-Bernstein Theorem discussed in Exercise 1.5.11 can now be applied to conclude that  $(0, 1) \sim S$ .

**Solution**

- (a) We scale and shift up into the square.  $f(x) = \frac{1}{2}x + \frac{1}{3}$

- (b) Let  $g : S \rightarrow (0, 1)$  be a function that interleaves decimals in the representation without trailing nines.  $g(0.32, 0.45) = 0.3425$  and  $g(0.1\bar{9}, 0.2) = g(0.2, 0.2) = 0.22$  etc.

Every real number can be written with two digit representations, one with trailing 9's and one without. However  $g(x, y) = 0.d_1d_2\ldots\bar{9}$  is impossible since it would imply  $x = 0.d_1\ldots\bar{9}$  and  $y = 0.d_2\ldots\bar{9}$  but the definition of  $g$  forbids this. Therefore  $g(s)$  is unique, and so  $g$  is 1-1.

Is  $g$  onto? No since  $g(x, y) = 0.1$  has no solutions, since we would want  $x = 0.1$  and  $y = 0$  but  $0 \notin (0, 1)$ .

### Exercise 1.5.8

Let  $B$  be a set of positive real numbers with the property that adding together any finite subset of elements from  $B$  always gives a sum of 2 or less. Show  $B$  must be finite or countable.

### Solution

Notice  $B \cap (a, 2)$  is finite for all  $a > 0$ , since if it was infinite we could make a set with sum greater than two. And since  $B$  is the countable union of finite sets  $\bigcup_{n=1}^{\infty} B \cap (1/n, 2)$ ,  $B$  must be countable or finite.

### Exercise 1.5.9

A real number  $x \in \mathbf{R}$  is called algebraic if there exist integers  $a_0, a_1, a_2, \dots, a_n \in \mathbf{Z}$ , not all zero, such that

$$a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0 = 0$$

Said another way, a real number is algebraic if it is the root of a polynomial with integer coefficients. Real numbers that are not algebraic are called *transcendental* numbers. Reread the last paragraph of Section 1.1. The final question posed here is closely related to the question of whether or not transcendental numbers exist.

- Show that  $\sqrt{2}$ ,  $\sqrt[3]{2}$ , and  $\sqrt{3} + \sqrt{2}$  are algebraic.
- Fix  $n \in \mathbf{N}$ , and let  $A_n$  be the algebraic numbers obtained as roots of polynomials with integer coefficients that have degree  $n$ . Using the fact that every polynomial has a finite number of roots, show that  $A_n$  is countable.
- Now, argue that the set of all algebraic numbers is countable. What may we conclude about the set of transcendental numbers?

### Solution

- $x^2 - 2 = 0$ ,  $x^3 - 2 = 0$  are obvious. Now consider  $\sqrt{3} + \sqrt{2}$ . The key is setting  $x = \sqrt{3} + \sqrt{2}$  then using algebra on  $x$  to concoct an integer, and thus find the polynomial with  $x$  as a root.

We have  $x^2 = 5 + 2\sqrt{6}$  meaning  $x^2 - 5 = 2\sqrt{6}$  and thus  $(x^2 - 5)^2 = 24$  so  $(x^2 - 5)^2 - 24 = 0$  is a polynomial with  $\sqrt{3} + \sqrt{2}$  as a root.

- Basically  $A_n \sim \mathbf{Z}^n \sim \mathbf{N}^n \sim \mathbf{N}$ .
  - $A_n \sim \mathbf{Z}^n$  since integer polynomials of degree  $n$  are identical to an ordered list of  $n$  integers.

- (ii)  $\mathbf{Z}^n \sim \mathbf{N}^n$  since  $f : \mathbf{N}^n \rightarrow \mathbf{Z}^n$  is just the piecewise application of  $g : \mathbf{N} \rightarrow \mathbf{Z}$ .
- (iii)  $\mathbf{N}^n \sim \mathbf{N}$  since it is the intersection of finite sets  $\bigcup_{n=2}^{\infty} \{(a, b) : a + b = n\}$ .

In general if  $V$  is countable then  $V^n = (v_1, \dots, v_n)$  is also countable.

- (c) By 1.5.3 the set of all algebraic numbers  $\bigcup_{n=1}^{\infty} A_n$  is countable.

### Exercise 1.5.10

- (a) Let  $C \subseteq [0, 1]$  be uncountable. Show that there exists  $a \in (0, 1)$  such that  $C \cap [a, 1]$  is uncountable.
- (b) Now let  $A$  be the set of all  $a \in (0, 1)$  such that  $C \cap [a, 1]$  is uncountable, and set  $\alpha = \sup A$ . Is  $C \cap [\alpha, 1]$  an uncountable set?
- (c) Does the statement in (a) remain true if “uncountable” is replaced by “infinite”?

### Solution

- (a) Suppose  $a$  does not exist, then  $C \cap [a, 1]$  is countable for all  $a \in (0, 1)$  meaning

$$\bigcup_{n=1}^{\infty} C \cap [1/n, 1] = C \cap [0, 1]$$

Is countable (by 1.5.3), contradicting our assumption that  $C \cap [0, 1]$  is uncountable.

- (b) If  $\alpha = 1$  then  $C \cap [\alpha, 1]$  is finite. Now if  $\alpha < 1$  we have  $C \cap [\alpha + \epsilon, 1]$  countable for  $\epsilon > 0$  (otherwise the set would be in  $A$ , and hence  $\alpha$  would not be an upper bound). Take

$$\bigcup_{n=1}^{\infty} C \cap [\alpha + 1/n, 1] = C \cap [\alpha, 1]$$

Which is countable by 1.5.3.

- (c) No, consider the set  $C = \{1/n : n \in \mathbf{N}\}$  it has  $C \cap [\alpha, 1]$  finite for every  $\alpha$ , but  $C \cap [0, 1]$  is infinite.

### Exercise 1.5.11 (Schröder-Bernstein Theorem)

Assume there exists a 1-1 function  $f : X \rightarrow Y$  and another 1-1 function  $g : Y \rightarrow X$ . Follow the steps to show that there exists a 1-1, onto function  $h : X \rightarrow Y$  and hence  $X \sim Y$ . The strategy is to partition  $X$  and  $Y$  into components

$$X = A \cup A' \quad \text{and} \quad Y = B \cup B'$$

with  $A \cap A' = \emptyset$  and  $B \cap B' = \emptyset$ , in such a way that  $f$  maps  $A$  onto  $B$ , and  $g$  maps  $B'$  onto  $A'$ .

- (a) Explain how achieving this would lead to a proof that  $X \sim Y$ .

- (b) Set  $A_1 = X \setminus g(Y) = \{x \in X : x \notin g(Y)\}$  (what happens if  $A_1 = \emptyset$ ?) and inductively define a sequence of sets by letting  $A_{n+1} = g(f(A_n))$ . Show that  $\{A_n : n \in \mathbf{N}\}$  is a pairwise disjoint collection of subsets of  $X$ , while  $\{f(A_n) : n \in \mathbf{N}\}$  is a similar collection in  $Y$ .
- (c) Let  $A = \bigcup_{n=1}^{\infty} A_n$  and  $B = \bigcup_{n=1}^{\infty} f(A_n)$ . Show that  $f$  maps  $A$  onto  $B$ .
- (d) Let  $A' = X \setminus A$  and  $B' = Y \setminus B$ . Show  $g$  maps  $B'$  onto  $A'$ .

**Solution**

- (a) **TODO**
- (b) **TODO**
- (c) **TODO**
- (d) **TODO**

## 1.6 Cantor's theorem

### Exercise 1.6.1

Show that  $(0, 1)$  is uncountable if and only if  $\mathbf{R}$  is uncountable.

### Solution

Exercise 1.5.4 tells us  $(0, 1)$  has the same cardinality as  $\mathbf{R}$ .

**TODO** Prove without using exercise 1.5.4 (probably what was intended)

### Exercise 1.6.2

Let  $f : \mathbf{N} \rightarrow \mathbf{R}$  be a way to list every real number (hence show  $\mathbf{R}$  is countable).

Define a new number  $x$  with digits  $b_1b_2 \dots$  given by

$$b_n = \begin{cases} 2 & \text{if } a_{nn} \neq 2 \\ 3 & \text{if } a_{nn} = 2 \end{cases}$$

- (a) Explain why the real number  $x = .b_1b_2b_3b_4 \dots$  cannot be  $f(1)$ .
- (b) Now, explain why  $x \neq f(2)$ , and in general why  $x \neq f(n)$  for any  $n \in \mathbf{N}$ .
- (c) Point out the contradiction that arises from these observations and conclude that  $(0, 1)$  is uncountable.

### Solution

- (a) The first digit is different
- (b) The  $n$ th digit is different
- (c) Therefore  $x$  is not in the list, since the  $n$ th digit is different

### Exercise 1.6.3

Supply rebuttals to the following complaints about the proof of Theorem 1.6.1.

- (a) Every rational number has a decimal expansion, so we could apply this same argument to show that the set of rational numbers between 0 and 1 is uncountable. However, because we know that any subset of  $\mathbf{Q}$  must be countable, the proof of Theorem 1.6.1 must be flawed.
- (b) Some numbers have two different decimal representations. Specifically, any decimal expansion that terminates can also be written with repeating 9's. For instance,  $1/2$  can be written as  $.5$  or as  $.4999 \dots$ . Doesn't this cause some problems?

### Solution

- (a) False, since the constructed number has an infinite number of decimals it is irrational.
- (b) No, since if we have  $9999 \dots$  and change the  $n$ th digit  $9992999 = 9993$  is still different.

**Exercise 1.6.4**

Let  $S$  be the set consisting of all sequences of 0's and 1's. Observe that  $S$  is not a particular sequence, but rather a large set whose elements are sequences; namely,

$$S = \{(a_1, a_2, a_3, \dots) : a_n = 0 \text{ or } 1\}$$

As an example, the sequence  $(1, 0, 1, 0, 1, 0, 1, 0, \dots)$  is an element of  $S$ , as is the sequence  $(1, 1, 1, 1, 1, 1, \dots)$ . Give a rigorous argument showing that  $S$  is uncountable.

**Solution**

We flip every bit in the diagonal just like with  $\mathbf{R}$ . Another way would be to show  $S \sim \mathbf{R}$  by writing real numbers in base 2.

**Exercise 1.6.5**

- (a) Let  $A = \{a, b, c\}$ . List the eight elements of  $P(A)$ . (Do not forget that  $\emptyset$  is considered to be a subset of every set.)
- (b) If  $A$  is finite with  $n$  elements, show that  $P(A)$  has  $2^n$  elements.

**Solution**

- (a)  $A = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}$ .
- (b) There are  $n$  elements, we can include or exclude each element so there are  $2^n$  subsets.

**Exercise 1.6.6**

- (a) Using the particular set  $A = \{a, b, c\}$ , exhibit two different 1 – 1 mappings from  $A$  into  $P(A)$ .
- (b) Letting  $C = \{1, 2, 3, 4\}$ , produce an example of a 1 – 1 map  $g : C \rightarrow P(C)$ .
- (c) Explain why, in parts (a) and (b), it is impossible to construct mappings that are onto.

**Solution**

- (a)  $f(x) = \{x\}$ ,  $f(x) = \{x, b\}$  for  $x \neq b$  and  $f(x) = \{a, b, c\}$  for  $x = b$ .
- (b)  $f(x) = \{x\}$ .
- (c) We can hit at most  $n$  elements in the power set out of the  $2^n$  total elements.

**Theorem (Cantor's Theorem)**

Given any set  $A$ , there does not exist a function  $f : A \rightarrow P(A)$  that is onto.

**Proof**

Suppose  $f : A \rightarrow P(A)$  is onto. We want to use the self referential nature of  $P(A)$  to find a contradiction. Define

$$B = \{a : a \notin f(a)\}$$

Since  $f$  is onto we must have  $f(a) = B$  for some  $a \in A$ . Then either

- (i)  $a \in B$  implies  $a \in f(a)$  which by the definition of  $B$  implies  $a \notin B$ , so  $a \in B$  is impossible.

- (ii)  $a \notin B$  implies  $a \notin f(a)$  since  $f(a) = B$ . but if  $a \notin f(a)$  then  $a \in B$  by the definition of  $B$ , contradicting  $a \notin B$ .

Therefore  $f$  cannot be onto, since we have found a  $B \in P(A)$  where  $f(a) = B$  is impossible.

Stepping back, the pearl of the argument is that if  $B = f(a)$  then  $B = \{a : a \notin B\}$  is undecidable/impossible.

### Exercise 1.6.7

See the proof of Cantor's theorem above (the rest is a computation)

### Exercise 1.6.8

See the proof of Cantor's theorem above

### Exercise 1.6.9

Using the various tools and techniques developed in the last two sections (including the exercises from Section 1.5), give a compelling argument showing that  $P(\mathbf{N}) \sim \mathbf{R}$ .

### Solution

I will show  $P(\mathbf{N}) \sim [0, 1]$  then use 1.5.3 to conclude  $P(\mathbf{N}) \sim \mathbf{R}$ .

Let  $A \subseteq \mathbf{N}$  and let  $a_n$  be the  $n$ th smallest element of  $A$ . We can write  $a_n$  via the digit representation as  $a_n = d_1 d_2 d_3 \dots d_m$ , concatenating the digits of every  $a_n$  in order gives a possibly infinite sequence of digits  $d_1 d_2 d_3 \dots$ .

This process is clearly 1-1, however it is not onto as  $\{1, 2\}$  and  $\{12\}$  both give the same digits. Thus  $P(\mathbf{N})$  is "greater than or equal"  $\mathbf{R}$ , if we show a 1-1 map  $\mathbf{R} \rightarrow P(\mathbf{N})$  we can complete the proof using 1.5.11.

**TODO** Finish (or take a different approach)

### Exercise 1.6.10

As a final exercise, answer each of the following by establishing a 1 – 1 correspondence with a set of known cardinality.

- (a) Is the set of all functions from  $\{0, 1\}$  to  $\mathbf{N}$  countable or uncountable?
- (b) Is the set of all functions from  $\mathbf{N}$  to  $\{0, 1\}$  countable or uncountable?
- (c) Given a set  $B$ , a subset  $\mathcal{A}$  of  $P(B)$  is called an antichain if no element of  $\mathcal{A}$  is a subset of any other element of  $\mathcal{A}$ . Does  $P(\mathbf{N})$  contain an uncountable antichain?

### Solution

- (a) The set of functions from  $\{0, 1\}$  to  $\mathbf{N}$  is the same as  $\mathbf{N}^2$  which we found was countable in Exercise 1.5.9.
- (b) This is the same as an infinite list of zeros and ones which we showed was uncountable in Exercise 1.6.4.
- (c) **TODO**





# Chapter 2

## Sequences and Series

### 2.2 The Limit of a Sequence

#### Exercise 2.2.1

What happens if we reverse the order of the quantifiers in Definition 2.2.3?

Definition: A sequence  $(x_n)$  *verconges* to  $x$  if *there exists* an  $\epsilon > 0$  such that *for all*  $N \in \mathbf{N}$  it is true that  $n \geq N$  implies  $|x_n - x| < \epsilon$

Give an example of a vercongent sequence. Is there an example of a vercongent sequence that is divergent? Can a sequence verconge to two different values? What exactly is being described in this strange definition?

#### Solution

Firstly, since we have *for all*  $N \in \mathbf{N}$  we can remove  $N$  entirely and just say  $n \in \mathbf{N}$ . Our new definition is

Definition: A sequence  $(x_n)$  *verconges* to  $x$  if *there exists* an  $\epsilon > 0$  such that *for all*  $n \in \mathbf{N}$  we have  $|x_n - x| < \epsilon$ .

In other words, a series  $(x_n)$  *verconges* to  $x$  if  $|x_n - x|$  is bounded. This is a silly definition though since if  $|x_n - x|$  is bounded, then  $|x_n - x'|$  is bounded for all  $x' \in \mathbf{R}$ , meaning if a sequence is vercongent it verconges to every  $x' \in \mathbf{R}$ .

Put another way, a sequence is vercongent *if and only if* it is bounded.

#### Exercise 2.2.2

Verify, using the definition of convergence of a sequence, that the following sequences converge to the proposed limit.

(a)  $\lim_{n \rightarrow \infty} \frac{2n+1}{5n+4} = \frac{2}{5}$ .

(b)  $\lim_{n \rightarrow \infty} \frac{2n^2}{n^3+3} = 0$ .

(c)  $\lim_{n \rightarrow \infty} \frac{\sin(n^2)}{\sqrt[3]{n}} = 0$ .

#### Solution

(a) We have

$$\left| \frac{2n+1}{5n+4} - \frac{2}{5} \right| = \left| \frac{5(2n+1) - 2(5n+4)}{5(5n+4)} \right| = \left| \frac{-3}{5(5n+4)} \right| = \frac{3}{5(5n+4)} < \epsilon$$

We now find  $n$  such that the distance is less than  $\epsilon$

$$\frac{3}{5(5n+4)} < \frac{1}{n} < \epsilon \implies n > \frac{1}{\epsilon}$$

You could also solve for the smallest  $n$ , which would give you

$$\frac{3}{5(5n+4)} < \epsilon \implies 5n+4 > \frac{3}{5\epsilon} \implies n > \frac{3}{25\epsilon} - \frac{4}{5}$$

I prefer the first approach, the second is better if you were doing numerical analysis and wanted a precise convergence rate.

(b) We have

$$\left| \frac{2n^2}{n^3+3} - 0 \right| = \frac{2n^2}{n^3+3} < \frac{2n^2}{n^3} = \frac{2}{n} < \epsilon \implies n > \frac{2}{\epsilon}$$

(c) We have

$$\frac{\sin(n^2)}{n^{1/3}} \leq \frac{1}{n^{1/3}} < \epsilon \implies n > \frac{1}{\epsilon^3}$$

Really slow convergence! if  $\epsilon = 10^{-2}$  we would require  $n > 10^6$

### Exercise 2.2.3

Describe what we would have to demonstrate in order to disprove each of the following statements.

- (a) At every college in the United States, there is a student who is at least seven feet tall.
- (b) For all colleges in the United States, there exists a professor who gives every student a grade of either A or B.
- (c) There exists a college in the United States where every student is at least six feet tall.

### Solution

- (a) Find a college in the United States with no students over seven feet tall.
- (b) Find a college in the United States with a professor who has given a grade other than an A or B.
- (c) Find a college in the United States with at least one student under six feet tall.

### Exercise 2.2.4

Give an example of each or state that the request is impossible. For any that are impossible, give a compelling argument for why that is the case.

- (a) A sequence with an infinite number of ones that does not converge to one.
- (b) A sequence with an infinite number of ones that converges to a limit not equal to one.
- (c) A divergent sequence such that for every  $n \in \mathbf{N}$  it is possible to find  $n$  consecutive ones somewhere in the sequence.

**Solution**

- (a)  $a_n = (-1)^n$
- (b) Impossible, if  $\lim a_n = a \neq 1$  then for any  $n \geq N$  we can find a  $n$  with  $a_n = 1$  meaning  $\epsilon < |1 - a|$  is impossible.
- (c)  $a_n = (1, 2, 1, 1, 3, 1, 1, 1, \dots)$

**Exercise 2.2.5**

Let  $\llbracket x \rrbracket$  be the greatest integer less than or equal to  $x$ . For example,  $\llbracket \pi \rrbracket = 3$  and  $\llbracket 3 \rrbracket = 3$ . For each sequence, find  $\lim a_n$  and verify it with the definition of convergence.

- (a)  $a_n = \llbracket 5/n \rrbracket$ ,
- (b)  $a_n = \llbracket (12 + 4n)/3n \rrbracket$ .

Reflecting on these examples, comment on the statement following Definition 2.2.3 that “the smaller the  $\epsilon$ -neighborhood, the larger  $N$  may have to be.”

**Solution**

- (a) For all  $n > 5$  we have  $\llbracket 5/n \rrbracket = 0$  meaning  $\lim a_n = 0$ .
- (b) The inside clearly converges to  $4/3$  from above, so  $\lim a_n = 1$ .  
Some sequences eventually reach their limit, meaning  $N$  no longer has to increase.

**Exercise 2.2.6**

**Theorem 2.2.7 (Uniqueness of Limits).** *The limit of a sequence, when it exists, must be unique.*

Prove Theorem 2.2.7. To get started, assume  $(a_n) \rightarrow a$  and also that  $(a_n) \rightarrow b$ . Now argue  $a = b$

**Solution**

If  $a \neq b$  then we can set  $\epsilon$  small enough that having both  $|a_n - a| < \epsilon$  and  $|a_n - b| < \epsilon$  is impossible. Therefore  $a = b$ .

(Making this rigorous is trivial and left as an exercise to the reader)

**Exercise 2.2.7**

Here are two useful definitions:

- (i) A sequence  $(a_n)$  is *eventually* in a set  $A \subseteq \mathbf{R}$  if there exists an  $N \in \mathbf{N}$  such that  $a_n \in A$  for all  $n \geq N$ .
- (ii) A sequence  $(a_n)$  is *frequently* in a set  $A \subseteq \mathbf{R}$  if, for every  $N \in \mathbf{N}$ , there exists an  $n \geq N$  such that  $a_n \in A$ .
  - (a) Is the sequence  $(-1)^n$  eventually or frequently in the set  $\{1\}$ ?
  - (b) Which definition is stronger? Does frequently imply eventually or does eventually imply frequently?
  - (c) Give an alternate rephrasing of Definition 2.2.3B using either frequently or eventually. Which is the term we want?

- (d) Suppose an infinite number of terms of a sequence  $(x_n)$  are equal to 2. Is  $(x_n)$  necessarily eventually in the interval  $(1.9, 2.1)$ ? Is it frequently in  $(1.9, 2.1)$ ?

**Solution**

- (a) Frequently, but not eventually.
- (b) Eventually is stronger, it implies frequently.
- (c)  $(x_n) \rightarrow x$  if and only if  $x_n$  is eventually in any  $\epsilon$ -neighborhood around  $x$ .
- (d)  $(x_n)$  is frequently in  $(1.9, 2.1)$  but not necessarily eventually (consider  $x_n = 2(-1)^n$ ).

**Exercise 2.2.8**

For some additional practice with nested quantifiers, consider the following invented definition:

Let's call a sequence  $(x_n)$  zero-heavy if there exists  $M \in \mathbf{N}$  such that for all  $N \in \mathbf{N}$  there exists  $n$  satisfying  $N \leq n \leq N + M$  where  $x_n = 0$

- (a) Is the sequence  $(0, 1, 0, 1, 0, 1, \dots)$  zero heavy?
- (b) If a sequence is zero-heavy does it necessarily contain an infinite number of zeros? If not, provide a counterexample.
- (c) If a sequence contains an infinite number of zeros, is it necessarily zeroheavy? If not, provide a counterexample.
- (d) Form the logical negation of the above definition. That is, complete the sentence: A sequence is not zero-heavy if ....

**Solution**

- (a) No.
- (b) Yes. as any finite number of zeros  $K$  would lead to a contradiction when  $M > K$ .
- (c) No, consider  $(0, 1, 0, \dots)$  from (a).
- (d) A sequence is not zero-heavy if there exists an  $M \in \mathbf{N}$  such that for all  $N \in \mathbf{N}$  there exists an  $n \in \mathbf{N}$  such that  $N \leq n \leq N + M$  but  $x_n \neq 0$ .

## 2.3 The Algebraic and Order Limit Theorems

### Exercise 2.3.1

Let  $x_n \geq 0$  for all  $n \in \mathbf{N}$ .

- (a) If  $(x_n) \rightarrow 0$ , show that  $(\sqrt{x_n}) \rightarrow 0$ .
- (b) If  $(x_n) \rightarrow x$ , show that  $(\sqrt{x_n}) \rightarrow \sqrt{x}$ .

### Solution

- (a) Setting  $x_n < \epsilon^2$  implies  $\sqrt{x_n} < \epsilon$  (for all  $n \geq N$  of course)
- (b) We want  $|\sqrt{x_n} - \sqrt{x}| < \epsilon$  multiplying by  $(\sqrt{x_n} + \sqrt{x})$  gives  $|x_n - x| < (\sqrt{x_n} + \sqrt{x})\epsilon$  since  $x_n$  is convergent, it is bounded  $|x_n| \leq M$  implying  $\sqrt{|x_n|} \leq \sqrt{M}$ , multiplying gives

$$|x_n - x| < (\sqrt{x_n} + \sqrt{x})\epsilon \leq (\sqrt{M} + \sqrt{x})\epsilon$$

Since  $|x_n - x|$  can be made arbitrarily small we can make this true for some  $n \geq N$ . Now dividing by  $\sqrt{M} + \sqrt{x}$  gives us

$$|\sqrt{x_n} - \sqrt{x}| \leq \frac{|x_n - x|}{\sqrt{M} + \sqrt{x}} < \epsilon$$

Therefor  $|\sqrt{x_n} - \sqrt{x}| < \epsilon$  completing the proof.

### Exercise 2.3.2

Using only Definition 2.2.3, prove that if  $(x_n) \rightarrow 2$ , then

- (a)  $(\frac{2x_n-1}{3}) \rightarrow 1$ ;
- (b)  $(1/x_n) \rightarrow 1/2$ .

(For this exercise the Algebraic Limit Theorem is off-limits, so to speak.)

### Solution

- (a) We have  $|\frac{2}{3}x_n - \frac{4}{3}| = \frac{2}{3}|x_n - 2| < \epsilon$  which can always be done since  $|x_n - 2|$  can be made arbitrarily small.
- (b) Since  $x_n$  is bounded we have  $|x_n| \leq M$

$$|1/x_n - 1/2| = \frac{|2 - x_n|}{|2x_n|} \leq \frac{|2 - x_n|}{|2M|} < \epsilon$$

Letting  $|2 - x_n| < \epsilon/|2M|$  gives  $|1/x_n - 1/2| < \epsilon$ .

### Exercise 2.3.3 (Squeeze Theorem)

Show that if  $x_n \leq y_n \leq z_n$  for all  $n \in \mathbf{N}$ , and if  $\lim x_n = \lim z_n = l$ , then  $\lim y_n = l$  as well.

### Solution

Let  $y = \lim y_n$ . By the order limit theorem we have  $l \leq y \leq l$  implying  $y = l$ .

**Exercise 2.3.4**

Let  $(a_n) \rightarrow 0$ , and use the Algebraic Limit Theorem to compute each of the following limits (assuming the fractions are always defined):

(a)  $\lim \left( \frac{1+2a_n}{1+3a_n-4a_n^2} \right)$

(b)  $\lim \left( \frac{(a_n+2)^2-4}{a_n} \right)$

(c)  $\lim \left( \frac{\frac{2}{a_n}+3}{\frac{1}{a_n}+5} \right).$

**Solution**

(a) Divide by  $a_n^2$ , then apply the ALT

$$\lim \left( \frac{1+2a_n}{1+3a_n-4a_n^2} \right) = \frac{\lim \left( \frac{1}{a_n^2} + \frac{2}{a_n} \right)}{\lim \left( \frac{1}{a_n^2} + \frac{3}{a_n} - 4 \right)} = \frac{0}{-4} = 0$$

(b)

$$\lim \left( \frac{(a_n+2)^2-4}{a_n} \right) = \lim \left( \frac{a_n^2+2a_n}{a_n} \right) = \lim (a_n+2) = \infty$$

(c) This one is a straightforward application of the algebraic limit theorem

$$\lim \left( \frac{\frac{2}{a_n}+3}{\frac{1}{a_n}+5} \right) = 3/5$$

**Exercise 2.3.5**

Let  $(x_n)$  and  $(y_n)$  be given, and define  $(z_n)$  to be the “shuffled” sequence  $(x_1, y_1, x_2, y_2, x_3, y_3, \dots, x_n, y_n, \dots)$ . Prove that  $(z_n)$  is convergent if and only if  $(x_n)$  and  $(y_n)$  are both convergent with  $\lim x_n = \lim y_n$ .

**Solution**

Obviously if  $\lim x_n = \lim y_n = l$  then  $z_n \rightarrow l$ . To show the other way suppose  $(z_n) \rightarrow l$ , then  $|z_n - l| < \epsilon$  for all  $n \geq N$  meaning  $|y_n - l| < \epsilon$  and  $|x_n - l| < \epsilon$  for  $n \geq N$  as well. Thus  $\lim x_n = \lim y_n = l$ .

**Exercise 2.3.6**

Consider the sequence given by  $b_n = n - \sqrt{n^2 + 2n}$ . Taking  $(1/n) \rightarrow 0$  as given, and using both the Algebraic Limit Theorem and the result in Exercise 2.3.1, show  $\lim b_n$  exists and find the value of the limit.

**Solution**

I’m going to find the value of the limit before proving it. We have

$$n - \sqrt{n^2 + 2n} = n - \sqrt{(n+1)^2 - 1}$$

For large  $n$ ,  $\sqrt{(n+1)^2 - 1} \approx n + 1$  so  $\lim b_n = -1$ .

Factoring out  $n$  we get  $n \left(1 - \sqrt{1 + 2/n}\right)$ . Tempting as it is to apply the ALT here to say  $(b_n) \rightarrow 0$  it doesn't work since  $n$  diverges.

How about if I get rid of the radical, then use the ALT to go back to what we had before?

$$(n - \sqrt{n^2 + 2n})(n + \sqrt{n^2 + 2n}) = n^2 - (n^2 + 2n) = -2n$$

Then we have

$$b_n = n - \sqrt{n^2 + 2n} = \frac{-2n}{n + \sqrt{n^2 + 2n}} = \frac{-2}{1 + \sqrt{1 + 2/n}}$$

Now we can finally use the algebraic limit theorem!

$$\lim \left( \frac{-2}{1 + \sqrt{1 + 2/n}} \right) = \frac{-2}{1 + \sqrt{1 + \lim (2/n)}} = \frac{-2}{1 + \sqrt{1 + 0}} = -1$$

Stepping back the key to this technique is removing the radicals via a difference of squares, then dividing both sides by the growthrate  $n$  and applying the ALT.

### Exercise 2.3.7

Give an example of each of the following, or state that such a request is impossible by referencing the proper theorem(s):

- (a) sequences  $(x_n)$  and  $(y_n)$ , which both diverge, but whose sum  $(x_n + y_n)$  converges;
- (b) sequences  $(x_n)$  and  $(y_n)$ , where  $(x_n)$  converges,  $(y_n)$  diverges, and  $(x_n + y_n)$  converges;
- (c) a convergent sequence  $(b_n)$  with  $b_n \neq 0$  for all  $n$  such that  $(1/b_n)$  diverges;
- (d) an unbounded sequence  $(a_n)$  and a convergent sequence  $(b_n)$  with  $(a_n - b_n)$  bounded;
- (e) two sequences  $(a_n)$  and  $(b_n)$ , where  $(a_n b_n)$  and  $(a_n)$  converge but  $(b_n)$  does not.

### Solution

- (a)  $(x_n) = n$  and  $(y_n) = -n$  diverge but  $x_n + y_n = 0$  converges
- (b) Impossible, the algebraic limit theorem implies  $\lim(x_n + y_n) - \lim(x_n) = \lim y_n$  therefore  $(y_n)$  must converge if  $(x_n)$  and  $(x_n + y_n)$  converge.
- (c) Impossible, the algebraic limit theorem implies that if  $(b_n) \rightarrow b$  then  $(1/b_n) \rightarrow 1/b$ .
- (d) Impossible, letting  $|b_n| \leq M$  we have  $|a_n - b_n| \leq |a_n - M|$  being bounded, which is impossible since  $a_n$  is unbounded and shifting by a constant  $M$  cannot not change that.
- (e)  $b_n = n$  and  $a_n = 0$  works. However if  $(a_n) \rightarrow a$ ,  $a \neq 0$  and  $(a_n b_n) \rightarrow p$  then the ALT would imply  $(b_n) \rightarrow p/a$ .

### Exercise 2.3.8

Let  $(x_n) \rightarrow x$  and let  $p(x)$  be a polynomial.

- (a) Show  $p(x_n) \rightarrow p(x)$ .
- (b) Find an example of a function  $f(x)$  and a convergent sequence  $(x_n) \rightarrow x$  where the sequence  $f(x_n)$  converges, but not to  $f(x)$ .

**Solution**

- (a) Applying the algebraic limit theorem multiple times gives  $(x_n^d) \rightarrow x^d$  meaning

$$\lim p(x_n) = \lim (a_d x_n^d + a_{d-1} x_n^{d-1} + \cdots + a_0) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_0 = p(x).$$

As a cute corollary, any continuous function  $f$  has  $\lim f(x_n) = f(x)$  since polynomials can approximate continuous functions arbitrarily well by the Weierstrass approximation theorem.

- (b) Let  $(x_n) = 1/n$  and define  $f$  as

$$f(x) = \begin{cases} 0 & \text{if } x = 0 \\ 1 & \text{otherwise} \end{cases}$$

We have  $f(1/n) = 1$  for all  $n$ , meaning  $\lim f(1/n) = 1$  but  $f(0) = 0$ .

**Exercise 2.3.9**

- (a) Let  $(a_n)$  be a bounded (not necessarily convergent) sequence, and assume  $\lim b_n = 0$ . Show that  $\lim (a_n b_n) = 0$ . Why are we not allowed to use the Algebraic Limit Theorem to prove this?
- (b) Can we conclude anything about the convergence of  $(a_n b_n)$  if we assume that  $(b_n)$  converges to some nonzero limit  $b$ ?
- (c) Use (a) to prove Theorem 2.3.3, part (iii), for the case when  $a = 0$ .

**Solution**

- (a) We can't use the ALT since  $a_n$  is not necessarily convergent.  $a_n$  being bounded gives  $|a_n| \leq M$  for some  $M$  giving

$$|a_n b_n| \leq M |b_n| < \epsilon$$

Which can be accomplished by letting  $|b_n| < \epsilon/M$  since  $(b_n) \rightarrow 0$ .

- (b) No
- (c) In (a) we showed  $\lim(a_n b_n) = 0 = ab$  for  $b = 0$  which proves part (iii) of the ALT.

**Exercise 2.3.10**

Consider the following list of conjectures. Provide a short proof for those that are true and a counterexample for any that are false.

- (a) If  $\lim (a_n - b_n) = 0$ , then  $\lim a_n = \lim b_n$ .
- (b) If  $(b_n) \rightarrow b$ , then  $|b_n| \rightarrow |b|$ .



- (c) If  $(a_n) \rightarrow a$  and  $(b_n - a_n) \rightarrow 0$ , then  $(b_n) \rightarrow a$ .
- (d) If  $(a_n) \rightarrow 0$  and  $|b_n - b| \leq a_n$  for all  $n \in \mathbf{N}$ , then  $(b_n) \rightarrow b$ .

**Solution**

- (a) False, consider  $a_n = n$  and  $b_n = -n$ .
- (b) True since if  $|b_n - b| < \epsilon$  then  $||b_n| - |b|| \leq |b_n - b| < \epsilon$  by Exercise 1.2.6 (d).
- (c) True by ALT since  $\lim(b_n - a_n) + \lim a_n = \lim b_n = a$ .
- (d) True, since  $0 \leq |b_n - b| \leq a_n$  we have  $a_n \geq 0$ . Let  $\epsilon > 0$  and pick  $N$  such that  $a_n < \epsilon$  for all  $n \geq N$ . Therefor

$$|b_n - b| \leq a_n < \epsilon$$

Proving  $(b_n) \rightarrow b$ .

**Exercise 2.3.11 (Cesaro Means)**

- (a) Show that if  $(x_n)$  is a convergent sequence, then the sequence given by the averages

$$y_n = \frac{x_1 + x_2 + \cdots + x_n}{n}$$

also converges to the same limit.

- (b) Give an example to show that it is possible for the sequence  $(y_n)$  of averages to converge even if  $(x_n)$  does not.

**Solution**

- (a) Let  $D = \sup\{|x_n - x| : n \in \mathbf{N}\}$  and let  $0 < \epsilon < D$ , we have

$$|y_n - x| = \left| \frac{x_1 + \cdots + x_n}{n} - x \right| \leq \left| \frac{|x_1 - x| + \cdots + |x_n - x|}{n} \right| \leq D$$

Let  $|x_n - x| < \epsilon/2$  for  $n > N_1$  giving

$$|y_n - x| \leq \left| \frac{|x_1 - x| + \cdots + |x_{N_1} - x| + \cdots + |x_n - x|}{n} \right| \leq \left| \frac{N_1 D + (n - N_1)\epsilon/2}{n} \right|$$

Let  $N_2$  be large enough that for all  $n > N_2$  (remember  $0 < \epsilon < D$  so  $(D - \epsilon/2) > 0$ .)

$$0 < \frac{N_1(D - \epsilon/2)}{n} < \epsilon/2$$

Therefor

$$|y_n - x| \leq \left| \frac{N_1(D - \epsilon/2)}{n} + \epsilon/2 \right| < \epsilon$$

Letting  $N = \max\{N_1, N_2\}$  completes the proof as  $|y_n - x| < \epsilon$  for all  $n > N$ .

(Note: I could have used any  $\epsilon' < \epsilon$  instead of  $\epsilon/2$ , I just needed some room.)

- (b)  $x_n = (-1)^n$  diverges but  $(y_n) \rightarrow 0$ .

### Exercise 2.3.12

A typical task in analysis is to decipher whether a property possessed by every term in a convergent sequence is necessarily inherited by the limit. Assume  $(a_n) \rightarrow a$ , and determine the validity of each claim. Try to produce a counterexample for any that are false.

- (a) If every  $a_n$  is an upper bound for a set  $B$ , then  $a$  is also an upper bound for  $B$ .
- (b) If every  $a_n$  is in the complement of the interval  $(0, 1)$ , then  $a$  is also in the complement of  $(0, 1)$ .
- (c) If every  $a_n$  is rational, then  $a$  is rational.

### Solution

- (a) True, let  $s = \sup B$ , we know  $s \leq a_n$  so by the order limit theorem  $s \leq a$  meaning  $a$  is also an upper bound on  $B$ .
- (b) True, since if  $a \in (0, 1)$  then there would exist an  $\epsilon$ -neighborhood inside  $(0, 1)$  that  $a_n$  would have to fall in, contradicting the fact that  $a_n \notin (0, 1)$ .
- (c) False, consider the sequence of rational approximations to  $\sqrt{2}$

### Exercise 2.3.13 (Iterated Limits)

Given a doubly indexed array  $a_{mn}$  where  $m, n \in \mathbf{N}$ , what should  $\lim_{m,n \rightarrow \infty} a_{mn}$  represent?

- (a) Let  $a_{mn} = m/(m+n)$  and compute the iterated limits

$$\lim_{n \rightarrow \infty} \left( \lim_{m \rightarrow \infty} a_{mn} \right) \quad \text{and} \quad \lim_{m \rightarrow \infty} \left( \lim_{n \rightarrow \infty} a_{mn} \right)$$

Define  $\lim_{m,n \rightarrow \infty} a_{mn} = a$  to mean that for all  $\epsilon > 0$  there exists an  $N \in \mathbf{N}$  such that if both  $m, n \geq N$ , then  $|a_{mn} - a| < \epsilon$

- (b) Let  $a_{mn} = 1/(m+n)$ . Does  $\lim_{m,n \rightarrow \infty} a_{mn}$  exist in this case? Do the two iterated limits exist? How do these three values compare? Answer these same questions for  $a_{mn} = mn/(m^2 + n^2)$
- (c) Produce an example where  $\lim_{m,n \rightarrow \infty} a_{mn}$  exists but where neither iterated limit can be computed.
- (d) Assume  $\lim_{m,n \rightarrow \infty} a_{mn} = a$ , and assume that for each fixed  $m \in \mathbf{N}$ ,  $\lim_{n \rightarrow \infty} (a_{mn}) \rightarrow b_m$ . Show  $\lim_{m \rightarrow \infty} b_m = a$
- (e) Prove that if  $\lim_{m,n \rightarrow \infty} a_{mn}$  exists and the iterated limits both exist, then all three limits must be equal.

### Solution

- (a)

$$\lim_{n \rightarrow \infty} \left( \lim_{m \rightarrow \infty} \frac{m}{m+n} \right) = 1, \quad \text{and} \quad \lim_{m \rightarrow \infty} \left( \lim_{n \rightarrow \infty} \frac{m}{m+n} \right) = 0$$

- (b) For  $a_{mn} = 1/(m+n)$  all three limits are zero. For  $a_{mn} = mn/(m^2 + n^2)$  iterated limits are zero, and  $\lim_{m,n \rightarrow \infty} a_{mn}$  does not exist since for  $m, n \geq N$  setting  $m = n$  gives

$$\frac{n^2}{n^2 + n^2} = \frac{1}{2}$$

Which cannot be made smaller then  $\epsilon = 1/2$ .

The reason you would think to set  $m = n$  is in trying to maximize  $mn/(m^2 + n^2)$  notice if  $m > n$  then  $mn > n^2$  so we are adding more to the numerator then the denominator, hence the ratio is increasing. And if  $m < n$  then the ratio is decreasing. Therefor the maximum point is at  $m = n$ .

- (c) **TODO**

- (d) We are given  $|b_m - a_{mn}| < \epsilon$  for all  $n > N$ , By the triangle inequality

$$|b_m - a| \leq |b_m - a_{mn}| + |a_{mn} - a| < \epsilon/2 + \epsilon/2 = \epsilon$$

- (e) Let  $b_m = \lim_{n \rightarrow \infty} (a_{mn})$  and  $a = \lim_{m,n \rightarrow \infty} (a_{mn})$ . In (d) we showed  $(b_m) \rightarrow a$ , A similar argument shows  $(c_n) \rightarrow a$ . Thus all three limits are equal to  $a$ .

## 2.4 The Monotone Convergence Theorem and a First Look at Infinite Series

### Exercise 2.4.1

- (a) Prove that the sequence defined by  $x_1 = 3$  and

$$x_{n+1} = \frac{1}{4 - x_n}$$

converges.

- (b) Now that we know  $\lim x_n$  exists, explain why  $\lim x_{n+1}$  must also exist and equal the same value.
- (c) Take the limit of each side of the recursive equation in part (a) to explicitly compute  $\lim x_n$ .

### Solution

- (a)  $x_2 = 1$  makes me conjecture  $x_n$  is monotonic. For induction suppose  $x_n > x_{n+1}$  then we have

$$4 - x_n < 4 - x_{n+1} \implies \frac{1}{4 - x_n} > \frac{1}{4 - x_{n+1}} \implies x_{n+1} > x_{n+2}$$

Thus  $x_n$  is decreasing, to show  $x_n$  is bounded notice  $x_n$  cannot be negative since  $x_n < 3$  means  $x_{n+1} = 1/(4 - x_n) > 0$ . Therefore by the monotone convergence theorem  $(x_n)$  converges.

- (b) Clearly skipping a single term does not change what the series converges to.
- (c) Since  $x = \lim(x_n) = \lim(x_{n+1})$  we must have

$$x = \frac{1}{4 - x} \iff x^2 - 4x + 1 = 0 \iff (x - 2)^2 = 3 \iff x = 2 \pm \sqrt{3}$$

$2 + \sqrt{3} > 3$  is impossible since  $x_n < 3$  thus  $x = 2 - \sqrt{3}$ .

### Exercise 2.4.2

- (a) Consider the recursively defined sequence  $y_1 = 1$

$$y_{n+1} = 3 - y_n$$

and set  $y = \lim y_n$ . Because  $(y_n)$  and  $(y_{n+1})$  have the same limit, taking the limit across the recursive equation gives  $y = 3 - y$ . Solving for  $y$ , we conclude  $\lim y_n = 3/2$ . What is wrong with this argument?

- (b) This time set  $y_1 = 1$  and  $y_{n+1} = 3 - \frac{1}{y_n}$ . Can the strategy in (a) be applied to compute the limit of this sequence?

### Solution

- (a) The series  $y_n = (1, 2, 1, 2, \dots)$  does not converge.

- (b) Yes,  $y_n$  converges by the monotone convergence theorem since  $0 < y_n < 3$  and  $y_n$  is increasing.

### Exercise 2.4.3

- (a) Show that

$$\sqrt{2}, \sqrt{2 + \sqrt{2}}, \sqrt{2 + \sqrt{2 + \sqrt{2}}}, \dots$$

converges and find the limit.

- (b) Does the sequence

$$\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \dots$$

converge? If so, find the limit.

### Solution

- (a) Let  $x_1 = \sqrt{2}$  and  $x_{n+1} = \sqrt{2 + x_n}$  clearly  $x_2 > x_1$ . assuming  $x_{n+1} > x_n$  gives

$$2 + x_{n+1} > 2 + x_n \iff \sqrt{2 + x_{n+1}} > \sqrt{2 + x_n} \iff x_{n+2} > x_{n+1}$$

Since  $x_n$  is monotonically increasing and bounded the monotone convergence theorem tells us  $(x_n) \rightarrow x$ . Equating both sides like in 2.4.1 gives

$$x = \sqrt{2 + x} \iff x^2 - x - 2 = 0 \iff x = \frac{1}{2} \pm \frac{\sqrt{5}}{2}$$

Since  $x > 0$  we must have  $x = (1 + \sqrt{5})/2$ .

- (b) We have  $x_1 = 2^{1/2}$  and  $x_{n+1} = (2x_n)^{1/2}$ . We have

$$x_{n+1} = (2x_n)^{1/2} \geq x_n \iff 2x_n \geq x_n^2 \iff 2 \geq x_n$$

Since  $x_1 = 2^{1/2} \leq 2$  induction implies  $x_n$  is increasing. Now to show  $x_n$  is bounded notice that  $x_1 \leq 2$  and if  $x_n \leq 2$  then

$$2x_n \leq 4 \implies (2x_n)^{1/2} \leq 2$$

Now the monotone convergence theorem tells us  $(x_n)$  converges. To find the limit use  $\lim x_n = \lim x_{n+1} = x$  to get

$$x = (2x)^{1/2} \implies x^2 = 2x \implies x = \pm 2$$

Since  $x_n \geq 0$  we have  $x = 2$ .

### Exercise 2.4.4

- (a) In Section 1.4 we used the Axiom of Completeness (AoC) to prove the Archimedean Property of  $\mathbf{R}$  (Theorem 1.4.2). Show that the Monotone Convergence Theorem can also be used to prove the Archimedean Property without making any use of AoC.

- (b) Use the Monotone Convergence Theorem to supply a proof for the Nested Interval Property (Theorem 1.4.1) that doesn't make use of AoC.

These two results suggest that we could have used the Monotone Convergence Theorem in place of AoC as our starting axiom for building a proper theory of the real numbers.

### Solution

- (a) MCT tells us  $(1/n)$  converges, obviously it must converge to zero therefor we have  $|1/n - 0| = 1/n < \epsilon$  for any  $\epsilon$ , which is the Archimedean Property.
- (b) We have  $I_n = [a_n, b_n]$  with  $a_n \leq b_n$  since  $I_n \neq \emptyset$ . Since  $I_{n+1} \subseteq I_n$  we must have  $b_{n+1} \leq b_n$  and  $a_{n+1} \geq a_n$  the MCT tells us that  $(a_n) \rightarrow a$  and  $(b_n) \rightarrow b$ . by the Order Limit Theorem we have  $a \leq b$  since  $a_n \leq b_n$ , therefor  $a \in I_n$  for all  $n$  meaning  $a \in \bigcap_{n=1}^{\infty} I_n$  and thus  $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$ .

### Exercise 2.4.5 (Calculating Square Roots)

Let  $x_1 = 2$ , and define

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{2}{x_n} \right)$$

- (a) Show that  $x_n^2$  is always greater than or equal to 2, and then use this to prove that  $x_n - x_{n+1} \geq 0$ . Conclude that  $\lim x_n = \sqrt{2}$ .
- (b) Modify the sequence  $(x_n)$  so that it converges to  $\sqrt{c}$ .

### Solution

- (a) Clearly  $x_1^2 \geq 2$ , now procede by induction. if  $x_n^2 \geq 2$  then we have

$$x_{n+1}^2 = \frac{1}{4} \left( \frac{x_n^2 + 2}{x_n} \right)^2 = \frac{1}{4} \left( \frac{(x_n^2 + 2)^2}{x_n^2} \right) \geq \frac{1}{4} \left( \frac{(x_n^2 + 2)^2}{2} \right)$$

Now since  $x_n^2 \geq 2$  we have  $(x_n^2 + 2)^2 \geq 16$  meaning

$$x_{n+1}^2 = \frac{1}{4} \left( \frac{(x_n^2 + 2)^2}{2} \right) \geq 2.$$

To show  $x_n - x_{n+1} \geq 0$  we examine  $(x_n - x_{n+1})^2 \geq 0$  with the hope of using  $x_n^2 \geq 2$ .

$$\begin{aligned} (x_n - x_{n+1})^2 &= x_n^2 - 2x_n x_{n+1} + x_{n+1}^2 \\ &= x_n^2 - 2x_n \frac{1}{2} \left( x_n + \frac{1}{x_n} \right) + x_{n+1}^2 \\ &= x_n^2 - (x_n^2 + 2) + x_{n+1}^2 \\ &= x_{n+1}^2 - 2 \geq 0. \end{aligned}$$

Now we know  $(x_n) \rightarrow x$  converges, to show  $x^2 = 2$  observe that  $\lim x_n = \lim x_{n+1}$  so

$$x = \frac{1}{2} \left( x + \frac{2}{x} \right) \iff x^2 = \frac{1}{2}x^2 + 1 \iff x^2 = 2$$

Therefor  $x = \pm\sqrt{2}$ , since every  $x_n$  is positive  $x = \sqrt{2}$ .

(b) Let

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{c}{x_n} \right)$$

I won't go through the convergence analysis again, but the only fixed point is

$$x = \frac{1}{2} \left( x + \frac{c}{x} \right) \implies \frac{1}{2}x^2 = \frac{1}{2}c \implies x^2 = c$$

So if  $x_n$  converges, it must converge to  $x^2 = c$ .

### Exercise 2.4.6 (Arithmetic-Geometric Mean)

(a) Explain why  $\sqrt{xy} \leq (x + y)/2$  for any two positive real numbers  $x$  and  $y$ . (The geometric mean is always less than the arithmetic mean.)

(b) Now let  $0 \leq x_1 \leq y_1$  and define

$$x_{n+1} = \sqrt{x_n y_n} \quad \text{and} \quad y_{n+1} = \frac{x_n + y_n}{2}$$

Show  $\lim x_n$  and  $\lim y_n$  both exist and are equal.

### Solution

(a) We have

$$\sqrt{xy} \leq (x + y)/2 \iff 4xy \leq x^2 + 2xy + y^2 \iff 0 \leq (x - y)^2$$

(b) The only fixed point is  $x_n = y_n$  so we only need to show both sequences converge.

The inequality  $x_1 \leq y_1$  is always true since

$$\sqrt{x_n y_n} \leq \frac{x_n + y_n}{2} \implies x_{n+1} \leq y_{n+1}$$

Also  $x_n \leq y_n$  implies  $(x_n + y_n)/2 = y_{n+1} \leq y_n$ , similarly  $\sqrt{x_n y_n} = x_{n+1} \geq x_n$  meaning both sequences converge by the monotone convergence theorem.

### Exercise 2.4.7 (Limit Superior)

Let  $(a_n)$  be a bounded sequence.

(a) Prove that the sequence defined by  $y_n = \sup \{a_k : k \geq n\}$  converges.

(b) The limit superior of  $(a_n)$ , or  $\limsup a_n$ , is defined by

$$\limsup a_n = \lim y_n$$

where  $y_n$  is the sequence from part (a) of this exercise. Provide a reasonable definition for  $\liminf a_n$  and briefly explain why it always exists for any bounded sequence.

(c) Prove that  $\liminf a_n \leq \limsup a_n$  for every bounded sequence, and give an example of a sequence for which the inequality is strict.

- (d) Show that  $\liminf a_n = \limsup a_n$  if and only if  $\lim a_n$  exists. In this case, all three share the same value.

**Solution**

- (a)  $(y_n)$  is decreasing and converges by the monotone convergence theorem.
- (b) Define  $\liminf a_n = \lim z_n$  for  $z_n = \inf\{a_k : k \geq n\}$ .  $z_n$  converges since it is increasing and bounded.
- (c) Obviously  $\inf\{a_k : k \geq n\} \leq \sup\{a_k : k \geq n\}$  so by the Order Limit Theorem  $\liminf a_n \leq \limsup a_n$ .
- (d) If  $\liminf a_n = \limsup a_n$  then the squeeze theorem (Exercise 2.3.3) implies  $a_n$  converges to the same value, since  $\inf\{a_{k \geq n}\} \leq a_n \leq \sup\{a_{k \geq n}\}$ .

**Exercise 2.4.8**

For each series, find an explicit formula for the sequence of partial sums and determine if the series converges.

- (a)  $\sum_{n=1}^{\infty} \frac{1}{2^n}$
- (b)  $\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$
- (c)  $\sum_{n=1}^{\infty} \log\left(\frac{n+1}{n}\right)$

(In (c),  $\log(x)$  refers to the natural logarithm function from calculus.)

**Solution**

- (a) This is a geometric series, we can use the usual trick to derive  $s_n$ . Let  $r = 1/2$  for convenience

$$\begin{aligned} s_n &= 1 + r + r^2 + \cdots + r^n \\ r s_n &= r + r^2 + \cdots + r^{n+1} \\ r s_n - s_n &= r^{n+1} - 1 \implies s_n = \frac{r^{n+1} - 1}{r - 1} \end{aligned}$$

So we have

$$\sum_{n=1}^{\infty} \frac{1}{2^n} = \lim_{n \rightarrow \infty} \frac{(1/2)^{n+1} - 1}{1/2 - 1} = \frac{-1}{-1/2} = 2$$

- (b) We can use partial fractions to get

$$\frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}$$

Which gives us a telescoping series, most of the terms cancel and we get

$$s_n = 1 - \frac{1}{n+1}$$

Therefor

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n+1}\right) = 1$$



(c) Another telescoping series, since

$$\log\left(\frac{n+1}{n}\right) = \log(n+1) - \log(n)$$

Therefore most of the terms cancel and we get

$$s_n = \log(n+1)$$

Which doesn't converge.

### Exercise 2.4.9

Complete the proof of Theorem 2.4.6 by showing that if the series  $\sum_{n=0}^{\infty} 2^n b_{2^n}$  diverges, then so does  $\sum_{n=1}^{\infty} b_n$ . Example 2.4.5 may be a useful reference.

### Solution

Let  $s_n = b_1 + b_2 + \cdots + b_n$  and  $t_k = b_1 + 2b_2 + \cdots + 2^k b_{2^k}$ .

We want to show  $s_n$  is unbounded, first we find a series similar to  $t_k$  that is less than  $s_n$ , then rewrite it in terms of  $t_k$ .

Let  $n = 2^k$  so things match up nicely. We get

$$\begin{aligned} s_n &= b_1 + b_2 + (b_3 + b_4) + \cdots + (b_{2^{k-1}} + \cdots + b_{2^k}) \\ &\leq b_1 + b_2 + (b_4 + b_4) + \cdots + 2^{k-1} b_{2^k} \end{aligned}$$

(Notice there are  $2^k - 2^{k-1} = 2^{k-1}$  terms in the last term)

Now define  $t'_k$  to be our new series  $b_1 + b_2 + 2b_4 + 4b_8 + \cdots + 2^{k-1} b_{2^k}$ . This looks a lot like  $t_k$ , and in fact some algebra gives

$$t'_k = \frac{1}{2} (b_1 + 2b_2 + 4b_4 + \cdots + 2^k b_k) + \frac{1}{2} b_1 = \frac{1}{2} t_k + \frac{1}{2} b_1$$

Therefore we are justified in writing

$$s_n \geq t'_k \geq \frac{1}{2} t_k$$

And since  $t_k/2$  diverges and  $s_n$  is bigger,  $s_n$  must also diverge.

**Summary:**  $s_n$  converges iff  $t_k$  conv since  $t_k \geq s_n \geq t_k/2$  for  $n = 2^k$ .

### Exercise 2.4.10 (Infinite Products)

A close relative of infinite series is the infinite product

$$\prod_{n=1}^{\infty} b_n = b_1 b_2 b_3 \cdots$$

which is understood in terms of its sequence of partial products

$$p_m = \prod_{n=1}^m b_n = b_1 b_2 b_3 \cdots b_m$$

Consider the special class of infinite products of the form

$$\prod_{n=1}^{\infty} (1 + a_n) = (1 + a_1)(1 + a_2)(1 + a_3) \cdots, \quad \text{where } a_n \geq 0$$

- (a) Find an explicit formula for the sequence of partial products in the case where  $a_n = 1/n$  and decide whether the sequence converges. Write out the first few terms in the sequence of partial products in the case where  $a_n = 1/n^2$  and make a conjecture about the convergence of this sequence.
- (b) Show, in general, that the sequence of partial products converges if and only if  $\sum_{n=1}^{\infty} a_n$  converges. (The inequality  $1 + x \leq 3^x$  for positive  $x$  will be useful in one direction.)

### Solution

- (a) This is a telescoping product, most of the terms cancel

$$p_m = \prod_{n=1}^m (1 + 1/n) = \prod_{n=1}^m \frac{n+1}{n} = \frac{2}{1} \cdot \frac{3}{2} \cdot \frac{4}{3} \cdots \frac{m+1}{m} = m+1$$

Therefore  $(p_m)$  diverges.

In the case  $a_n = 1/n^2$  we get

$$\prod_{n=1}^{\infty} (1 + 1/n^2) = \prod_{n=1}^{\infty} \frac{1+n^2}{n^2} = \frac{2}{1} \cdot \frac{5}{4} \cdot \frac{10}{9} \cdots$$

The growth seems slower, I conjecture it converges now.

- (b) Using the inequality suggested we have  $1 + a_n \leq 3^{a_n}$  letting  $s_m = a_1 + \cdots + a_m$  we get

$$p_m = (1 + a_1) \cdots (1 + a_m) \leq 3^{a_1} 3^{a_2} \cdots 3^{a_m} = 3^{s_m}$$

Now if  $s_m$  converges it is bounded by some  $M$  meaning  $p_m$  is bounded by  $3^M$ . and because  $a_n \geq 0$  the partial products  $p_m$  are increasing, so they converge by the MCT. This shows  $s_m$  converging implies  $p_m$  converges.

For the other direction suppose  $p_m \rightarrow p$ . Distributing inside the products gives  $p_2 = a_1 + a_2 + 1 + a_1 a_2 > s_2$  and in general  $p_m > s_m$  implying that if  $p_m$  is bounded then  $s_n$  is bounded as well. This completes the proof.

**Summary:** Convergence is if and only if because  $s_m \leq p_m \leq 3^{s_m}$ .

(By the way the inequality  $1 + x \leq 3^x$  can be derived from  $\log(1+x) \leq x$  implying  $1+x \leq e^x$ , I assume abbot rounded up to 3.)

## 2.5 Subsequences and the Bolzano–Weierstrass Theorem

### Exercise 2.5.1

Give an example of each of the following, or argue that such a request is impossible.

- (a) A sequence that has a subsequence that is bounded but contains no subsequence that converges.
- (b) A sequence that does not contain 0 or 1 as a term but contains subsequences converging to each of these values.
- (c) A sequence that contains subsequences converging to every point in the infinite set  $\{1, 1/2, 1/3, 1/4, 1/5, \dots\}$ .
- (d) A sequence that contains subsequences converging to every point in the infinite set  $\{1, 1/2, 1/3, 1/4, 1/5, \dots\}$ , and no subsequences converging to points outside of this set.

### Solution

- (a) Impossible, the Bolzano–Weierstrass theorem tells us a convergent subsequence of that subsequence exists, and that sub-sub sequence is also a subsequence of the original sequence.
- (b)  $(1 + 1/n) \rightarrow 1$  and  $(1/n) \rightarrow 0$  so  $(1/2, 1 + 1/2, 1/3, 1 + 1/3, \dots)$  has subsequences converging to 0 and 1.
- (c) Copy the finitely many previous terms before proceeding to a new term

$$(1, 1/2, 1, 1/3, 1, 1/2, 1/4, 1, 1/2, 1/3, \dots)$$

The sequence contains infinitely many terms in  $\{1, 1/2, 1/3, \dots\}$  hence subsequences exist converging to each of these values.

- (d) Impossible, the sequence must converge to zero which is not in the set.

Proof: Let  $\epsilon > 0$  be arbitrary, pick  $N$  large enough that  $1/n < \epsilon/2$  for  $n > N$ . We can find a subsequence  $(b_m) \rightarrow 1/n$  meaning  $|b_m - 1/n| < \epsilon/2$  for some  $m$ . using the triangle inequality we get

$$|b_m - 0| \leq |b_m - 1/n| + |1/n - 0| < \epsilon/2 + \epsilon/2 = \epsilon$$

Therefor we have found a number  $b_m$  in the sequence  $a_m$  with  $|b_m| < \epsilon$ . This process can be repeated for any  $\epsilon$  therefor a sequence which converges to zero can be constructed.

### Exercise 2.5.2

Decide whether the following propositions are true or false, providing a short justification for each conclusion.

- (a) If every proper subsequence of  $(x_n)$  converges, then  $(x_n)$  converges as well.

- (b) If  $(x_n)$  contains a divergent subsequence, then  $(x_n)$  diverges.
- (c) If  $(x_n)$  is bounded and diverges, then there exist two subsequences of  $(x_n)$  that converge to different limits.
- (d) If  $(x_n)$  is monotone and contains a convergent subsequence, then  $(x_n)$  converges.

**Solution**

- (a) True, removing the first term gives us the proper subsequence  $(x_2, x_3, \dots)$  which converges, implying  $(x_1, x_2, \dots)$  also converges.
- (b) True, the divergent subsequence is unbounded, hence  $(x_n)$  is also unbounded and divergent.
- (c) True, since  $x_n$  is bounded  $\limsup x_n$  and  $\liminf x_n$  both converge. And since  $x_n$  diverges Exercise 2.4.7 tells us  $\limsup x_n \neq \liminf x_n$ .
- (d) True, The subsequence  $(x_{n_k})$  converges meaning it is bounded  $|x_{n_k}| \leq M$ . Suppose  $(x_n)$  is increasing, then  $x_n$  is bounded since picking  $k$  so that  $n_k > n$  we have  $x_n \leq x_{n_k} \leq M$ . A similar argument applies if  $x_n$  is decreasing, Therefore  $x_n$  is monotonic bounded and so must converge.

**Exercise 2.5.3**

- (a) Prove that if an infinite series converges, then the associative property holds. Assume  $a_1 + a_2 + a_3 + a_4 + a_5 + \dots$  converges to a limit  $L$  (i.e., the sequence of partial sums  $(s_n) \rightarrow L$ ). Show that any regrouping of the terms

$$(a_1 + a_2 + \dots + a_{n_1}) + (a_{n_1+1} + \dots + a_{n_2}) + (a_{n_2+1} + \dots + a_{n_3}) + \dots$$

leads to a series that also converges to  $L$ .

- (b) Compare this result to the example discussed at the end of Section 2.1 where infinite addition was shown not to be associative. Why doesn't our proof in (a) apply to this example?

**Solution**

- (a) Let  $s_n$  be the original partial sums, and let  $s'_m$  be the regrouping. Since  $s'_m$  is a subsequence of  $s_n$ ,  $(s_n) \rightarrow s$  implies  $(s'_m) \rightarrow s$ .
- (b) The subsequence  $s'_m = (1 - 1) + \dots = 0$  converging does not imply the parent sequence  $s_n$  converges. In fact BW tells us any bounded sequence of partial sums will have a convergent subsequence (regrouping in this case).

**Exercise 2.5.4**

The Bolzano-Weierstrass Theorem is extremely important, and so is the strategy employed in the proof. To gain some more experience with this technique, assume the Nested Interval Property is true and use it to provide a proof of the Axiom of Completeness. To prevent the argument from being circular, assume also that  $(1/2^n) \rightarrow 0$ . (Why precisely is this last assumption needed to avoid circularity?)

**Solution**

Let  $A$  be a bounded set, we're basically going to binary search for  $\sup A$  and then use NIP to prove the limit exists.

Let  $M$  be an upper bound on  $A$ , and pick any  $L \in A$  as our starting lower bound for  $\sup A$  and define  $I_1 = [L, M]$ . Doing binary search gives  $I_{n+1} \subseteq I_n$  with length proportional to  $(1/2)^n$ . Applying the Nested Interval Property gives

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset$$

As the length  $(1/2)^n$  goes to zero, there is a single  $s \in \bigcap_{n=1}^{\infty} I_n$  which must be the least upper bound since  $I_n = [L_n, M_n]$  gives  $L_n \leq x \leq M_n$  for all  $n$  meaning  $s = \sup A$  since

- (i)  $s \geq L_n$  implies  $s$  is an upper bound
- (ii)  $s \leq M_n$  implies  $s$  is the least upper bound

**TODO** Draw a diagram with tikz, Make this rigorous

**Exercise 2.5.5**

Assume  $(a_n)$  is a bounded sequence with the property that every convergent subsequence of  $(a_n)$  converges to the same limit  $a \in \mathbf{R}$ . Show that  $(a_n)$  must converge to  $a$ .

**Solution**

$(a_2, a_3, \dots)$  Is a convergent subsequence, so obviously if  $(a_2, a_3, \dots) \rightarrow a$  then  $(a_n) \rightarrow a$  also.

**Exercise 2.5.6**

Use a similar strategy to the one in Example 2.5.3 to show  $\lim b^{1/n}$  exists for all  $b \geq 0$  and find the value of the limit. (The results in Exercise 2.3.1 may be assumed.)

**Solution**

Intuitively  $\lim b^{1/n} = 1$

Two facts I'll take as granted (you can prove them if you wish)

- (i) If  $b < 1$  then  $b^{1/n}$  is increasing
- (ii) If  $b > 1$  then  $b^{1/n}$  is decreasing

Thus  $b^{1/n}$  is monotonic, and bounded since

- (i) If  $b > 1$  then  $b^{1/n} > 1$  since  $b > 1^n$
- (ii) If  $b < 1$  then  $b^{1/n} < 1$  since  $b < 1^n$

Therefor  $b^{1/n}$  converges by the monotone convergence theorem. to find the limit equate terms

$$b^{1/n+1} = b^{1/n} \implies b^1 = b^{\frac{n+1}{n}} = b^2 \implies b = 1$$

**TODO** Shorten this

**Exercise 2.5.7**

Extend the result proved in Example 2.5.3 to the case  $|b| < 1$ ; that is, show  $\lim(b^n) = 0$  if and only if  $-1 < b < 1$ .

**Solution**

If  $|b| \geq 1$  then  $\lim(b^n) \neq 0$  (diverges for  $b \neq 1$ ).

Now for the other direction, if  $|b| < 1$  we immediately get  $|b^n| < 1$  thus  $b^n$  is bounded. Since it is decreasing the monotone convergence theorem implies it converges. To find the limit equating terms  $b^{n+1} = b^n$  gives  $b = 0$  or  $b = 1$ , since  $b$  is *strictly* decreasing we have  $b = 0$ .

**Exercise 2.5.8**

Another way to prove the Bolzano-Weierstrass Theorem is to show that every sequence contains a monotone subsequence. A useful device in this endeavor is the notion of a peak term. Given a sequence  $(x_n)$ , a particular term  $x_m$  is a peak term if no later term in the sequence exceeds it; i.e., if  $x_m \geq x_n$  for all  $n \geq m$ .

- (a) Find examples of sequences with zero, one, and two peak terms. Find an example of a sequence with infinitely many peak terms that is not monotone.
- (b) Show that every sequence contains a monotone subsequence and explain how this furnishes a new proof of the Bolzano-Weierstrass Theorem.

**Solution**

- (a)  $(1, 2, \dots)$  has zero peak terms,  $(1, 0, 1/2, 2/3, 3/4, \dots)$  has a single peak term,  $(2, 1, 1/2, 2/3, \dots)$  has two peak terms (a similar argument works for  $k$  peak terms) and  $(1, 1/2, 1/3, \dots)$  has infinitely many peak terms. The sequence  $(1, -1/2, 1/3, -1/4, \dots)$  has infinitely many peak terms, but is not monotone.
- (b) The sequence of peak terms is monotonic decreasing, thus if the parent sequence is bounded we have found a subsequence which converges, hence proving BW. (If there aren't infinitely many peak terms, then take the sequence of valley terms)

**Exercise 2.5.9**

Let  $(a_n)$  be a bounded sequence, and define the set

$$S = \{x \in \mathbf{R} : x < a_n \text{ for infinitely many terms } a_n\}$$

Show that there exists a subsequence  $(a_{n_k})$  converging to  $s = \sup S$ . (This is a direct proof of the Bolzano-Weierstrass Theorem using the Axiom of Completeness.)

**Solution**

For every  $\epsilon > 0$  there exists an  $x \in S$  with  $x > s - \epsilon$  implying  $|s - x| < \epsilon$ . Therefor we can get arbitrarily close to  $s = \sup S$  so there is a subsequence converging to this value.

To make this more rigorous, pick  $x_n \in S$  such that  $|x_n - s| < 1/n$  then pick  $N > 1/\epsilon$  to get  $|x_n - s| < \epsilon$  for all  $n > N$ .

## 2.6 The Cauchy Criterion

### Exercise 2.6.1

Prove every convergent sequence is a Cauchy sequence. (Theorem 2.6.2)

### Solution

Suppose  $(x_n)$  is convergent, we must show that for  $m, n > N$  we have  $|x_n - x_m| < \epsilon$

Set  $|x_n - x| < \epsilon/2$  for  $n > N$ .

We get  $|x_n - x_m| \leq |x_n - x| + |x - x_m| \leq \epsilon/2 + \epsilon/2 = \epsilon$

### Exercise 2.6.2

Give an example of each of the following, or argue that such a request is impossible.

- (a) A Cauchy sequence that is not monotone.
- (b) A Cauchy sequence with an unbounded subsequence.
- (c) A divergent monotone sequence with a Cauchy subsequence.
- (d) An unbounded sequence containing a subsequence that is Cauchy.

### Solution

- (a)  $x_n = (-1)^n/n$  is cauchy by Theorem 2.6.2.
- (b) Impossible since all cauchy sequences converge.
- (c) Impossible, If a subsequence was cauchy it would converge, implying the subsequence would be bounded and therefor the parent sequence would be bounded (because it is monotone) and thus would converge.
- (d)  $(2, 1/2, 3, 1/3, \dots)$  has subsequence  $(1/2, 1/3, \dots)$  which is cauchy.

### Exercise 2.6.3

If  $(x_n)$  and  $(y_n)$  are Cauchy sequences, then one easy way to prove that  $(x_n + y_n)$  is Cauchy is to use the Cauchy Criterion. By Theorem 2.6.4,  $(x_n)$  and  $(y_n)$  must be convergent, and the Algebraic Limit Theorem then implies  $(x_n + y_n)$  is convergent and hence Cauchy.

- (a) Give a direct argument that  $(x_n + y_n)$  is a Cauchy sequence that does not use the Cauchy Criterion or the Algebraic Limit Theorem.
- (b) Do the same for the product  $(x_n y_n)$ .

### Solution

- (a) We have  $|(x_n + y_n) - (x_m + y_m)| \leq |x_n - x_m| + |y_n - y_m| < \epsilon/2 + \epsilon/2 = \epsilon$
- (b) Bound  $|x_n| \leq M_1$ , and  $|y_n| \leq M_2$  then

$$\begin{aligned} |x_n y_n - x_m y_m| &= |(x_n y_n - x_n y_m) + (x_n y_m - x_m y_m)| \\ &\leq |x_n(y_n - y_m)| + |y_m(x_n - x_m)| \\ &\leq M_1|y_n - y_m| + M_2|x_n - x_m| \\ &< \epsilon/2 + \epsilon/2 = \epsilon \end{aligned}$$

After setting  $|y_n - y_m| < \epsilon/(2M_1)$  and  $|x_n - x_m| < \epsilon/(2M_2)$ .

**Exercise 2.6.4**

Let  $(a_n)$  and  $(b_n)$  be Cauchy sequences. Decide whether each of the following sequences is a Cauchy sequence, justifying each conclusion.

- (a)  $c_n = |a_n - b_n|$
- (b)  $c_n = (-1)^n a_n$
- (c)  $c_n = [[a_n]]$ , where  $[[x]]$  refers to the greatest integer less than or equal to  $x$ .

**Solution**

- (a) Yes, since  $|(a_n - b_n) - (a_m - b_m)| \leq |a_n - a_m| + |b_m - b_n| < \epsilon/2 + \epsilon/2 = \epsilon$
- (b) No, if  $a_n = 1$  then  $(-1)^n a_n$  diverges, and thus is not Cauchy.
- (c) No, if  $a_n = 1 - (-1)^n/n$  then  $[[a_n]]$  fluctuates between 0 and 1 and so cannot be Cauchy.

**Exercise 2.6.5**

Consider the following (invented) definition: A sequence  $(s_n)$  is pseudo-Cauchy if, for all  $\epsilon > 0$ , there exists an  $N$  such that if  $n \geq N$ , then  $|s_{n+1} - s_n| < \epsilon$

Decide which one of the following two propositions is actually true. Supply a proof for the valid statement and a counterexample for the other.

- (i) Pseudo-Cauchy sequences are bounded.
- (ii) If  $(x_n)$  and  $(y_n)$  are pseudo-Cauchy, then  $(x_n + y_n)$  is pseudo-Cauchy as well.

**Solution**

- (i) False, consider  $s_n = \log n$ . clearly  $|s_{n+1} - s_n|$  can be made arbitrarily small but  $s_n$  is unbounded.
- (ii) True, as  $|(x_{n+1} + y_{n+1}) - (x_n + y_n)| \leq |x_{n+1} - x_n| + |y_{n+1} - y_n| < \epsilon/2 + \epsilon/2 = \epsilon$ .

**Exercise 2.6.6**

Let's call a sequence  $(a_n)$  quasi-increasing if for all  $\epsilon > 0$  there exists an  $N$  such that whenever  $n > m \geq N$  it follows that  $a_n > a_m - \epsilon$

- (a) Give an example of a sequence that is quasi-increasing but not monotone or eventually monotone.
- (b) Give an example of a quasi-increasing sequence that is divergent and not monotone or eventually monotone.
- (c) Is there an analogue of the Monotone Convergence Theorem for quasiincreasing sequences? Give an example of a bounded, quasi-increasing sequence that doesn't converge, or prove that no such sequence exists.

**Solution**

- (a)  $a_n = (-1)^n/n$  is quasi-increasing since we can get  $(-1)^m/m - (-1)^n/n \leq 1/m + 1/n < \epsilon$  for large enough  $n > m \geq N$ .



- (b)  $a_n = (2, 1/2, 3, 1/3, \dots)$  is quasi-increasing since  $n > m - \epsilon$  is clearly true as  $(n)$  is increasing. And  $1/n > 1/m - \epsilon$  is true after picking  $N$  large enough that for  $m \geq N$  we have  $1/m < \epsilon$  and thus  $1/n > 1/m - \epsilon$ .
- (c) In (b) I gave such an example, so there is no Monotone Convergence Theorem for quasiincreasing sequences (without modifying the definition that is.)

**Exercise 2.6.7**

Exercises 2.4.4 and 2.5.4 establish the equivalence of the Axiom of Completeness and the Monotone Convergence Theorem. They also show the Nested Interval Property is equivalent to these other two in the presence of the Archimedean Property.

- (a) Assume the Bolzano-Weierstrass Theorem is true and use it to construct a proof of the Monotone Convergence Theorem without making any appeal to the Archimedean Property. This shows that BW, AoC, and MCT are all equivalent.
- (b) Use the Cauchy Criterion to prove the Bolzano-Weierstrass Theorem, and find the point in the argument where the Archimedean Property is implicitly required. This establishes the final link in the equivalence of the five characterizations of completeness discussed at the end of Section 2.6.
- (c) How do we know it is impossible to prove the Axiom of Completeness starting from the Archimedean Property?

**Solution**

- (a) Suppose  $(x_n)$  is increasing and bounded, BW tells us there exists a convergent subsequence  $(x_{n_k}) \rightarrow x$ . We will show  $(x_n) \rightarrow x$ . First note  $x_k \leq x_{n_k}$  implies  $x_n \leq x$  by the Order Limit Theorem.

Pick  $K$  such that for  $k \geq K$  we have  $|x_{n_k} - x| < \epsilon$ . Since  $(x_n)$  is increasing and  $x_n \leq x$  every  $n \geq n_K$  satisfies  $|x_n - x| < \epsilon$  as well. Thus  $(x_n)$  converges, completing the proof.

- (b) We're basically going to use the Cauchy criterion as a replacement for NIP in the proof of BW. Recall we had  $I_{n+1} \subseteq I_n$  with  $a_{n_k} \in I_k$ , we will show  $a_{n_k}$  is Cauchy.

The length of  $I_k$  is  $M(1/2)^{k-1}$  by construction, so clearly  $|a_{n_k} - a_{n_j}| < M(1/2)^{N-1}$  for  $k, j \geq N$ , implying  $(a_{n_k})$  converges by the Cauchy criterion.

We needed the Archimedean Property to conclude  $M(1/2)^{N-1} \in \mathbf{Q}$  can be made smaller than any  $\epsilon \in \mathbf{R}^+$ .

- (c) The Archimedean Property is true for  $\mathbf{Q}$  meaning it cannot prove AoC which is only true for  $\mathbf{R}$ . (If we did, then we would have proved AoC for  $\mathbf{Q}$  which is obviously false.)

## 2.7 Properties of Infinite Series

### Exercise 2.7.1

Proving the Alternating Series Test (Theorem 2.7.7) amounts to showing that the sequence of partial sums

$$s_n = a_1 - a_2 + a_3 - \cdots \pm a_n$$

converges. (The opening example in Section 2.1 includes a typical illustration of  $(s_n)$ .) Different characterizations of completeness lead to different proofs.

- (a) Prove the Alternating Series Test by showing that  $(s_n)$  is a Cauchy sequence.
- (b) Supply another proof for this result using the Nested Interval Property (Theorem 1.4.1).
- (c) Consider the subsequences  $(s_{2n})$  and  $(s_{2n+1})$ , and show how the Monotone Convergence Theorem leads to a third proof for the Alternating Series Test.

### Solution

- (a) We would like to show  $|a_{m+1} - a_{m+2} + \cdots \pm a_n|$  becomes arbitrarily small. **TODO**
- (b) Let  $I_1$  be the interval  $[a_1 - a_2, a_1]$  and in general  $I_n = [a_n - a_{n+1}, a_n]$  we have  $I_{n+1} \subseteq I_n$  since  $(a_n)$  is decreasing. The nested interval property gives

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset$$

Let  $x \in \bigcap_{n=1}^{\infty} I_n$ , since  $a_n \in I_n$  and  $x \in I_n$  the distance  $|a_n - x|$  must be less than the length  $|I_n|$ . and since the length goes to zero  $|a_n - x|$  can be made less than any  $\epsilon$ .

- (c) If we can show  $\lim s_{2n} = \lim s_{2n+1} = s$  that will imply  $\lim s_n = s$  since each  $s_n$  is either in  $(s_{2n+1})$  or in  $(s_{2n})$  as  $n$  must be even or odd.

We have  $s_{2n+1} \leq a_1$  since

$$s_{2n+1} = a_1 - (a_2 - a_3) - \cdots - (a_{2n} - a_{2n+1}) \leq a_1$$

Thus  $s_{2n+1} \rightarrow s$  by the Monotone Convergence Theorem, to show  $(s_{2n}) \rightarrow s$  notice  $s_{2n} = s_{2n+1} - a_{2n+1}$  with  $(a_{2n+1}) \rightarrow 0$  meaning we can use the triangle inequality

$$|s_{2n} - s| \leq \underbrace{|s_{2n} - s_{2n+1}|}_{a_{2n+1}} + |s_{2n+1} - s| < \epsilon/2 + \epsilon/2 < \epsilon$$

Thus  $(s_{2n}) \rightarrow s$  as well finally implying  $(s_n) \rightarrow s$ .

**Summary:** Partition the alternating series into two subsequences of partial sums, then use MCT to show they both converge to the same limit.

### Exercise 2.7.2

Decide whether each of the following series converges or diverges:

- (a)  $\sum_{n=1}^{\infty} \frac{1}{2^n + n}$
- (b)  $\sum_{n=1}^{\infty} \frac{\sin(n)}{n^2}$
- (c)  $1 - \frac{3}{4} + \frac{4}{6} - \frac{5}{8} + \frac{6}{10} - \frac{7}{12} + \dots$
- (d)  $1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} + \frac{1}{8} - \frac{1}{9} + \dots$
- (e)  $1 - \frac{1}{2^2} + \frac{1}{3} - \frac{1}{4^2} + \frac{1}{5} - \frac{1}{6^2} + \frac{1}{7} - \frac{1}{8^2} + \dots$

**Solution**

- (a) Converges by a comparison test with  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- (b) Converges by a comparison test with  $\sum_{n=1}^{\infty} \frac{1}{n^2}$ .
- (c) Diverges since  $(n+1)/2n = 1/2 + 1/2n$  never gets smaller than  $1/2$ .
- (d) The first thing I notice is the series cannot converge absolutely (as then it would be harmonic).

Grouping terms gives

$$\begin{aligned} \frac{1}{n} + \frac{1}{n+1} - \frac{1}{n+2} &= \frac{(n+1)(n+2) + n(n+2) - n(n+1)}{n(n+1)(n+2)} \\ &= \frac{(n^2 + 3n + 2) + (n^2 + 2n) - (n^2 + n)}{n(n+1)(n+2)} \\ &= \frac{n^2 + 4n + 2}{n(n+1)(n+2)} \end{aligned}$$

Which diverges since

$$\frac{n^2 + 4n + 2}{n(n+1)(n+2)} \geq \frac{n^2 + 4n + 2}{n^3} \geq \frac{1}{n}$$

In other words, If we take terms three at a time (a subsequence of the partial sums) grows faster than the harmonic series implying the subsequence (and thus the parent sequence) of partial sums diverges.

**TODO** Find better proof

- (e) Intuitively this should diverge since it is a mixture of  $1/n$  (divergent) and  $1/n^2$  (convergent). To make this rigorous examine the subsequence  $(s_{2n})$

$$s_{2n} = 1 - 1/2^2 + 1/3 - 1/4^2 + \dots + 1/(2n)^2$$

We can rearrange terms in the partial sum to get

$$s_{2n} = \sum_{k=1}^n \frac{1}{2k-1} - \sum_{k=1}^n \frac{1}{(2k)^2} \equiv t_n - v_n$$

Now if  $\lim(s_{2n})$  converges then  $\lim(t_n - v_n)$  also converges. This is impossible however as  $(t_n)$  diverging and  $(v_n)$  converging would imply  $\lim(t_n - v_n) + \lim(v_n) = \lim(t_n)$  converges by the ALT, but  $(t_n)$  diverges so we have a contradiction. Thus  $(s_{2n})$  diverges and so the parent sequence  $(s_n)$  must also diverge.

**Exercise 2.7.3**

- (a) Provide the details for the proof of the Comparison Test (Theorem 2.7.4) using the Cauchy Criterion for Series.
- (b) Give another proof for the Comparison Test, this time using the Monotone Convergence Theorem.

**Solution**

Suppose  $a_n, b_n \geq 0$ ,  $a_n \leq b_n$  and define  $s_n = a_1 + \cdots + a_n$ ,  $t_n = b_1 + \cdots + b_n$ .

- (a) We have  $|a_{m+1} + \cdots + a_n| \leq |b_{m+1} + \cdots + b_n| < \epsilon$  implying  $\sum_{n=1}^{\infty} a_n$  converges by the cauchy criterion. The other direction is analogous, if  $(s_n)$  diverges then  $(t_n)$  must also diverge since  $s_n \leq t_n$ .
- (b) Since  $(t_n) \rightarrow t$ . This implies that  $s_n$  is bounded, and since  $s_n \leq t_n$  implies  $s_n \leq t$  by the order limit theorem, we can use the monotone convergence theorem to conclude  $(s_n)$  converges.

**Exercise 2.7.4**

Give an example of each or explain why the request is impossible referencing the proper theorem(s).

- (a) Two series  $\sum x_n$  and  $\sum y_n$  that both diverge but where  $\sum x_n y_n$  converges.
- (b) A convergent series  $\sum x_n$  and a bounded sequence  $(y_n)$  such that  $\sum x_n y_n$  diverges.
- (c) Two sequences  $(x_n)$  and  $(y_n)$  where  $\sum x_n$  and  $\sum (x_n + y_n)$  both converge but  $\sum y_n$  diverges.
- (d) A sequence  $(x_n)$  satisfying  $0 \leq x_n \leq 1/n$  where  $\sum (-1)^n x_n$  diverges.

**Solution**

- (a)  $x_n = 1/n$  and  $y_n = 1/n$  have their respective series diverge, but  $\sum x_n y_n = \sum 1/n^2$  converges since it is a p-series with  $p > 1$ .
- (b) Let  $x_n = (-1)^n/n$  and  $y_n = (-1)^n$ .  $\sum x_n$  converges but  $\sum x_n y_n = \sum 1/n$  diverges.
- (c) Impossible as the algebraic limit theorem for series implies  $\sum (x_n + y_n) - \sum x_n = \sum y_n$  converges.
- (d) Impossible as the alternating series test implies it converges.

**Exercise 2.7.5**

Prove the series  $\sum_{n=1}^{\infty} 1/n^p$  converges if and only if  $p > 1$ . (Corollary 2.4.7)

**Solution**

Eventually we have  $1/n^p < 1/p^n$  for  $p > 1$  (polynomial vs exponential) meaning we can use the comparison test to conclude  $\sum_{n=1}^{\infty} 1/n^p$  converges if  $p > 1$ .

Now suppose  $p \leq 1$ , since  $1/n^p \leq 1/n$  a comparison test with the harmonic series implies  $p \leq 1$  diverges.

**Exercise 2.7.6**

Let's say that a series subverges if the sequence of partial sums contains a subsequence that converges. Consider this (invented) definition for a moment, and then decide which of the following statements are valid propositions about subvergent series:

- (a) If  $(a_n)$  is bounded, then  $\sum a_n$  subverges.
- (b) All convergent series are subvergent.
- (c) If  $\sum |a_n|$  subverges, then  $\sum a_n$  subverges as well.
- (d) If  $\sum a_n$  subverges, then  $(a_n)$  has a convergent subsequence.

**Solution**

- (a) False, consider  $a_n = 1$  then  $s_n = n$  does not have a convergent subsequence.
- (b) True, every subsequence converges to the same limit in fact.
- (c) True, since  $s_n = \sum_{k=1}^n |a_k|$  converges it is bounded  $|s_n| \leq M$ , and since  $t_n = \sum_{k=1}^n a_k$  is smaller  $t_n \leq s_n$  it is bounded  $t_n \leq M$  which by BW implies there exists a convergent subsequence  $(t_{n_k})$ .
- (d) False,  $a_n = (1, -1, 2, -2, \dots)$  has no convergent subsequence but the sum  $s_n = \sum_{k=1}^n a_k$  has the subsequence  $(s_{2n}) \rightarrow 0$ .

**Exercise 2.7.7**

- (a) Show that if  $a_n > 0$  and  $\lim(na_n) = l$  with  $l \neq 0$ , then the series  $\sum a_n$  diverges.
- (b) Assume  $a_n > 0$  and  $\lim(n^2a_n)$  exists. Show that  $\sum a_n$  converges.

**Solution**

Note: This is kind of like a wierd way to do a comparison with  $1/n$  and  $1/n^2$ .

- (a) If  $\lim(na_n) = l \neq 0$  then  $na_n \in (l - \epsilon, l + \epsilon)$ , setting  $\epsilon = l/2$  gives  $na_n \in (l/2, 3l/2)$  implying  $a_n > (l/2)(1/n)$ . But if  $a_n > (l/2)(1/n)$  then  $\sum a_n$  diverges as it is a multiple of the harmonic series. (note that  $a_n > 0$  ensures  $l \geq 0$ .)
- (b) Letting  $l = \lim(n^2a_n)$  we have  $n^2a_n \in (l - \epsilon, l + \epsilon)$  setting  $\epsilon = l$  gives  $n^2a_n \in (0, 2l)$  implying  $0 \leq a_n \leq 2l/n^2$  and so  $\sum a_n$  converges by a comparsion test with  $\sum 2l/n^2$ .

**Exercise 2.7.8**

Consider each of the following propositions. Provide short proofs for those that are true and counterexamples for any that are not.

- (a) If  $\sum a_n$  converges absolutely, then  $\sum a_n^2$  also converges absolutely.
- (b) If  $\sum a_n$  converges and  $(b_n)$  converges, then  $\sum a_nb_n$  converges.
- (c) If  $\sum a_n$  converges conditionally, then  $\sum n^2a_n$  diverges.

**Solution**

- (a) True since  $(a_n) \rightarrow 0$  so eventually  $a_n^2 \leq |a_n|$  meaning  $\sum a_n^2$  converges by a comparison test with  $\sum |a_n|$ .
- (b) False, let  $a_n = (-1)^n/\sqrt{n}$  and  $b_n = (-1)^n/\sqrt{n}$ .  $\sum a_n$  converges by the alternating series test, but  $\sum a_n b_n = \sum 1/n$  diverges.
- (c) True, suppose  $(n^2 a_n)$  converges, since  $(n^2 a_n) \rightarrow 0$  we have  $|n^2 a_n| < 1$  for  $n > N$ , implying  $|a_n| < 1/n^2$ . But if  $|a_n| < 1/n^2$  then a comparison test with  $1/n^2$  implies  $a_n$  converges absolutely, contradicting the assumption that  $a_n$  converges conditionally. Therefore  $\sum n^2 a_n$  must diverge.

**Exercise 2.7.9 (Ratio Test)**

Given a series  $\sum_{n=1}^{\infty} a_n$  with  $a_n \neq 0$ , the Ratio Test states that if  $(a_n)$  satisfies

$$\lim \left| \frac{a_{n+1}}{a_n} \right| = r < 1$$

then the series converges absolutely.

- (a) Let  $r'$  satisfy  $r < r' < 1$ . Explain why there exists an  $N$  such that  $n \geq N$  implies  $|a_{n+1}| \leq |a_n| r'$ .
- (b) Why does  $|a_n| \sum (r')^n$  converge?
- (c) Now, show that  $\sum |a_n|$  converges, and conclude that  $\sum a_n$  converges.

**Solution**

- (a) We are given

$$\left| \frac{a_{n+1}}{a_n} - r \right| < \epsilon$$

Since  $1 > r' > r$  we can set  $\epsilon = r' - r$  meaning the neighborhood

$$\frac{a_{n+1}}{a_n} \in (r - \epsilon, r + \epsilon) = (2r - r', r')$$

is all less than  $r'$  meaning

$$\left| \frac{a_{n+1}}{a_n} \right| \leq r' \implies |a_{n+1}| \leq r' |a_n|$$

- (b) Let  $N$  be large enough that for  $n > N$  we have  $|a_{n+1}| \leq |a_n| r'$ . Applying this multiple times gives  $|a_n| \leq (r')^{n-N} |a_N|$  which gives

$$|a_N| + |a_{N+1}| + \cdots + |a_n| \leq |a_N| + r'|a_N| + \cdots + (r')^{n-N} |a_N|$$

Factoring out  $|a_N|$  and writing with sums gives

$$\sum_{k=N}^n |a_k| \leq |a_N| \sum_{k=0}^{n-N} (r')^k$$

Which converges as  $n \rightarrow \infty$  since  $|r'| < 1$  and  $|a_N|$  is constant. Implying  $\sum_{k=N}^n |a_k|$  converges and thus  $\sum_{k=1}^n |a_k|$  also converges since we only omitted finitely many terms.

(c) See (b)

### Exercise 2.7.10 (Infinite Products)

Review Exercise 2.4.10 about infinite products and then answer the following questions:

- (a) Does  $\frac{2}{1} \cdot \frac{3}{2} \cdot \frac{5}{4} \cdot \frac{9}{8} \cdot \frac{17}{16} \cdots$  converge?
- (b) The infinite product  $\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdot \frac{7}{8} \cdot \frac{9}{10} \cdots$  certainly converges. (Why?) Does it converge to zero?
- (c) In 1655, John Wallis famously derived the formula

$$\left(\frac{2 \cdot 2}{1 \cdot 3}\right) \left(\frac{4 \cdot 4}{3 \cdot 5}\right) \left(\frac{6 \cdot 6}{5 \cdot 7}\right) \left(\frac{8 \cdot 8}{7 \cdot 9}\right) \cdots = \frac{\pi}{2}$$

Show that the left side of this identity at least converges to something. (A complete proof of this result is taken up in Section 8.3.)

### Solution

- (a) Rewriting the terms as  $a_n = (1 + 1/n)$  and using the result from 2.4.10 implies the product diverges since  $\sum 1/n$  diverges.
- (b) Converges by the monotone convergence theorem since the partial products are decreasing and greater than zero.
- (c) In component form we have

$$a_n = \frac{(2n)^2}{(2n-1)(2n+1)} = \frac{(2n)^2}{(2n)^2 - 1} = 1 + \frac{(2n)^2 - ((2n)^2 - 1)}{(2n)^2 - 1} = 1 + \frac{1}{(2n)^2 - 1}$$

And since  $\sum 1/((2n)^2 - 1)$  converges by a comparison test with  $1/n^2$  2.4.10 implies

$$\prod_{n=1}^{\infty} \left(1 + \frac{1}{(2n)^2 - 1}\right)$$

also converges.

### Exercise 2.7.11

Find examples of two series  $\sum a_n$  and  $\sum b_n$  both of which diverge but for which  $\sum \min\{a_n, b_n\}$  converges. To make it more challenging, produce examples where  $(a_n)$  and  $(b_n)$  are strictly positive and decreasing.

### Solution

TODO

### Exercise 2.7.12 (Summation-by-parts)

Let  $(x_n)$  and  $(y_n)$  be sequences, let  $s_n = x_1 + x_2 + \cdots + x_n$  and set  $s_0 = 0$ . Use the observation that  $x_j = s_j - s_{j-1}$  to verify the formula

$$\sum_{j=m}^n x_j y_j = s_n y_{n+1} - s_{m-1} y_m + \sum_{j=m}^n s_j (y_j - y_{j+1})$$

**Solution**

Since  $x_j = s_j - s_{j-1}$  we can rewrite the sum as

$$\begin{aligned}
 &= \sum_{j=m}^n (s_j - s_{j-1}) y_j \\
 &= \sum_{j=m}^n s_j y_j - \sum_{j=m}^n s_{j-1} y_j \\
 &= (s_m y_m + \cdots + s_n y_n) - (s_{m-1} y_m + \cdots + s_{n-1} y_n) && \text{Factor out each } s_j \\
 &= s_n (y_n) - s_{m-1} (y_m) + s_m (y_m - y_{m+1}) + \cdots + s_{n-1} (y_{n-1} - y_n) && \text{Add and subtract } s_n y_{n+1} \text{ for factoring} \\
 &= s_n (y_{n+1}) - s_{m-1} (y_m) + s_m (y_m - y_{m+1}) + \cdots + s_n (y_n - y_{n+1}) && \text{Rewrite as a sum} \\
 &= s_n (y_{n+1}) - s_{m-1} (y_m) + \sum_{j=m}^n s_j (y_j - y_{j+1}) && \text{Done :) }
 \end{aligned}$$

**Exercise 2.7.13 (Abel's Test)**

Abel's Test for convergence states that if the series  $\sum_{k=1}^{\infty} x_k$  converges, and if  $(y_k)$  is a sequence satisfying

$$y_1 \geq y_2 \geq y_3 \geq \cdots \geq 0$$

then the series  $\sum_{k=1}^{\infty} x_k y_k$  converges.

(a) Use Exercise 2.7.12 to show that

$$\sum_{k=1}^n x_k y_k = s_n y_{n+1} + \sum_{k=1}^n s_k (y_k - y_{k+1})$$

where  $s_n = x_1 + x_2 + \cdots + x_n$ .

(b) Use the Comparison Test to argue that  $\sum_{k=1}^{\infty} s_k (y_k - y_{k+1})$  converges absolutely, and show how this leads directly to a proof of Abel's Test.

**Solution**

Intuitively, Abel's test is saying that multiplying by a positive decreasing sequence  $(y_n)$  is basically the same as multiplying by a constant in that it doesn't effect convergence.

(a) Exercise 2.7.12 combined with  $s_0 = 0$  gives

$$\sum_{k=1}^n x_k y_k = s_n y_{n+1} + \sum_{k=1}^n s_k (y_k - y_{k+1})$$

as desired.

(b)  $s_n y_{n+1}$  clearly converges since  $y_{n+1}$  is "eventually constant", so we must only show the right hand side converges.



We will show absolute convergence, note  $y_k - y_{k+1} \geq 0$  and so

$$\sum_{k=1}^n |s_k|(y_k - y_{k+1}) \geq 0$$

Bounding  $|s_n| \leq M$  gives

$$\sum_{k=1}^n |s_k|(y_k - y_{k+1}) \leq M \sum_{k=1}^n (y_k - y_{k+1})$$

Since  $\sum_{k=1}^n (y_k - y_{k+1}) = y_1 - y_{n+1}$  is telescoping we can write

$$\sum_{k=1}^n |s_k|(y_k - y_{k+1}) \leq M(y_1 - y_{n+1}) \leq My_1$$

Implying  $\sum_{k=1}^{\infty} |s_k|(y_k - y_{k+1})$  converges since it is bounded and increasing. And since the series converges absolutely so does the original  $\sum_{k=1}^{\infty} s_k(y_k - y_{k+1})$ .

**Summary:** Bound  $|s_k| \leq M$  and use the fact that  $\sum (y_k - y_{k+1})$  is telescoping.

### Exercise 2.7.14 (Dirichlet's Test)

Dirichlet's Test for convergence states that if the partial sums of  $\sum_{k=1}^{\infty} x_k$  are bounded (but not necessarily convergent), and if  $(y_k)$  is a sequence satisfying  $y_1 \geq y_2 \geq y_3 \geq \cdots \geq 0$  with  $\lim y_k = 0$ , then the series  $\sum_{k=1}^{\infty} x_k y_k$  converges.

- Point out how the hypothesis of Dirichlet's Test differs from that of Abel's Test in Exercise 2.7.13, but show that essentially the same strategy can be used to provide a proof.
- Show how the Alternating Series Test (Theorem 2.7.7) can be derived as a special case of Dirichlet's Test.

### Solution

- Abel's test assumed  $\sum x_n$  converged but did not require  $\lim y_k = 0$ . To prove it we use summation by parts to get

$$\sum_{k=1}^n x_k y_k = s_n y_{n+1} + \sum_{k=1}^n s_k (y_k - y_{k+1})$$

Clearly  $(s_n y_{n+1}) \rightarrow 0$  since  $|s_n| \leq M$  and  $(y_{n+1}) \rightarrow 0$ . Now we show absolute convergence of the right hand side (note  $|y_k - y_{k+1}| = (y_k - y_{k+1})$  as  $(y_n)$  is decreasing)

$$\sum_{k=1}^n |s_k|(y_k - y_{k+1}) \leq M \sum_{k=1}^n (y_k - y_{k+1}) = M(y_1 - y_{n+1}) \leq My_1$$

Thus  $\sum s_k(y_k - y_{k+1})$  converges absolutely and so finally

$$s_n y_{n+1} + \sum_{k=1}^n s_k (y_k - y_{k+1}) = \sum_{k=1}^n x_k y_k$$

converges.

- (b) Let  $a_n \geq 0$  with  $a_1 \geq a_2 \geq \cdots \geq 0$  and  $\lim a_n = 0$ . The series  $\sum (-1)^n \leq 1$  is bounded, so Dirichlet's test implies  $\sum (-1)^n a_n$  converges.

In fact any periodic  $b_n$  with  $\sum b_n$  bounded has  $\sum a_n b_n$  converging by Dirichlet.

# Chapter 3

## Basic Topology of $\mathbf{R}$

### 3.2 Open and Closed Sets

#### Exercise 3.2.1

- (a) Where in the proof of Theorem 3.2.3 part (ii) does the assumption that the collection of open sets be finite get used?
- (b) Give an example of a countable collection of open sets  $\{O_1, O_2, O_3, \dots\}$  whose intersection  $\bigcap_{n=1}^{\infty} O_n$  is closed, not empty and not all of  $\mathbf{R}$ .

#### Solution

- (a) Taking  $\min\{\epsilon_1, \dots, \epsilon_N\}$  is only possible for finite sets.
- (b)  $O_n = (-1/n, 1 + 1/n)$  has  $\bigcap_{n=1}^{\infty} O_n = [0, 1]$ .

#### Exercise 3.2.2

Let

$$A = \left\{ (-1)^n + \frac{2}{n} : n = 1, 2, 3, \dots \right\} \quad \text{and} \quad B = \{x \in \mathbf{Q} : 0 < x < 1\}$$

Answer the following questions for each set:

- (a) What are the limit points?
- (b) Is the set open? Closed?
- (c) Does the set contain any isolated points?
- (d) Find the closure of the set.

#### Solution

- (a) The set of  $B$ 's limit points is  $[0, 1]$ . The set of  $A$ 's limit points is  $\{1, -1\}$ .
- (b)  $B$  is not open since every  $(a, b) \not\subseteq B$  and  $B$  is not closed since we can construct limits to irrational values outside  $B$ .  $A$  is closed since  $\{1, -1\} \subseteq A$ , but not open as it does not contain any irrationals meaning  $(a, b) \not\subseteq A$  for all  $a, b \in \mathbf{R}$ .

- (c) Every point of  $A$  except the limit points  $\{1, -1\}$  is isolated, as if it were not isolated it would be a limit point.  $B$  has no isolated points since  $B \setminus [0, 1] = \emptyset$ , or in other words since  $B$  is dense in  $[0, 1]$  every  $b \in B \subseteq [0, 1]$  can be reached via a limit.
- (d)  $\overline{A} = A$  as  $A$  is already closed, and  $\overline{B} = B \cup [0, 1] = [0, 1]$ .

**Exercise 3.2.3**

Decide whether the following sets are open, closed, or neither. If a set is not open, find a point in the set for which there is no  $\epsilon$ -neighborhood contained in the set. If a set is not closed, find a limit point that is not contained in the set.

- (a)  $\mathbf{Q}$ .
- (b)  $\mathbf{N}$ .
- (c)  $\{x \in \mathbf{R} : x \neq 0\}$ .
- (d)  $\{1 + 1/4 + 1/9 + \cdots + 1/n^2 : n \in \mathbf{N}\}$
- (e)  $\{1 + 1/2 + 1/3 + \cdots + 1/n : n \in \mathbf{N}\}$

**Solution**

- (a) Neither, not open as  $(a, b) \subseteq \mathbf{Q}$  is impossible since  $\mathbf{Q}$  contains no irrationals but  $(a, b)$  does. and not closed since every irrational can be reached as a limit of rationals ( $\sqrt{2}$  is a simple example).
- (b) Clearly not open, but ironically closed since it has no limit points.
- (c) Open since every  $x \in \{x \in \mathbf{R} : x \neq 0\}$  has an  $\epsilon$ -neighborhood around it excluding zero. But closed since  $(1/n) \rightarrow 0$ .
- (d) Neither, not closed, as the limit  $\sum_k^n 1/n^2 = \pi^2/6$  is irrational but every term is rational. and not open as it does not contain any irrationals.
- (e) Closed as it has no limit points, every sequence diverges. Not open because it contains no irrationals.

**Exercise 3.2.4**

Let  $A$  be nonempty and bounded above so that  $s = \sup A$  exists.

- (a) Show that  $s \in \overline{A}$ .
- (b) Can an open set contain its supremum?

**Solution**

- (a) Since every  $s - \epsilon$  has an  $a \in A$  with  $a > s - \epsilon$  we can find  $a \in V_\epsilon(s)$  for any  $\epsilon > 0$ , meaning  $s$  is a limit point of  $A$  and thus contained in  $\overline{A}$ .
- (b) No, as  $(s, s + \epsilon)$  contains no  $a$ 's we have  $V_\epsilon(s) \not\subseteq A$  for every  $\epsilon > 0$ .

**Exercise 3.2.5**

Prove that a set  $F \subseteq \mathbf{R}$  is closed *if and only if* every Cauchy sequence contained in  $F$  has a limit that is also an element of  $F$ .

**Solution**

Let  $F \subseteq \mathbf{R}$  be closed and suppose  $(x_n)$  is a Cauchy sequence in  $F$ , since Cauchy sequences converge  $(x_n) \rightarrow x$  and finally since  $x \in F$  since  $F$  contains its limit points.

Now suppose every Cauchy sequence  $(x_n)$  in  $F$  converges to a limit in  $F$  and let  $l$  be a limit point of  $F$ , as  $l$  is a limit point of  $F$  there exists a sequence  $(y_n)$  in  $F$  with  $\lim(y_n) = l$ . since  $(y_n)$  converges it must be Cauchy, and since every Cauchy sequence converges to a limit inside  $F$  we have  $l \in F$ .

**Exercise 3.2.6**

Decide whether the following statements are true or false. Provide counterexamples for those that are false, and supply proofs for those that are true.

- (a) An open set that contains every rational number must necessarily be all of  $\mathbf{R}$ .
- (b) The Nested Interval Property remains true if the term “closed interval” is replaced by “closed set.”
- (c) Every nonempty open set contains a rational number.
- (d) Every bounded infinite closed set contains a rational number.
- (e) The Cantor set is closed.

**Solution**

- (a) False,  $A = (-\infty, \sqrt{2}) \cup (\sqrt{2}, \infty)$  contains every rational number but not  $\sqrt{2}$ .
- (b) False,  $C_n = [n, \infty)$  is closed, has  $C_{n+1} \subseteq C_n$  and  $C_n \neq \emptyset$  but  $\bigcap_{n=1}^{\infty} C_n = \emptyset$ .
- (c) True, let  $x \in A$  since  $A$  is open we have  $(a, b) \subseteq A$  with  $x \in (a, b)$  the density theorem implies there exists an  $r \in \mathbf{Q}$  with  $r \in (a, b)$  and thus  $r \in A$ .
- (d) False,  $A = \{1/n + \sqrt{2} : n \in \mathbf{N}\} \cup \{\sqrt{2}\}$  is closed and contains no rational numbers.
- (e) True, as it is the intersection of countably many closed intervals.

**Exercise 3.2.7**

Given  $A \subseteq \mathbf{R}$ , let  $L$  be the set of all limit points of  $A$ .

- (a) Show that the set  $L$  is closed.
- (b) Argue that if  $x$  is a limit point of  $A \cup L$ , then  $x$  is a limit point of  $A$ . Use this observation to furnish a proof for Theorem 3.2.12.

**Solution**

- (a) Every  $x_n \in L$  is  $x_n = \lim_{m \rightarrow \infty} a_{mn}$  for  $a_{mn} \in A$ . meaning if  $\lim x_n = x \notin L$  then for  $n > N$  and  $m > M$  we have

$$|a_{mn} - x| \leq |a_{mn} - x_n| + |x_n - x| < \epsilon/2 + \epsilon/2 = \epsilon$$

For  $n > \max\{N, M\}$  we get  $|a_{nn} - x| < \epsilon$  meaning  $x \in L$  since  $x$  is a limit point of  $A$ .

- (b) Let  $x_n \in A \cup L$  and  $x = \lim x_n$ . since  $x_n$  is infinite there must be at least one subsequence  $(x_{n_k}) \rightarrow x$  which is either all in  $A$  or all in  $L$ . If every  $x_{n_k} \in L$  then we know  $x \in L$  from (a), and if every  $x_{n_k} \in A$  then  $x \in L$  aswell.

### Exercise 3.2.8

Assume  $A$  is an open set and  $B$  is a closed set. Determine if the following sets are definitely open, definitely closed, both, or neither.

- (a)  $\overline{A \cup B}$
- (b)  $A \setminus B = \{x \in A : x \notin B\}$
- (c)  $(A^c \cup B)^c$
- (d)  $(A \cap B) \cup (A^c \cap B)$
- (e)  $\overline{A}^c \cap \overline{A}^c$

### Solution

For all of these keep in mind the only open and closed sets are  $\mathbf{R}$  and  $\emptyset$ , and if  $A$  is open  $A^c$  is closed and vice versa.

- (a) Closed, since the closure of a set is closed.
- (b) Open since  $B$  being closed implies  $B^c$  is open and thus  $A \cap B^c$  is open as it is an intersection of open sets.
- (c) Demorgan's laws give  $(A^c \cup B)^c = A \cap B^c$  which is the same as (b)
- (d) Both since  $(A \cap B) \cup (A^c \cap B) = \mathbf{R}$
- (e) Neither in general. Note that  $\overline{A}^c \neq \overline{A^c}$  consider how  $A = \{1/n : n \in \mathbf{N}\}$  has  $\overline{A^c} = \mathbf{R}$  but  $\overline{A}^c \neq \mathbf{R}$ .

### Exercise 3.2.9 (De Morgan's Laws)

A proof for De Morgan's Laws in the case of two sets is outlined in Exercise 1.2.5. The general argument is similar.

- (a) Given a collection of sets  $\{E_\lambda : \lambda \in \Lambda\}$ , show that

$$\left( \bigcup_{\lambda \in \Lambda} E_\lambda \right)^c = \bigcap_{\lambda \in \Lambda} E_\lambda^c \quad \text{and} \quad \left( \bigcap_{\lambda \in \Lambda} E_\lambda \right)^c = \bigcup_{\lambda \in \Lambda} E_\lambda^c$$

- (b) Now, provide the details for the proof of Theorem 3.2.14.

**Solution**

- (a) If  $x \in \left(\bigcup_{\lambda \in \Lambda} E_\lambda\right)^c$  then  $x \notin \bigcup_{\lambda \in \Lambda} E_\lambda$  meaning  $x \notin E_\lambda$  for all  $\lambda \in \Lambda$  implying  $x \in E_\lambda^c$  for all  $\lambda \in \Lambda$  and so finally  $x \in \bigcap_{\lambda \in \Lambda} E_\lambda^c$ . This shows

$$\left(\bigcup_{\lambda \in \Lambda} E_\lambda\right)^c \subseteq \bigcap_{\lambda \in \Lambda} E_\lambda^c$$

To show the reverse inclusion suppose  $x \in \bigcap_{\lambda \in \Lambda} E_\lambda^c$  then  $x \notin E_\lambda$  for all  $\lambda$  meaning  $x \notin \bigcup_{\lambda \in \Lambda} E_\lambda$  and so the reverse inclusion

$$\bigcap_{\lambda \in \Lambda} E_\lambda^c \subseteq \left(\bigcup_{\lambda \in \Lambda} E_\lambda\right)^c$$

Is true, completing the proof.

- (b) Let  $F = F_1 \cup F_2$ , if  $x_n \in F$  and  $x = \lim x_n$ . Let  $(x_{n_k})$  be a subsequence of  $(x_n)$  fully contained in  $F_1$  or  $F_2$ . the subsequence  $(x_{n_k})$  must also converge to  $x$ , meaning  $x$  is in  $F_1$  or  $F_2$ , the rest is by induction.

Now let  $F = \bigcap_{\lambda \in \Lambda} F_\lambda$

$$F^c = \bigcup_{\lambda \in \Lambda} F_\lambda^c$$

Each  $F_\lambda^c$  is open by Theorem 3.2.13, thus Theorem 3.2.3 (ii) implies  $F^c$  is open, and so  $(F^c)^c = F$  is closed.

**Exercise 3.2.10**

Only one of the following three descriptions can be realized. Provide an example that illustrates the viable description, and explain why the other two cannot exist.

- (i) A countable set contained in  $[0, 1]$  with no limit points.
- (ii) A countable set contained in  $[0, 1]$  with no isolated points.
- (iii) A set with an uncountable number of isolated points.

**Solution**

- (i) Cannot exist because taking any sequence  $(x_n)$  BW tells us there exists a convergent subsequence.
- (ii)  $\mathbf{Q} \cap [0, 1]$  is countable and has no isolated points.
- (iii) Impossible, let  $A \subseteq \mathbf{R}$  and let  $x$  be an isolated point of  $A$ . From the definition there exists a  $\delta > 0$  with  $V_\delta(x) \cap A = \{x\}$ . in Exercise 1.5.3 we proved there cannot exist an uncountable collection of disjoint open intervals, meaning we cannot have an uncountable set of isolated points as we can map them to open sets in a 1-1 fashion.

**Exercise 3.2.11**

- (a) Prove that  $\overline{A \cup B} = \overline{A} \cup \overline{B}$ .

(b) Does this result about closures extend to infinite unions of sets?

**Solution**

(a) Recall that the set of limit points of a set is closed (Exercise 3.2.7). Let  $L$  be the set of limit points of  $A \cup B$  and let  $L_a, L_b$  be the set of limit points for  $A$  and  $B$  respectively.

Let  $x \in L$ , thus there exists a sequence  $x_n \in A \cup B$  with  $x = \lim x_n$ , since  $(x_n)$  is infinite there exists a subsequence  $(x_{n_k})$  where every term is in  $A$  or  $B$ . Thus the limit  $\lim(x_{n_k}) = x$  must be a limit point of  $A$  or  $B$  meaning  $x \in L_a \cup L_b$ . This shows  $\overline{A \cup B} \subseteq \overline{A} \cup \overline{B}$ .

Now let  $x \in L_a$  ( $L_b$  is the same). there exists a sequence  $x_n \in A$  with  $x = \lim x_n$ , now since  $x_n \in A \cup B$  as well,  $x \in L$ . Thus we have shown  $\overline{A} \cup \overline{B} \subseteq \overline{A \cup B}$  completing the proof.

(b) False, take  $A_n = \{1/n\}$  as a counterexample

$$\bigcup_{n=1}^{\infty} A_n = \{1/n : n \in \mathbf{N}\} \cup \{0\}, \text{ but } \bigcup_{n=1}^{\infty} \overline{A_n} = \{1/n : n \in \mathbf{N}\}$$

**Exercise 3.2.12**

Let  $A$  be an uncountable set and let  $B$  be the set of real numbers that divides  $A$  into two uncountable sets; that is,  $s \in B$  if both  $\{x : x \in A \text{ and } x < s\}$  and  $\{x : x \in A \text{ and } x > s\}$  are uncountable. Show  $B$  is nonempty and open.

**Solution**

**TODO**

**Exercise 3.2.13**

Prove that the only sets that are both open and closed are  $\mathbf{R}$  and the empty set  $\emptyset$ .

**Solution**

**TODO**

**Exercise 3.2.14**

A dual notion to the closure of a set is the *interior* of a set. The interior of  $E$  is denoted  $E^\circ$  and is defined as

$$E^\circ = \{x \in E : \text{there exists } V_\epsilon(x) \subseteq E\}$$

Results about closures and interiors possess a useful symmetry.

(a) Show that  $E$  is closed if and only if  $\bar{E} = E$ . Show that  $E$  is open if and only if  $E^\circ = E$ .

(b) Show that  $\bar{E}^c = (E^c)^\circ$ , and similarly that  $(E^\circ)^c = \overline{E^c}$ .

**Solution**

(a) **TODO**

(b) **TODO**



**Exercise 3.2.15**

A set  $A$  is called an  $F_\sigma$  set if it can be written as the countable union of closed sets. A set  $B$  is called a  $G_\delta$  set if it can be written as the countable intersection of open sets.

- (a) Show that a closed interval  $[a, b]$  is a  $G_\delta$  set.
- (b) Show that the half-open interval  $(a, b]$  is both a  $G_\delta$  and an  $F_\sigma$  set.
- (c) Show that  $\mathbf{Q}$  is an  $F_\sigma$  set, and the set of irrationals  $\mathbf{I}$  forms a  $G_\delta$  set. (We will see in Section 3.5 that  $\mathbf{Q}$  is not a  $G_\delta$  set, nor is  $\mathbf{I}$  an  $F_\sigma$  set.)

**Solution**

- (a) **TODO**
- (b) **TODO**
- (c) **TODO**

### 3.3 Compact Sets

**Exercise 3.3.1**

Show that if  $K$  is compact and nonempty, then  $\sup K$  and  $\inf K$  both exist and are elements of  $K$ .

**Solution**

Let  $s = \sup K$ , since  $s$  is the least upper bound for every  $\epsilon > 0$  there exists an  $x \in K$  with  $s - \epsilon < x$ . Picking  $\epsilon_n = 1/n$  and  $x_n$  such that  $s - \epsilon_n < x_n$  we get that  $(x_n) \rightarrow s$  since  $(\epsilon_n) \rightarrow 0$ , and thus  $s \in K$ .

A similar argument applies to  $\inf K$ .

**Exercise 3.3.2**

Decide which of the following sets are compact. For those that are not compact, show how Definition 3.3.1 breaks down. In other words, give an example of a sequence contained in the given set that does not possess a subsequence converging to a limit in the set.

- (a)  $\mathbf{N}$ .
- (b)  $\mathbf{Q} \cap [0, 1]$ .
- (c) The Cantor set.
- (d)  $\{1 + 1/2^2 + 1/3^2 + \cdots + 1/n^2 : n \in \mathbf{N}\}$ .
- (e)  $\{1, 1/2, 2/3, 3/4, 4/5, \dots\}$ .

**Solution**

- (a) Not compact, the sequence  $x_n = n$  in  $\mathbf{N}$  has no convergent subsequence in  $\mathbf{N}$ .
- (b) Not compact, as we can construct a sequence  $(x_n) \rightarrow 1/\sqrt{2} \notin \mathbf{Q} \cap [0, 1]$  implying  $K$  is not closed, and thus cannot be compact.

- (c) Compact, since the cantor set is bounded and closed since it is the infinite intersection of closed sets  $\bigcap_{n=1}^{\infty} C_n$  where  $C_1 = [0, 1/3] \cup [2/3, 1]$  etc where you keep removing the middle thirds of each interval.
- (d) Not compact as every sequence  $(x_n)$  contained in the set converges to  $\pi^2/6$  which is not in the set, meaning the set isn't closed and thus cannot be compact.
- (e) Compact since it is bounded and closed, with every sequence in the set converging to one.

**Exercise 3.3.3**

Prove the converse of Theorem 3.3.4 by showing that if a set  $K \subseteq \mathbf{R}$  is closed and bounded, then it is compact.

**Solution**

Let  $K$  be closed and bounded and let  $(x_n)$  be a sequence contained in  $K$ . BW tells us a convergent subsequence  $(x_{n_k}) \rightarrow x$  exists since  $K$  is bounded, and since  $K$  is closed  $x \in K$ . Thus every sequence in  $K$  contains a subsequence converging to a limit in  $K$ , which is the definition of  $K$  being compact.

**Exercise 3.3.4**

Assume  $K$  is compact and  $F$  is closed. Decide if the following sets are definitely compact, definitely closed, both, or neither.

- (a)  $K \cap F$
- (b)  $\overline{F^c \cup K^c}$
- (c)  $K \setminus F = \{x \in K : x \notin F\}$
- (d)  $\overline{K \cap F^c}$

**Solution**

- (a) Compact since  $K \cap F$  is closed (finite intersection of closed sets) and bounded (since  $K$  is bounded)
- (b) Closed but not Compact since  $K$  being bounded implies  $K^c$  is unbounded, meaning  $\overline{F^c \cup K^c}$  is unbounded.
- (c)  $K \setminus F = K \cap F^c$  could be either, if  $K = [0, 1]$ ,  $F^c = (0, 1)$  then  $K \cap F^c$  is open, but if  $K = [0, 1]$  and  $F^c = (-1, 2)$  then  $K \cap F^c = [0, 1]$  is compact.
- (d) Compact since  $K \cap F^c$  is bounded (since  $K$  is bounded) implies  $\overline{K \cap F^c}$  is closed (closure of a set is closed) and bounded (if  $A$  is bounded then  $\overline{A}$  is also bounded).

**Exercise 3.3.5**

Decide whether the following propositions are true or false. If the claim is valid, supply a short proof, and if the claim is false, provide a counterexample.

- (a) The arbitrary intersection of compact sets is compact.

- (b) The arbitrary union of compact sets is compact.
- (c) Let  $A$  be arbitrary, and let  $K$  be compact. Then, the intersection  $A \cap K$  is compact.
- (d) If  $F_1 \supseteq F_2 \supseteq F_3 \supseteq F_4 \supseteq \cdots$  is a nested sequence of nonempty closed sets, then the intersection  $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$ .

**Solution**

- (a) True, as it will be bounded and closed (since arbitrary intersections of closed sets are closed).
- (b) False,  $\bigcup_{n=1}^{\infty} [0, n]$  is unbounded and thus not compact.
- (c) False, let  $K = [0, 1]$  and  $A = (0, 1)$ . The intersection  $K \cap A = (0, 1)$  is not compact.
- (d) False as  $\bigcap_{n=1}^{\infty} [n, \infty) = \emptyset$  (It is true for compact sets though)

**Exercise 3.3.6**

This exercise is meant to illustrate the point made in the opening paragraph to Section 3.3. Verify that the following three statements are true if every blank is filled in with the word “finite.” Which are true if every blank is filled in with the word “compact”? Which are true if every blank is filled in with the word “closed”?

- (a) Every \_\_\_\_\_ set has a maximum.
- (b) If  $A$  and  $B$  are \_\_\_\_\_, then  $A + B = \{a + b : a \in A, b \in B\}$  is also \_\_\_\_\_
- (c) If  $\{A_n : n \in \mathbf{N}\}$  is a collection of \_\_\_\_\_ sets with the property that every finite subcollection has a nonempty intersection, then  $\bigcap_{n=1}^{\infty} A_n$  is nonempty as well.

**Solution**

- (a) Compact
- (b) Closed, since if  $a + b \in A + B$  then we can find sequences  $(a_n) \rightarrow a$  and  $(b_n) \rightarrow b$  in  $A$  and  $B$  respectively, with  $(a_n + b_n)$  contained in  $A + B$  and  $(a_n + b_n) \rightarrow a + b$ .
- (c) Compact, since letting  $K_n = \bigcap_{k=1}^n A_k$  gives  $K_n \subseteq K_{n-1}$ , we also have  $K_n \neq \emptyset$  since every finite intersection is known to be nonempty. Applying the Nested Compact Set Property allows us to conclude

$$\bigcap_{n=1}^{\infty} A_n = \bigcap_{n=1}^{\infty} K_n \neq \emptyset$$

**Exercise 3.3.7**

As some more evidence of the surprising nature of the Cantor set, follow these steps to show that the sum  $C + C = \{x + y : x, y \in C\}$  is equal to the closed interval  $[0, 2]$ . (Keep in mind that  $C$  has zero length and contains no intervals.)

Because  $C \subseteq [0, 1]$ ,  $C + C \subseteq [0, 2]$ , so we only need to prove the reverse inclusion  $[0, 2] \subseteq \{x + y : x, y \in C\}$ . Thus, given  $s \in [0, 2]$ , we must find two elements  $x, y \in C$  satisfying  $x + y = s$

- (a) Show that there exist  $x_1, y_1 \in C_1$  for which  $x_1 + y_1 = s$ . Show in general that, for an arbitrary  $n \in \mathbf{N}$ , we can always find  $x_n, y_n \in C_n$  for which  $x_n + y_n = s$ .
- (b) Keeping in mind that the sequences  $(x_n)$  and  $(y_n)$  do not necessarily converge, show how they can nevertheless be used to produce the desired  $x$  and  $y$  in  $C$  satisfying  $x + y = s$ .

**Solution**

- (a) Since  $y = s - x$  we must show there exists an  $x \in C_1$  such that  $s - x \in C_1$ . If  $s \in C_1$  then set  $x = 0$ , if  $s \notin C_1$  then either  $s \in (1/3, 2/3)$  and  $x = 1/3$  or  $s \in (1, 2]$  and  $x = 1$ .

**TODO**

- (b) Since  $C$  is compact, there exists a subsequence  $(x_{n_k}) \rightarrow x$  with  $x \in C$ . Now since  $x_{n_k} + y_{n_k} = s$  for all  $k$ , we have  $\lim y_{n_k} = \lim s - x_{n_k} = s - x$ . Now since each  $y_{n_k} \in C$  the limit  $y = s - x \in C$  as well, thus we have found  $x, y \in C$  with  $x + y = s$ .

**Exercise 3.3.8**

Let  $K$  and  $L$  be nonempty compact sets, and define

$$d = \inf\{|x - y| : x \in K \text{ and } y \in L\}$$

This turns out to be a reasonable definition for the distance between  $K$  and  $L$ .

- (a) If  $K$  and  $L$  are disjoint, show  $d > 0$  and that  $d = |x_0 - y_0|$  for some  $x_0 \in K$  and  $y_0 \in L$ .
- (b) Show that it's possible to have  $d = 0$  if we assume only that the disjoint sets  $K$  and  $L$  are closed.

**Solution**

- (a) The set  $|K - L| = \{|x - y| : x \in K, y \in L\}$  is compact since  $K - L$  is compact by 3.3.6 (b) and  $|\cdot|$  preserves compactness. Thus  $d = \inf |K - L|$  has  $d = |x_0 - y_0|$  for some  $x_0 \in K$  and  $y_0 \in L$ .
- (b)  $K = \{n : n \in \mathbf{N}\}$  and  $L = \{n + 1/n : n \in \mathbf{N}\}$  have  $d = 0$ , and both are closed since every limit diverges.

**Exercise 3.3.9**

Follow these steps to prove that being compact implies every open cover has a finite subcover.

Assume  $K$  is compact, and let  $\{O_\lambda : \lambda \in \Lambda\}$  be an open cover for  $K$ . For contradiction, let's assume that no finite subcover exists. Let  $I_0$  be a closed interval containing  $K$ .

- (a) Show that there exists a nested sequence of closed intervals  $I_0 \supseteq I_1 \supseteq I_2 \supseteq \cdots$  with the property that, for each  $n$ ,  $I_n \cap K$  cannot be finitely covered and  $\lim |I_n| = 0$ .
- (b) Argue that there exists an  $x \in K$  such that  $x \in I_n$  for all  $n$ .
- (c) Because  $x \in K$ , there must exist an open set  $O_{\lambda_0}$  from the original collection that contains  $x$  as an element. Explain how this leads to the desired contradiction.

**Solution**

- (a) Bisect  $I_0$  into two intervals, and let  $I_1$  be the interval where  $I_1 \cap K$  cannot be finitely covered. Repeating in this fashion we have  $\lim |I_n| = \lim |I_0|(1/2)^n = 0$ .
- (b) The nested compact set property with  $K_n = I_n \cap K$  gives  $x \in \bigcap_{n=1}^{\infty} K_n$  meaning  $x \in K$  and  $x \in I_n$  for all  $n$ .
- (c) Since  $x \in O_{\lambda_0}$  and  $|I_n| \rightarrow 0$  with  $x \in I_n$  for all  $n$ , there exists an  $N$  where  $n > N$  implies  $I_n \subseteq O_{\lambda_0}$  contradicting the assumption that  $I_n \cap K$  cannot be finitely covered since  $\{O_{\lambda_0}\}$  is a finite subcover for  $I_n \cap K$ .

**Exercise 3.3.10**

Here is an alternate proof to the one given in Exercise 3.3.9 for the final implication in the Heine-Borel Theorem.

Consider the special case where  $K$  is a closed interval. Let  $\{O_\lambda : \lambda \in \Lambda\}$  be an open cover for  $[a, b]$  and define  $S$  to be the set of all  $x \in [a, b]$  such that  $[a, x]$  has a finite subcover from  $\{O_\lambda : \lambda \in \Lambda\}$ .

- (a) Argue that  $S$  is nonempty and bounded, and thus  $s = \sup S$  exists.
- (b) Now show  $s = b$ , which implies  $[a, b]$  has a finite subcover.
- (c) Finally, prove the theorem for an arbitrary closed and bounded set  $K$ .

**Solution**

- (a) **TODO**
- (b) **TODO**
- (c) **TODO**

**Exercise 3.3.11**

Consider each of the sets listed in Exercise 3.3.2. For each one that is not compact, find an open cover for which there is no finite subcover.

**Solution**

- (a)  $\mathbf{N}$  and  $\{V_1(n) : n \in \mathbf{N}\}$  has no finite subcover since each  $V_1(n)$  covers exactly one  $n \in \mathbf{N}$ , meaning there are no subcovers at all!
- (b)  $\mathbf{Q} \cap [0, 1]$  and  $\{O(x) : x \in \mathbf{Q} \cap [0, 1]\}$  where  $O(x) = V_{\frac{1}{2}|x-y|}(x)$  for  $y \in \mathbf{Q}^c \cap [0, 1]$  since any finite subcover  $\{O(x_1), \dots, O(x_n)\}$  has  $V_\epsilon(y) \not\subseteq O(x_i)$  for all  $i$  when  $\epsilon = \min_i \{\frac{1}{2}|y - x_i|\}$ . Thus the density theorem gives an  $x \in V_\epsilon(y)$  with  $x \in \mathbf{Q} \cap [0, 1]$  where  $x$  is not in the finite cover. **TODO** clean this up
- (c) The Cantor is compact
- (d)  $K = \{1 + 1/2^2 + 1/3^2 + \dots + 1/n^2 : n \in \mathbf{N}\}$  and  $\{V_{\frac{1}{2}|x-L|}(x) : x \in K\}$  for  $L = \pi^2/6$  since any finite cover  $\{V_{\frac{1}{2}|x_1-L|}(x_1), \dots, V_{\frac{1}{2}|x_n-L|}(x_n)\}$ , letting  $\epsilon = \min\{\frac{1}{2}|x_i - L|\}$  will make  $V_\epsilon(L)$  not in the finite cover, meaning there exists an  $x \in V_\epsilon(L)$  with  $x \in K$  (since  $K$  gets arbitrarily close to  $L$ ) but  $x$  not in the finite cover.

(e)  $\{1, 1/2, 2/3, 3/4, 4/5, \dots\}$  is compact

### Exercise 3.3.12

Using the concept of open covers (and explicitly avoiding the Bolzano-Weierstrass Theorem), prove that every bounded infinite set has a limit point.

**Solution**

**TODO**

### Exercise 3.3.13

Let's call a set *clompact* if it has the property that every *closed* cover (i.e., a cover consisting of closed sets) admits a finite subcover. Describe all of the clompact subsets of  $\mathbf{R}$ .

**Solution**

$K$  is clompact if and only if  $K$  is finite, since the closed cover  $\{[x, x] : x \in K\} = K$  having a finite subcover implies  $\{[x, x] : x \in K\}$  is finite (since it is the only subcover that works) therefor  $K$  is finite. If  $K$  is finite then it obviously permits a finite subcover.

## 3.4 Perfect Sets and Connected Sets

### Exercise 3.4.1

If  $P$  is a perfect set and  $K$  is compact, is the intersection  $P \cap K$  always compact? Always perfect?

**Solution**

Recall a perfect set is a closed set with no isolated points. Thus the intersection of a closed set  $P$  and a closed bounded set  $K$  gives a closed bounded (and thus compact) set  $P \cap K$ .

Now take  $P = \mathbf{R}$ , we get  $P \cap K = K$  which is not necessarily perfect.

### Exercise 3.4.2

Does there exist a perfect set consisting of only rational numbers?

**Solution**

**TODO**

### Exercise 3.4.3

Review the portion of the proof given in Example 3.4.2 and follow these steps to complete the argument.

- (a) Because  $x \in C_1$ , argue that there exists an  $x_1 \in C \cap C_1$  with  $x_1 \neq x$  satisfying  $|x - x_1| \leq 1/3$ .
- (b) Finish the proof by showing that for each  $n \in \mathbf{N}$ , there exists  $x_n \in C \cap C_n$ , different from  $x$ , satisfying  $|x - x_n| \leq 1/3^n$ .

**Solution**

- (a) **TODO**

(b) **TODO**

#### Exercise 3.4.4

Repeat the Cantor construction from Section 3.1 starting with the interval  $[0, 1]$ . This time, however, remove the open middle *fourth* from each component.

(a) Is the resulting set compact? Perfect?

(b) Using the algorithms from Section 3.1, compute the length and dimension of this Cantor-like set.

#### Solution

(a) **TODO**

(b) **TODO**

#### Exercise 3.4.5

Let  $A$  and  $B$  be nonempty subsets of  $\mathbf{R}$ . Show that if there exist disjoint open sets  $U$  and  $V$  with  $A \subseteq U$  and  $B \subseteq V$ , then  $A$  and  $B$  are separated.

#### Solution

**TODO**

#### Exercise 3.4.6

Prove that a set  $E \subseteq \mathbf{R}$  is connected if and only if, for all nonempty disjoint sets  $A$  and  $B$  satisfying  $E = A \cup B$ , there always exists a convergent sequence  $(x_n) \rightarrow x$  with  $(x_n)$  contained in one of  $A$  or  $B$ , and  $x$  an element of the other. (Theorem 3.4.6)

#### Solution

**TODO**

#### Exercise 3.4.7

A set  $E$  is totally disconnected if, given any two distinct points  $x, y \in E$ , there exist separated sets  $A$  and  $B$  with  $x \in A, y \in B$ , and  $E = A \cup B$ .

(a) Show that  $\mathbf{Q}$  is totally disconnected.

(b) Is the set of irrational numbers totally disconnected?

#### Solution

(a) **TODO**

(b) **TODO**

#### Exercise 3.4.8

Follow these steps to show that the Cantor set is totally disconnected in the sense described in Exercise 3.4.7. Let  $C = \bigcap_{n=0}^{\infty} C_n$ , as defined in Section 3.1.

(a) Given  $x, y \in C$ , with  $x < y$ , set  $\epsilon = y - x$ . For each  $n = 0, 1, 2, \dots$ , the set  $C_n$  consists of a finite number of closed intervals. Explain why there must exist an  $N$  large enough so that it is impossible for  $x$  and  $y$  both to belong to the same closed interval of  $C_N$ .

- (b) Show that  $C$  is totally disconnected.

**Solution**

(a) **TODO**

(b) **TODO**

**Exercise 3.4.9**

Let  $\{r_1, r_2, r_3, \dots\}$  be an enumeration of the rational numbers, and for each  $n \in \mathbb{N}$  set  $\epsilon_n = 1/2^n$ . Define  $O = \bigcup_{n=1}^{\infty} V_{\epsilon_n}(r_n)$ , and let  $F = O^c$ .

- (a) Argue that  $F$  is a closed, nonempty set consisting only of irrational numbers.
- (b) Does  $F$  contain any nonempty open intervals? Is  $F$  totally disconnected? (See Exercise 3.4.7 for the definition.)
- (c) Is it possible to know whether  $F$  is perfect? If not, can we modify this construction to produce a nonempty perfect set of irrational numbers?

**Solution**

(a) **TODO**

(b) **TODO**

(c) **TODO**



# Chapter 4

## Functional Limits

### 4.2 Functional Limits

#### Exercise 4.2.1

- (a) Supply the details for how Corollary 4.2.4 part (ii) follows from the Sequential Criterion for Functional Limits in Theorem 4.2.3 and the Algebraic Limit Theorem for sequences proved in Chapter 2.
- (b) Now, write another proof of Corollary 4.2.4 part (ii) directly from Definition 4.2.1 without using the sequential criterion in Theorem 4.2.3.
- (c) Repeat (a) and (b) for Corollary 4.2.4 part (iii).

#### Solution

- (a) **TODO**
- (b) **TODO**
- (c) **TODO**

#### Exercise 4.2.2

For each stated limit, find the largest possible  $\delta$ -neighborhood that is a proper response to the given  $\epsilon$  challenge.

- (a)  $\lim_{x \rightarrow 3} (5x - 6) = 9$ , where  $\epsilon = 1$ .
- (b)  $\lim_{x \rightarrow 4} \sqrt{x} = 2$ , where  $\epsilon = 1$ .
- (c)  $\lim_{x \rightarrow \pi} \lceil x \rceil = 3$ , where  $\epsilon = 1$ . (The function  $\lceil x \rceil$  returns the greatest integer less than or equal to  $x$ .)
- (d)  $\lim_{x \rightarrow \pi} \lfloor x \rfloor = 3$ , where  $\epsilon = .01$ .

#### Solution

- (a) **TODO**
- (b) **TODO**

(c) **TODO**(d) **TODO****Exercise 4.2.3**

Review the definition of Thomae's function  $t(x)$  from Section 4.1.

- (a) Construct three different sequences  $(x_n)$ ,  $(y_n)$ , and  $(z_n)$ , each of which converges to 1 without using the number 1 as a term in the sequence.
- (b) Now, compute  $\lim t(x_n)$ ,  $\lim t(y_n)$ , and  $\lim t(z_n)$ .
- (c) Make an educated conjecture for  $\lim_{x \rightarrow 1} t(x)$ , and use Definition 4.2.1 B to verify the claim. (Given  $\epsilon > 0$ , consider the set of points  $\{x \in \mathbf{R} : t(x) \geq \epsilon\}$ . Argue that all the points in this set are isolated.)

**Solution**(a) **TODO**(b) **TODO**(c) **TODO****Exercise 4.2.4**

Consider the reasonable but erroneous claim that

$$\lim_{x \rightarrow 10} 1/[x] = 1/10$$

- (a) Find the largest  $\delta$  that represents a proper response to the challenge of  $\epsilon = 1/2$
- (b) Find the largest  $\delta$  that represents a proper response to  $\epsilon = 1/50$ .
- (c) Find the largest  $\epsilon$  challenge for which there is no suitable  $\delta$  response possible.

**Solution**(a) **TODO**(b) **TODO**(c) **TODO****Exercise 4.2.5**

Use Definition 4.2.1 to supply a proper proof for the following limit statements.

- (a)  $\lim_{x \rightarrow 2} (3x + 4) = 10$
- (b)  $\lim_{x \rightarrow 0} x^3 = 0$
- (c)  $\lim_{x \rightarrow 2} (x^2 + x - 1) = 5$ .
- (d)  $\lim_{x \rightarrow 3} 1/x = 1/3$

**Solution**

- (a) **TODO**
- (b) **TODO**
- (c) **TODO**
- (d) **TODO**

**Exercise 4.2.6**

Decide if the following claims are true or false, and give short justifications for each conclusion.

- (a) If a particular  $\delta$  has been constructed as a suitable response to a particular  $\epsilon$  challenge, then any smaller positive  $\delta$  will also suffice.
- (b) If  $\lim_{x \rightarrow a} f(x) = L$  and  $a$  happens to be in the domain of  $f$ , then  $L = f(a)$
- (c) If  $\lim_{x \rightarrow a} f(x) = L$ , then  $\lim_{x \rightarrow a} 3[f(x) - 2]^2 = 3(L - 2)^2$
- (d) If  $\lim_{x \rightarrow a} f(x) = 0$ , then  $\lim_{x \rightarrow a} f(x)g(x) = 0$  for any function  $g$  (with domain equal to the domain of  $f$ .)

**Solution**

- (a) **TODO**
- (b) **TODO**
- (c) **TODO**
- (d) **TODO**

**Exercise 4.2.7**

Let  $g : A \rightarrow \mathbf{R}$  and assume that  $f$  is a bounded function on  $A$  in the sense that there exists  $M > 0$  satisfying  $|f(x)| \leq M$  for all  $x \in A$ . Show that if  $\lim_{x \rightarrow c} g(x) = 0$ , then  $\lim_{x \rightarrow c} g(x)f(x) = 0$  as well.

**Solution**

**TODO**

**Exercise 4.2.8**

Compute each limit or state that it does not exist. Use the tools developed in this section to justify each conclusion.

- (a)  $\lim_{x \rightarrow 2} \frac{|x-2|}{x-2}$
- (b)  $\lim_{x \rightarrow 7/4} \frac{|x-2|}{x-2}$
- (c)  $\lim_{x \rightarrow 0} (-1)^{[1/x]}$
- (d)  $\lim_{x \rightarrow 0} \sqrt[3]{x}(-1)^{[1/x]}$

**Solution**(a) **TODO**(b) **TODO**(c) **TODO**(d) **TODO**