

# Project 2 in FYS3150

Bendik Steinsvåg Dalen, Ulrik Seip

October 3, 2018

## 1 ABSTRACT

In this project we have coded a implementation of Jacobi's rotation algorithm, that finds an approximation to differential equations. We then then used this to model a harmonic oscillator problem in three dimensions, with one and two electrons.

## 2 INTRODUCTION

## 3 METHOD

### 3.a Mathematical basis

### 3.b Basic code, eller noe

### 3.c Testing the code

### 3.d Quantum dots in three dimensions, one electron

Now that we had a general algorithm we used it to model a electron that moves in a three-dimensional harmonic oscillator potential. In other words, we looked for the solution of the radial part of Schroedinger's equation for one electron, which reads

$$-\frac{\hbar^2}{2m} \left( \frac{1}{r^2} \frac{d}{dr} r^2 \frac{d}{dr} - \frac{l(l+1)}{r^2} \right) R(r) + V(r)R(r) = ER(r). \quad (1)$$

This problem also has analytical solutions, so we can test how accurate our algorithm is.  
(Some math-stuff).

The Schroedinger's equation then becomes

$$-\frac{d^2}{d\rho^2} u(\rho) + \rho^2 u(\rho) = \lambda y(\rho). \quad (2)$$

Since we are working in radial coordinates we have  $\rho \in [0, \infty)$ . Since we can't represent infinity on a computer we have to find an approximation, which we will come back to later. For now we define  $\rho_{min} = 0$  and  $\rho_{max}$  to represent the minimum and maximum values of  $\rho$ .

Function 2 is an differential equation that can be modeled similarly to (ting). If we have  $n$  mesh points we get a step length

$$h = \frac{\rho_{max} - \rho_{min}}{n}. \quad (3)$$

### 3.e Quantum dots in three dimensions, two electrons

## 4 RESULTS

## 5 CONCLUSIONS

## 6 APENDICES

## 7 REFERENCES