



COMP9020

Foundations of Computer Science
Term 3, 2024

Lecture 8: Boolean Logic

Outline

What is logic?

Boolean Logic

Boolean Functions

Conjunctive and Disjunctive Normal Form

Karnaugh Maps

Boolean Algebras

Outline

What is logic?

Boolean Logic

Boolean Functions

Conjunctive and Disjunctive Normal Form

Karnaugh Maps

Boolean Algebras

What is logic?

Logic is about **formalizing reasoning** and **defining truth**

- Adding rigour
- Removing ambiguity
- Mechanizing the process of reasoning

Loose history of logic

- (Ancient times): Logic exclusive to philosophy
- Mid-19th Century: Logical foundations of Mathematics (Boole, Jevons, Schröder, etc)
- 1910: Russell and Whitehead's Principia Mathematica
- 1928: Hilbert proposes *Entscheidungsproblem*
- 1931: Gödel's Incompleteness Theorem
- 1935: Church's Lambda calculus
- 1936: Turing's Machine-based approach
- 1930s: Shannon develops Circuit logic
- 1960s: Formal verification; Relational databases

Logic in Computer Science

Computation = Calculation + Symbolic manipulation

Logic in Computer Science

Computation = Calculation + Symbolic manipulation

Logic as 2-valued computation (Boolean logic):

- Circuit design
- Code optimization
- Boolean algebra
- Nand game

Logic in Computer Science

Computation = Calculation + Symbolic manipulation

Logic as symbolic reasoning (Propositional logic, and beyond):

- Formal verification
- Proof assistance
- Knowledge Representation and Reasoning
- Automated reasoning
- Databases

Outline

What is logic?

Boolean Logic

Boolean Functions

Conjunctive and Disjunctive Normal Form

Karnaugh Maps

Boolean Algebras

Boolean logic

Boolean logic is about performing calculations in a “simple” mathematical structure.

- complex calculations can be built entirely from these simple ones
- can help identify simplifications that improve performance at the circuit level
- can help identify simplifications that improve presentation at the programming level

The Boolean Algebra \mathbb{B}

Definition

The (two-element) **Boolean algebra** is defined to be the set $\mathbb{B} = \{0, 1\}$, together with the functions $! : \mathbb{B} \rightarrow \mathbb{B}$, $\&\& : \mathbb{B}^2 \rightarrow \mathbb{B}$, and $\| : \mathbb{B}^2 \rightarrow \mathbb{B}$, defined as follows:

$$!x = (1 - x) \quad x \&\& y = \min\{x, y\} \quad x \| y = \max\{x, y\}$$

Alternative notation

Commonly, the following alternative notation is used:

For \mathbb{B} : $\{\text{false}, \text{true}\}, \{F, T\}, \{0, 1\}, \{\perp, \top\}$

For $\neg x$: $\bar{x}, x', \sim x, \neg x, \text{NOT}(x)$

For $x \&\& y$: $xy, x \wedge y, (x \text{ AND } y)$

For $x \parallel y$: $x + y, x \vee y, (x \text{ OR } y)$

The Boolean Algebra \mathbb{B} – Alternative definition

Definition

The (two-element) **Boolean algebra** is defined to be the set $\mathbb{B} = \{\text{false}, \text{true}\}$, together with the functions $! : \mathbb{B} \rightarrow \mathbb{B}$, $\&\& : \mathbb{B}^2 \rightarrow \mathbb{B}$, and $\| : \mathbb{B}^2 \rightarrow \mathbb{B}$, defined as follows:

x	$!x$	x	y	$x \&\& y$	x	y	$x \ y$
		false	false	false	false	false	false
false	true	false	true	false	false	true	true
true	false	true	false	false	true	false	true
		true	true	true	true	true	true

Properties

We observe that $!$, $\&\&$, and \parallel satisfy the following:

For all $x, y, z \in \mathbb{B}$:

Commutativity

$$\begin{aligned}x \parallel y &= y \parallel x \\x \&\& y &= y \&\& x\end{aligned}$$

Associativity

$$\begin{aligned}(x \parallel y) \parallel z &= x \parallel (y \parallel z) \\(x \&\& y) \&\& z &= x \&\& (y \&\& z)\end{aligned}$$

Distribution

$$\begin{aligned}x \parallel (y \&\& z) &= (x \parallel y) \&\& (x \parallel z) \\x \&\& (y \parallel z) &= (x \&\& y) \parallel (x \&\& z)\end{aligned}$$

Identity

$$\begin{aligned}x \parallel 0 &= x \\x \&\& 1 &= x\end{aligned}$$

Complementation

$$\begin{aligned}x \parallel (!x) &= 1 \\x \&\& (!x) &= 0\end{aligned}$$

Examples

Examples

- Calculate $x \ \&\& \ x$ for all $x \in \mathbb{B}$
- Calculate $((1 \ \&\& \ 0) \parallel ((!1) \ \&\& \ (!0)))$

Outline

What is logic?

Boolean Logic

Boolean Functions

Conjunctive and Disjunctive Normal Form

Karnaugh Maps

Boolean Algebras

Boolean Functions

Definition

An **n -ary Boolean function** is a map $f : \mathbb{B}^n \rightarrow \mathbb{B}$.

Question

How many unary Boolean functions are there?

How many binary functions?

n -ary?

Examples

Examples

- $!$ is a unary Boolean function
- $\&\&$, \parallel are binary Boolean functions
- $f(x, y) = !(x \&\& y)$ is a binary boolean function (NAND)
- $\text{AND}(x_0, x_1, \dots) = (\dots((x_0 \&\& x_1) \&\& x_2) \dots)$ is a (family) of Boolean functions
- $\text{OR}(x_0, x_1, \dots) = (\dots((x_0 \parallel x_1) \parallel x_2) \dots)$ is a (family) of Boolean functions

Application: Adding two one-bit numbers

Question. How can we implement:

$$\text{add} : \mathbb{B}^2 \rightarrow \mathbb{B}^2$$

defined as

x	y	$\text{add}(x, y)$
0	0	00
0	1	01
1	0	01
1	1	10

Take Notice

Observe that this is **not** a Boolean function. Boolean functions can output either 0 or 1 (a single bit). This function outputs 2 bits.

(Short) Answer. Use two Boolean functions!

Take Notice

Digital circuits are just sequences of Boolean functions.

Outline

What is logic?

Boolean Logic

Boolean Functions

Conjunctive and Disjunctive Normal Form

Karnaugh Maps

Boolean Algebras

Conjunctive and Disjunctive normal form

Definition

- A **literal** is a unary Boolean function
- A **minterm** is a Boolean function of the form $\text{AND}(l_1(x_1), l_2(x_2), \dots, l_n(x_n))$ where the l_i are literals
- A **maxterm** is a Boolean function of the form $\text{OR}(l_1(x_1), l_2(x_2), \dots, l_n(x_n))$ where the l_i are literals
- A **CNF Boolean function** is a function of the form $\text{AND}(m_1, m_2, \dots)$, where the m_i are maxterms.
- A **DNF Boolean function** is a function of the form $\text{OR}(m_1, m_2, \dots)$, where the m_i are minterms.

Examples

Examples

Question. Are these functions in CNF? Are they in DNF?

- $f(x, y, z) = (x \ \&\& \ (!y) \ \&\& \ z) \ || \ (x \ \&\& \ (!y) \ \&\& \ (!z)) = x \ \bar{y} \ z + x \ \bar{y} \ \bar{z}$: DNF , but not CNF
- $g(x, y, z) = (x \ || \ (!y) \ || \ z) \ \&\& \ (x \ || \ (!y) \ || \ (!z)) = (x + \bar{y} + z)(x + \bar{y} + \bar{z})$: CNF function, but not DNF
- $h(x, y, z) = (x \ \&\& \ (!y) \ \&\& \ z) = x \ \bar{y} \ z$: both CNF and DNF
- $j(x, y, z) = x + y(z + x)$: Neither CNF nor DNF

Take Notice

CNF: product of sums; DNF: sum of products

Theorem

Every Boolean function can be written as a function in DNF.

Theorem

Every Boolean function can be written as a function in CNF.

Proof...

Canonical DNF

Given an n -ary Boolean function $f : \mathbb{B}^n \rightarrow \mathbb{B}$ we construct an equivalent DNF Boolean function as follows:

For each $\mathbf{b} = (b_1, \dots, b_n) \in \mathbb{B}^n$ we define the minterm

$$m_{\mathbf{b}} = \text{AND}(l_1(x_1), l_2(x_2), \dots, l_n(x_n))$$

where

$$l_i(x_i) = \begin{cases} x_i & \text{if } b_i = 1 \\ \neg x_i & \text{if } b_i = 0 \end{cases}$$

We then define the DNF formula:

$$f_{\text{DNF}} = \sum_{f(\mathbf{b})=1} m_{\mathbf{b}},$$

that is, f_{DNF} is the disjunction (or) over all minterms corresponding to elements $\mathbf{b} \in \mathbb{B}$ where $f(\mathbf{b}) = 1$.

Canonical DNF

Theorem

f and f_{DNF} are the same function.

Exercise

Exercises

RW: 10.2.3 Find the canonical DNF form of each of the following expressions in variables x, y, z

- xy
- \bar{z}
- $xy + \bar{z}$
- $f(x, y, z) = 1$

Outline

What is logic?

Boolean Logic

Boolean Functions

Conjunctive and Disjunctive Normal Form

Karnaugh Maps

Boolean Algebras

Karnaugh Maps

For up to four variables (propositional symbols) a diagrammatic method of simplification called **Karnaugh maps** works quite well.

- For every propositional function of $k = 2, 3, 4$ variables we construct a rectangular array of 2^k cells.
- Column labels and row labels are ordered by **Gray code**.
- Squares corresponding to the value **true** are marked with eg “+”.
- We try to cover these squares with as few rectangles with sides 1 or 2 or 4 as possible.

Example

	yz	$y\bar{z}$	$\bar{y}\bar{z}$	$\bar{y}z$
x	+	+		+
\bar{x}	+		+	+

$$E = (\textcolor{red}{x}\textcolor{red}{y}) \vee (\textcolor{brown}{\bar{x}}\textcolor{brown}{\bar{y}}) \vee \textcolor{blue}{z}$$

Canonical form would consist of writing all cells separately (6 clauses).

Karnaugh Maps

For optimisation, the idea is to cover the + squares with the minimum number of rectangles. One *cannot* cover any empty cells.

- The rectangles can go 'around the corner'/the actual map should be seen as a *torus*.
- Rectangles must have sides of 1, 2 or 4 squares (three adjacent cells are useless).

Example

	yz	$y\bar{z}$	$\bar{y}\bar{z}$	$\bar{y}z$
x	+	+		+
\bar{x}	+		+	+

$$E = (\textcolor{red}{x}\textcolor{red}{y}) \vee (\textcolor{brown}{\bar{x}}\textcolor{brown}{\bar{y}}) \vee \textcolor{blue}{z}$$

Canonical form would consist of writing all cells separately (6 clauses).

Exercise

Exercise

RW: 10.6.6(c)

	yz	$y\bar{z}$	$\bar{y}\bar{z}$	$\bar{y}z$
wx	+	+		+
$w\bar{x}$	+	+	+	+
$\bar{w}\bar{x}$			+	+
$\bar{w}x$	+			+

Outline

What is logic?

Boolean Logic

Boolean Functions

Conjunctive and Disjunctive Normal Form

Karnaugh Maps

Boolean Algebras

Definition: Boolean Algebra

Definition

A **Boolean algebra** is a structure $(T, \vee, \wedge, ', 0, 1)$ where

- $0, 1 \in T$
- $\vee, \wedge : T \times T \rightarrow T$ (called **join** and **meet** respectively)
- $' : T \rightarrow T$ (called **complementation**)

and the following laws hold for all $x, y, z \in T$:

Commutativity: $x \vee y = y \vee x, \quad x \wedge y = y \wedge x$

Associativity: $(x \vee y) \vee z = x \vee (y \vee z)$
 $(x \wedge y) \wedge z = x \wedge (y \wedge z)$

Distributivity: $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$
 $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$

Identity: $x \vee 0 = x, \quad x \wedge 1 = x$

Complementation: $x \vee x' = 1, \quad x \wedge x' = 0$

Examples of Boolean Algebras

Example

The set of subsets of a (singleton) set $X = \{x\}$:

$$\mathcal{T} : \text{Pow}(X) = \{\{x\}, \emptyset\}$$

$$\vee \text{ (join)} : \cup$$

$$\wedge \text{ (meet)} : \cap$$

$$' \text{ (complement)} : \cdot^c$$

$$0 : \emptyset$$

$$1 : X \quad (\mathcal{U})$$

The Laws of Boolean algebra follow from the Laws of Set Operations.

Examples of Boolean Algebras

Example

The two element Boolean Algebra :

$$\mathbb{B} = (\{\text{true}, \text{false}\}, ||, \&\&, !, \text{false}, \text{true})$$

where $!$, $\&\&$, $||$ are defined as:

- $!\text{true} = \text{false}; !\text{false} = \text{true},$
- $\text{true} \&\& \text{true} = \text{true}; \dots$
- $\text{true} || \text{true} = \text{true}; \dots$

Examples of Boolean Algebras

Example

Cartesian products of \mathbb{B} , that is n -tuples of 0's and 1's with Boolean operations, e.g. \mathbb{B}^4 :

$$\text{join: } (1, 0, 0, 1) \vee (1, 1, 0, 0) = (1, 1, 0, 1)$$

$$\text{meet: } (1, 0, 0, 1) \wedge (1, 1, 0, 0) = (1, 0, 0, 0)$$

$$\text{complement: } (1, 0, 0, 1)' = (0, 1, 1, 0)$$

$$0 : (0, 0, 0, 0)$$

$$1 : (1, 1, 1, 1).$$

Examples of Boolean Algebras

Example

Functions from any set S to \mathbb{B} ; that is, \mathbb{B}^S

If $f, g : S \rightarrow \mathbb{B}$ then

$(f \vee g) : S \rightarrow \mathbb{B}$ defined by $s \mapsto f(s) \parallel g(s)$

$(f \wedge g) : S \rightarrow \mathbb{B}$ defined by $s \mapsto f(s) \&\& g(s)$

$f' : S \rightarrow \mathbb{B}$ defined by $s \mapsto !f(s)$

$0 : S \rightarrow \mathbb{B}$ is the function $s \mapsto 0$

$1 : S \rightarrow \mathbb{B}$ is the function $s \mapsto 1$

Proofs in Boolean Algebras

If you can show that an identity holds using the laws of Boolean Algebra, then that identity holds **in all Boolean Algebras**.

Example

Claim. In all Boolean Algebras

$$x \wedge x = x$$

for all $x \in T$.

Proof:

x	$= x \wedge 1$	[Identity]
	$= x \wedge (x \vee x')$	[Complement]
	$= (x \wedge x) \vee (x \wedge x')$	[Distributivity]
	$= (x \wedge x) \vee 0$	[Complement]
	$= (x \wedge x)$	[Identity]

Duality

Definition

If E is an expression defined using variables (x, y, z , etc), constants (0 and 1), and the operations of Boolean Algebra (\wedge, \vee , and $'$) then $\text{dual}(E)$ is the expression obtained by replacing \wedge with \vee (and vice-versa) and 0 with 1 (and vice-versa).

Definition

If $(T, \vee, \wedge, ', 0, 1)$ is a Boolean Algebra, then $(T, \wedge, \vee, ', 1, 0)$ is also a Boolean algebra, known as the **dual** Boolean algebra.

Theorem (Principle of duality)

If you can show $E_1 = E_2$ using the laws of Boolean Algebra, then $\text{dual}(E_1) = \text{dual}(E_2)$.

Duality

Example

We have shown $x \wedge x = x$.

By duality: $x \vee x = x$.