

The broken stick problem as the continuum limit of a Bernoulli process

Ulysses Zhan¹

¹Department of Physics, University of California, Santa Barbara

May 13, 2026

Run

A **run** is consecutive identical elements in a sequence.

Sequence of Bernoulli trials:



Run of “●”:



The length of the longest run is 5.

The Research Question

Behavior of the longest run when sequence length $n \rightarrow \infty$ while the probability of getting a perfect sequence is constant:

$$n \rightarrow \infty,$$

$$p^n = P = \text{const.}$$

The Discrete Model

Consider a Bernoulli sequence of length n , where each element is either success (S) or failure (F), with probabilities p and $1 - p$ respectively, and independent of each other. Let $P_{n,k}(p)$ be the probability that the length of the longest run of S's is k . What is the expression of $P_{n,k}(p)$?

The Recurrence Relation


$$\underbrace{\bullet \bullet \dots \bullet}_k \underbrace{\circ \circ \dots \circ}_k \quad p^k (1-p) \sum_{s=0}^k P_{n-k-1,s}$$

$$\underbrace{\bullet \bullet \dots \bullet}_s \underbrace{\circ \circ \dots \circ}_k \quad \sum_{s=0}^{k-1} p^s (1-p) P_{n-s-1,k}$$

$$P_{n,0 < k < n} = (1-p) \left(p^k \sum_{s=0}^k P_{n-k-1,s} + \sum_{s=0}^{k-1} p^s P_{n-s-1,k} \right).$$

$$P_{n,0} = (1-p)^n, \quad P_{n,n} = p^n, \quad P_{n,k > n} = 0.$$

Visualizing the Discrete Distribution



Defining the Continuum Limit

Define $P := p^n$ and keep it constant as $n \rightarrow \infty$. The probability distribution of $x := k/n$ is

$$f(P, x) := \lim_{n \rightarrow \infty} n P_{n, xn} \left(P^{1/n} \right) \quad (0 \leq P \leq 1, \quad x \geq 0).$$

The Integral Equation

$$g(P, x) := \frac{1}{P} f(P, x) - \delta(x - 1).$$

$$g(P, 0 \leq x \leq 1) = -\ln P \left(\int_0^{x/(1-x)} (g(P^{1-x}, u) + \delta(u - 1)) \, du \right. \\ \left. + \int_x^{x/(1-x)} \left(g(P^{x/v}, v) + \delta(v - 1) \right) \frac{dv}{v} \right),$$

$$g(P, x > 1) = 0.$$

$$\int_0^\infty g(P, x) \, dx = \frac{1}{P} - 1.$$

Overview of the ADM

The Adomian decomposition method (ADM):

$$g = \Phi(g) .$$


$$g = g_0 + \lambda g_1 + \lambda^2 g_2 + \lambda^3 g_3 + \cdots ,$$

$$\Phi(g) = \Phi_0 + \lambda \Phi_1(g) + \lambda^2 \Phi_2(g, g) + \lambda^3 \Phi_3(g, g, g) + \cdots .$$

$$g_j = \sum_{l=0}^j \sum_{m_1 + \cdots + m_l = j-l} \Phi_l(g_{m_1}, \dots, g_{m_l}) .$$

Piece-by-Piece Solution

$$g^{(q)}(P, x) := g(P, x) \quad \left(\frac{1}{q+1} < x < \frac{1}{q} \right).$$




Solution

$$F(P, x) := \int_0^x dx' \int_0^{x'} dx'' f(P, x'') , \quad r_s := \ln P^{sx-1}.$$

$$F(P, x) = \frac{P}{\ln P} \sum_{s=1}^{\lfloor 1/x \rfloor} \frac{(-1)^s}{s! s} e^{r_s} r_s^s - \frac{\operatorname{li} P - \gamma - \ln(-\ln P)}{\ln P} + x.$$

Visualization of the Solution



The Broken Stick Problem


Break a unit-length stick at k random points. The distribution of the longest segment is $f_k(x)$.

$$f(e^{-\lambda}, x) = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} f_k(x),$$

$$f(P, x) = P \sum_{k=0}^{\infty} \frac{(-\ln P)^k}{k!} f_k(x).$$


Run Statistics to the Gumbel Distribution

$f(P, x)$ tends to the Gumbel distribution as $P \rightarrow 0$.



Broken Stick Problem to the Gumbel Distribution

$f_k(x)$ tends to the Gumbel distribution as $k \rightarrow \infty$.



Summary of Findings

- Derive a recurrence relation for the longest run distribution and formulate its continuum limit to an integral equation.
- Derive a piecewise analytical solution using the ADM.
- Formally establish the link between run statistics and the broken stick problem.
- Formally derive the Gumbel distribution in limiting cases.