Seaborn Assignment

Bivariate_Categorical scatteredPlot:

- Swarmplot:

  sb.swarmplot(x='gender',y='ssc_p',data=dataset)

- Stripplot and swarmplot are Seaborn categorical scatterplots used to visualize data distribution.
- swarmplot adjusts points to avoid overlap, creating a "beeswarm" effect that shows density, while stripplot often overlaps points.
- Swarmplots are best for smaller datasets, while stripplots scale better for large, dense datasets.
- jitter is used to reduce overlap in stripplot.

- Distribution observation-Boxplot:

  sb.boxplot(x='ssc_b',y='ssc_p',data=dataset,hue='gender')



- Boxplot is a percentile method--initial value,Q1,Q2,Q3,Q4. Q2 is a median value.
- The graph shows x axis="ssc_b" and Y axis="ssc_p", blue color is male and brown color is female

- The ssc_b has two types of data they are others and central and the ssc_p students gender both male and female.
- The graph shows in ssc_b others category, male initial score is 50 and Q4 is 90 but female initial score is 45 and Q4 is 85. So when compared with female, male scored higher marks. But the average mark of female 75 is higher than male average score is 65.
- In ssc_b other category the female has lesser outliers.
- The graph shows in ssc_b central category, male initial score is 43and Q4 is 85 but female initial score is 45 and Q4 is 90. So when compared with male, female scored higher marks. But the average mark of female 68 is higher than male average score is 65.