

# Ridge Regression: Regularizing Linear Models

---

Nipun Batra

IIT Gandhinagar

July 30, 2025

# Outline

# When Linear Regression Goes Wrong

**Problem:** What happens when we have too many features or complex polynomials?

## High-Degree Polynomial

$$f(x) = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots + c_nx^n$$

# When Linear Regression Goes Wrong

**Problem:** What happens when we have too many features or complex polynomials?

## High-Degree Polynomial

$$f(x) = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots + c_nx^n$$

# When Linear Regression Goes Wrong

**Problem:** What happens when we have too many features or complex polynomials?

## High-Degree Polynomial

$$f(x) = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots + c_nx^n$$

### Issues:

- Large coefficients  $|c_i|$  can cause instability

# When Linear Regression Goes Wrong

**Problem:** What happens when we have too many features or complex polynomials?

## High-Degree Polynomial

$$f(x) = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots + c_nx^n$$

### Issues:

- Large coefficients  $|c_i|$  can cause instability
- Overfitting to training data

# When Linear Regression Goes Wrong

**Problem:** What happens when we have too many features or complex polynomials?

## High-Degree Polynomial

$$f(x) = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots + c_nx^n$$

### Issues:

- **Large coefficients**  $|c_i|$  can cause instability
- **Overfitting** to training data
- **Poor generalization** to new data

# When Linear Regression Goes Wrong

**Problem:** What happens when we have too many features or complex polynomials?

## High-Degree Polynomial

$$f(x) = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots + c_nx^n$$

### Issues:

- **Large coefficients**  $|c_i|$  can cause instability
- **Overfitting** to training data
- **Poor generalization** to new data



# When Linear Regression Goes Wrong

**Problem:** What happens when we have too many features or complex polynomials?

## High-Degree Polynomial

$$f(x) = c_0 + c_1x + c_2x^2 + c_3x^3 + \dots + c_nx^n$$

### Issues:

- **Large coefficients**  $|c_i|$  can cause instability
- **Overfitting** to training data
- **Poor generalization** to new data

## The Key Insight

**Solution:** Penalize large coefficients to encourage simpler models!

# Pop Quiz: Overfitting Intuition

## Quick Quiz 1

A linear model with coefficients  $[0.1, 0.2, -0.1]$  vs  $[10.2, -15.6, 23.4]$ . Which is likely to generalize better?

a)  $[10.2, -15.6, 23.4]$  (larger coefficients)

**Answer:** b) Smaller coefficients typically indicate a more stable, generalizable model!

# Pop Quiz: Overfitting Intuition

## Quick Quiz 1

A linear model with coefficients  $[0.1, 0.2, -0.1]$  vs  $[10.2, -15.6, 23.4]$ . Which is likely to generalize better?

- a)  $[10.2, -15.6, 23.4]$  (larger coefficients)
- b)  $[0.1, 0.2, -0.1]$  (smaller coefficients)

**Answer:** b) Smaller coefficients typically indicate a more stable, generalizable model!

# Pop Quiz: Overfitting Intuition

## Quick Quiz 1

A linear model with coefficients  $[0.1, 0.2, -0.1]$  vs  $[10.2, -15.6, 23.4]$ . Which is likely to generalize better?

- a)  $[10.2, -15.6, 23.4]$  (larger coefficients)
- b)  $[0.1, 0.2, -0.1]$  (smaller coefficients)
- c) Both are equivalent

**Answer:** b) Smaller coefficients typically indicate a more stable, generalizable model!

# Two Ways to Think About Ridge Regression

# Two Ways to Think About Ridge Regression

# Two Ways to Think About Ridge Regression

## Constrained Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$   
subject to  $\|\boldsymbol{\theta}\|^2 \leq S$

**Interpretation:** Find best fit while keeping coefficients "small"

# Two Ways to Think About Ridge Regression

## Constrained Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$   
subject to  $\|\boldsymbol{\theta}\|^2 \leq S$

**Interpretation:** Find best fit while keeping coefficients "small"



# Two Ways to Think About Ridge Regression

## Constrained Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$   
subject to  $\|\boldsymbol{\theta}\|^2 \leq S$

**Interpretation:** Find best fit while keeping coefficients "small"

# Two Ways to Think About Ridge Regression

## Constrained Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$   
subject to  $\|\boldsymbol{\theta}\|^2 \leq S$

**Interpretation:** Find best fit while keeping coefficients "small"

## Penalized Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 + \lambda\|\boldsymbol{\theta}\|^2$

**Interpretation:** Balance fit quality vs coefficient size

# Two Ways to Think About Ridge Regression

## Constrained Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$   
subject to  $\|\boldsymbol{\theta}\|^2 \leq S$

**Interpretation:** Find best fit while keeping coefficients "small"

## Penalized Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 + \lambda\|\boldsymbol{\theta}\|^2$

**Interpretation:** Balance fit quality vs coefficient size

# Two Ways to Think About Ridge Regression

## Constrained Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$   
subject to  $\|\boldsymbol{\theta}\|^2 \leq S$

**Interpretation:** Find best fit while keeping coefficients "small"

## Penalized Form

Minimize  $\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 + \lambda\|\boldsymbol{\theta}\|^2$

**Interpretation:** Balance fit quality vs coefficient size

## Key Insight

These formulations are **equivalent!** Different  $\lambda$  values correspond to different constraint budgets  $S$ .

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda\|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda\|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda\|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

### Components:

- **Data Fit:** How well model fits training data

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda\|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

### Components:

- **Data Fit**: How well model fits training data
- **Penalty**: Discourages large coefficients



# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda\|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

### Components:

- **Data Fit**: How well model fits training data
- **Penalty**: Discourages large coefficients
- $\lambda$  (**lambda**): Controls the trade-off

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda\|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

### Components:

- **Data Fit**: How well model fits training data
- **Penalty**: Discourages large coefficients
- $\lambda$  (**lambda**): Controls the trade-off

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda \|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

### Components:

- **Data Fit**: How well model fits training data
- **Penalty**: Discourages large coefficients
- $\lambda$  (**lambda**): Controls the trade-off

### $\lambda$ behavior:

- $\lambda = 0$ : No regularization (ordinary least squares)

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda\|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

### Components:

- **Data Fit**: How well model fits training data
- **Penalty**: Discourages large coefficients
- $\lambda$  (**lambda**): Controls the trade-off

### $\lambda$ behavior:

- $\lambda = 0$ : No regularization (ordinary least squares)
- $\lambda$  small: Slight regularization

# Understanding the Ridge Penalty

## Ridge Regression Objective

$$\mathcal{L}(\boldsymbol{\theta}) = \underbrace{\|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2}_{\text{Data Fit}} + \underbrace{\lambda \|\boldsymbol{\theta}\|^2}_{\text{Complexity Penalty}}$$

### Components:

- **Data Fit**: How well model fits training data
- **Penalty**: Discourages large coefficients
- $\lambda$  (**lambda**): Controls the trade-off

### $\lambda$ behavior:

- $\lambda = 0$ : No regularization (ordinary least squares)
- $\lambda$  small: Slight regularization
- $\lambda$  large: Heavy regularization (coefficients shrink toward 0)

# Pop Quiz: Ridge Parameter

## Quick Quiz 2

What happens to the coefficients as  $\lambda$  increases in Ridge regression?

a) Coefficients become larger to minimize error

**Answer:** b) Ridge penalty  $\lambda \|\theta\|^2$  forces coefficients to shrink!

# Pop Quiz: Ridge Parameter

## Quick Quiz 2

What happens to the coefficients as  $\lambda$  increases in Ridge regression?

- a) Coefficients become larger to minimize error
- b) Coefficients shrink toward zero

**Answer:** b) Ridge penalty  $\lambda \|\theta\|^2$  forces coefficients to shrink!

# Pop Quiz: Ridge Parameter

## Quick Quiz 2

What happens to the coefficients as  $\lambda$  increases in Ridge regression?

- a) Coefficients become larger to minimize error
- b) Coefficients shrink toward zero
- c) Coefficients remain unchanged

**Answer:** b) Ridge penalty  $\lambda \|\theta\|^2$  forces coefficients to shrink!



# Lagrangian Approach

**Starting from the constrained formulation:**

$$\begin{aligned} &\text{Minimize } \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 \\ &\text{subject to } \|\boldsymbol{\theta}\|^2 \leq S \end{aligned}$$

# Lagrangian Approach

**Starting from the constrained formulation:**

$$\begin{aligned} &\text{Minimize } \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 \\ &\text{subject to } \|\boldsymbol{\theta}\|^2 \leq S \end{aligned}$$

**Lagrangian:**

$$L(\boldsymbol{\theta}, \mu) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 + \mu(\|\boldsymbol{\theta}\|^2 - S)$$

where  $\mu \geq 0$  is the Lagrange multiplier.

# Lagrangian Approach

**Starting from the constrained formulation:**

$$\begin{aligned} &\text{Minimize } \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 \\ &\text{subject to } \|\boldsymbol{\theta}\|^2 \leq S \end{aligned}$$

**Lagrangian:**

$$L(\boldsymbol{\theta}, \mu) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2 + \mu(\|\boldsymbol{\theta}\|^2 - S)$$

where  $\mu \geq 0$  is the Lagrange multiplier.

## Key Insight

Setting  $\lambda = \mu$  gives us the penalized form! Different constraint budgets  $S$  correspond to different penalty strengths  $\lambda$ .