if class loaded beamer

Coordinate Descent

Nipun Batra

IIT Gandhinagar

July 29, 2025

random, round-robin

No step-size to choose!

Converges for Lasso objective

Learn $y = \theta_0 + \theta_1 x$ on following dataset, using coordinate descent where initially $(\theta_0, \theta_1) = (2, 3)$ for 2 iterations.

| X | у |
|---|---|
| 1 | 1 |
| 2 | 2 |
| 3 | 3 |

Our predictor,
$$\hat{y} = \theta_0 + \theta_1 x$$

Error for
$$i^{th}$$
 datapoint, $\epsilon_i = y_i - \hat{y}_i$
 $\epsilon_1 = 1 - \theta_0 - \theta_1$
 $\epsilon_2 = 2 - \theta_0 - 2\theta_1$
 $\epsilon_3 = 3 - \theta_0 - 3\theta_1$

$$\mathsf{MSE} = \tfrac{\epsilon_1^2 + \epsilon_2^2 + \epsilon_3^2}{3} = \tfrac{14 + 3\theta_0^2 + 14\theta_1^2 - 12\theta_0 - 28\theta_1 + 12\theta_0\theta_1}{3}$$

CostFunction

Iteration 1

INIT: $\theta_0 = 2$ and $\theta_1 = 3$

 $\theta_1=3$ optimize for θ_0

INIT:
$$\theta_0=2$$
 and $\theta_1=3$

$$\theta_1 = 3$$
 optimize for θ_0

$$\frac{\partial \, \mathsf{MSE}}{\partial \theta_0} = 6\theta_0 + 24 = 0$$

$$\theta_0 = -4$$

CostFunction

INIT:
$$\theta_0 = -4$$
 and $\theta_1 = 3$

$$heta_0 = -4$$
 optimize for $heta_1$

INIT:
$$\theta_0 = -4$$
 and $\theta_1 = 3$

$$\theta_0 = -4$$
 optimize for θ_1

$$\theta_1 = 2.7$$

CostFunction

INIT:
$$\theta_0 = -4$$
 and $\theta_1 = 2.7$

$$\theta_1=2.7$$
 optimize for θ_0

INIT:
$$\theta_0 = -4$$
 and $\theta_1 = 2.7$

$$\theta_1=$$
 2.7 optimize for θ_0

$$\theta_0 = -3.4$$

CostFunction