

[BoldFont=Fira Sans SemiBold]Fira Sans Book Fira Mono

popquizbox[1] colback=nipun-lightblue!10, colframe=nipun-blue,  
boxrule=2pt, arc=3pt, left=8pt, right=8pt, top=8pt,  
bottom=8pt, title= **Quick Quiz 1**, fonttitle=,  
**coltitle=nipun-white, colbacktitle=nipun-blue,**  
**enhanced, attach boxed title to top left=xshift=0pt,**  
**yshift=-2pt, boxed title style=arc=3pt, boxrule=0pt**


definitionbox[1] colback=nipun-green!8, colframe=nipun-green,  
boxrule=1.5pt, arc=2pt, left=6pt, right=6pt, top=6pt,  
bottom=6pt, title= **Definition: 1**, fonttitle=,  
**coltitle=nipun-white, colbacktitle=nipun-green**

examplebox[1] colback=nipun-orange!8, colframe=nipun-orange,  
boxrule=1.5pt, arc=2pt, left=6pt, right=6pt, top=6pt,  
bottom=6pt, title= **Example: 1**, fonttitle=,  
**coltitle=nipun-white, colbacktitle=nipun-orange**

keypointsbox colback=nipun-blue!8, colframe=nipun-blue,  
boxrule=1.5pt, arc=2pt, left=6pt, right=6pt, top=6pt,  
bottom=6pt, title= **Key Points**, fonttitle=,  
**coltitle=nipun-white, colbacktitle=nipun-blue**


# Definition

- Convexity is defined on an interval  $[\alpha, \beta]$




# Definition

- Convexity is defined on an interval  $[\alpha, \beta]$
- The line segment joining  $(a, f(a))$  and  $(b, f(b))$  should be *above or on* the function  $f$  for all points in interval  $[\alpha, \beta]$ .




**Example:**  $y = x^2$

Convex on the entire real line i.e.  $(-\infty, \infty)$




**Example:**  $y = |x|$

Convex on the entire real line i.e.  $(-\infty, \infty)$




**Example:**  $y = e^x$

Convex on the entire real line i.e.  $(-\infty, \infty)$




## Example: $y = \ln x$

Not convex on the entire real line i.e.  $(-\infty, \infty)$



**Example:**  $y = x^3$


It is convex for the interval  $[0, \infty)$






**Example:**  $y = x^3$

It is concave for the interval  $(-\infty, 0]$



**Example:**  $y = x^3$


But, it is not convex for the interval  $(-\infty, \infty)$



# Mathematical Formulation

Function  $f$  is convex on set  $X$ , if  $\forall x_1, x_2 \in X$  and  $\forall t \in [0, 1]$

$$f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2)$$



**Question: Prove that  $f(x) = x^2$  is convex**

**Question: Prove that  $f(x) = x^2$  is convex**

To prove:

$$f(tx_1 + (1 - t)x_2) \leq tf(x_1) + (1 - t)f(x_2)$$

## Question: Prove that $f(x) = x^2$ is convex

To prove:

$$f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2)$$

$$\text{LHS} = f(tx_1 + (1-t)x_2) = t^2x_1^2 + (1-t)^2x_2^2 + 2t(1-t)x_1x_2$$

$$\text{RHS} = tf(x_1) + (1-t)f(x_2) = tx_1^2 + (1-t)x_2^2$$

## Question: Prove that $f(x) = x^2$ is convex

To prove:

$$f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2)$$

$$\text{LHS} = f(tx_1 + (1-t)x_2) = t^2x_1^2 + (1-t)^2x_2^2 + 2t(1-t)x_1x_2$$

$$\text{RHS} = tf(x_1) + (1-t)f(x_2) = tx_1^2 + (1-t)x_2^2$$

Here,

$$\begin{aligned}\text{LHS} - \text{RHS} &= (t^2 - t)x_1^2 + [(1-t)^2 - (1-t)]x_2^2 + 2t(1-t)x_1x_2 \\ &= (t^2 - t)x_1^2 + (t^2 - t)x_2^2 - 2(t^2 - t)x_1x_2 \\ &= (t^2 - t)(x_1 - x_2)^2\end{aligned}$$

## Question: Prove that $f(x) = x^2$ is convex

To prove:

$$f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2)$$

$$\text{LHS} = f(tx_1 + (1-t)x_2) = t^2x_1^2 + (1-t)^2x_2^2 + 2t(1-t)x_1x_2$$

$$\text{RHS} = tf(x_1) + (1-t)f(x_2) = tx_1^2 + (1-t)x_2^2$$

Here,

$$\begin{aligned}\text{LHS} - \text{RHS} &= (t^2 - t)x_1^2 + [(1-t)^2 - (1-t)]x_2^2 + 2t(1-t)x_1x_2 \\ &= (t^2 - t)x_1^2 + (t^2 - t)x_2^2 - 2(t^2 - t)x_1x_2 \\ &= (t^2 - t)(x_1 - x_2)^2\end{aligned}$$

Here,  $(t^2 - t) \leq 0$  since  $t \in [0, 1]$  and  $(x_1 - x_2)^2 \geq 0$

Hence,  $\text{LHS} - \text{RHS} \leq 0$

Hence  $\text{LHS} \leq \text{RHS}$

Hence proved.



## Alternative ways to prove convexity

### The Double-Derivative Test

If  $f''(x) > 0$ , the function is convex.

For example,

$$\frac{\partial^2(x^2)}{\partial x^2} = 2 > 0 \Rightarrow x^2 \text{ is a convex function.}$$

## Alternative ways to prove convexity

The double derivative test for multi-parameter function is equal to using the Hessian Matrix

A function  $f(x_1, x_2, \dots, x_n)$  is convex iff its  $n \times n$  Hessian Matrix is positive semidefinite for all possible values of  $(x_1, x_2, \dots, x_n)$

$$\mathbf{H} = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{bmatrix}$$

## Alternative ways to prove convexity

Show that  $f(x_1, x_2) = x_1^2 + x_2^2$  is convex.

## Alternative ways to prove convexity

Show that  $f(x_1, x_2) = x_1^2 + x_2^2$  is convex.

$$\mathbf{H} = \begin{bmatrix} \frac{\partial^2(x_1^2+x_2^2)}{\partial x_1^2} & \frac{\partial^2(x_1^2+x_2^2)}{\partial x_1 \partial x_2} \\ \frac{\partial^2(x_1^2+x_2^2)}{\partial x_2 \partial x_1} & \frac{\partial^2(x_1^2+x_2^2)}{\partial x_2^2} \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

## Alternative ways to prove convexity

Show that  $f(x_1, x_2) = x_1^2 + x_2^2$  is convex.

$$\mathbf{H} = \begin{bmatrix} \frac{\partial^2(x_1^2+x_2^2)}{\partial x_1^2} & \frac{\partial^2(x_1^2+x_2^2)}{\partial x_1 \partial x_2} \\ \frac{\partial^2(x_1^2+x_2^2)}{\partial x_2 \partial x_1} & \frac{\partial^2(x_1^2+x_2^2)}{\partial x_2^2} \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

Eigenvalues of  $\mathbf{H}$  are 2 and  $2 > 0 \Rightarrow \mathbf{H}$  is positive semidefinite.

Hence,  $f(x_1, x_2) = x_1^2 + x_2^2$  is convex.

## Convexity of linear least squares

Prove the convexity of linear least squares i.e.  $f(\boldsymbol{\theta}) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$

## Convexity of linear least squares

Prove the convexity of linear least squares i.e.  $f(\boldsymbol{\theta}) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$

We will use the double derivative (Hessian)

## Convexity of linear least squares

Prove the convexity of linear least squares i.e.  $f(\boldsymbol{\theta}) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$

We will use the double derivative (Hessian)

$$\frac{df}{d\boldsymbol{\theta}} = \frac{d(\|\mathbf{y}\|^2 - 2\mathbf{y}^T \mathbf{X}\boldsymbol{\theta} + \|\mathbf{X}\boldsymbol{\theta}\|^2)}{d\boldsymbol{\theta}} = -2\mathbf{y}^T \mathbf{X} + 2(\mathbf{X}\boldsymbol{\theta})^T \mathbf{X}$$



## Convexity of linear least squares

Prove the convexity of linear least squares i.e.  $f(\boldsymbol{\theta}) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$

We will use the double derivative (Hessian)

$$\frac{df}{d\boldsymbol{\theta}} = \frac{d(\|\mathbf{y}\|^2 - 2\mathbf{y}^T \mathbf{X}\boldsymbol{\theta} + \|\mathbf{X}\boldsymbol{\theta}\|^2)}{d\boldsymbol{\theta}} = -2\mathbf{y}^T \mathbf{X} + 2(\mathbf{X}\boldsymbol{\theta})^T \mathbf{X}$$

$$\frac{d^2 f}{d\boldsymbol{\theta}^2} = \mathbf{H} = 2\mathbf{X}^T \mathbf{X}$$

## Convexity of linear least squares

Prove the convexity of linear least squares i.e.  $f(\boldsymbol{\theta}) = \|\mathbf{y} - \mathbf{X}\boldsymbol{\theta}\|^2$

We will use the double derivative (Hessian)

$$\frac{df}{d\boldsymbol{\theta}} = \frac{d(\|\mathbf{y}\|^2 - 2\mathbf{y}^T\mathbf{X}\boldsymbol{\theta} + \|\mathbf{X}\boldsymbol{\theta}\|^2)}{d\boldsymbol{\theta}} = -2\mathbf{y}^T\mathbf{X} + 2(\mathbf{X}\boldsymbol{\theta})^T\mathbf{X}$$

$$\frac{d^2f}{d\boldsymbol{\theta}^2} = \mathbf{H} = 2\mathbf{X}^T\mathbf{X}$$

$\mathbf{X}^T\mathbf{X}$  is positive semidefinite for any  $\mathbf{X} \in \mathbb{R}^{m \times n}$ .

Hence, linear least squares function is convex.

# Properties of Convex Functions

- If  $f(x)$  is convex, then  $kf(x)$  is also convex, for some constant  $k$

## Properties of Convex Functions

- If  $f(x)$  is convex, then  $kf(x)$  is also convex, for some constant  $k$
- If  $f(x)$  and  $g(x)$  are convex, then  $f(x) + g(x)$  is also convex.

# Properties of Convex Functions

- If  $f(x)$  is convex, then  $kf(x)$  is also convex, for some constant  $k$
- If  $f(x)$  and  $g(x)$  are convex, then  $f(x) + g(x)$  is also convex.

Using this we can say that:

- $(\mathbf{y} - \mathbf{X}\boldsymbol{\theta})^T(\mathbf{y} - \mathbf{X}\boldsymbol{\theta}) + \boldsymbol{\theta}^T\boldsymbol{\theta}$  is convex
- $(\mathbf{y} - \mathbf{X}\boldsymbol{\theta})^T(\mathbf{y} - \mathbf{X}\boldsymbol{\theta}) + \|\boldsymbol{\theta}\|_1$  is convex