

## สรุป Chapter 8 (ต่อ)

### Bayes' Theorem: Basics

- Total probability Theorem:

$$p(B) = \sum_i p(B|A_i)p(A_i)$$

- Bayes' Theorem:

$$p(H|X) = \frac{p(X|H)P(H)}{p(X)} \propto p(X|H)P(H)$$

posteriori probability      likelihood      prior probability

What we should choose      What we just see      What we knew previously

- X: a data sample ("evidence")

Prediction can be done based on Bayes' Theorem:

- H: X belongs to class C

Classification is to derive the maximum posteriori

- สูตรในการหาค่า

### Naïve Bayes Classifier: Making a Naïve Assumption

- Practical difficulty of Naïve Bayes inference: It requires initial knowledge of many probabilities, which may not be available or involving significant computational cost

- A Naïve Special Case

- Make an additional assumption to simplify the model, but achieve comparable performance.

attributes are conditionally independent  
(i.e., no dependence relation between attributes)

$$p(X|C_i) = \prod_k p(x_k|C_i) = p(x_1|C_i) \cdot p(x_2|C_i) \cdots p(x_n|C_i)$$

- Only need to count the class distribution w.r.t. features

-จะต้องมีความสัมพันธ์กันถึงจะหาค่าได้

## Naïve Bayes Classifier: Categorical vs. Continuous Valued Features

- If feature  $x_k$  is categorical,  $p(x_k = v_k | C_i)$  is the # of tuples in  $C_i$  with  $x_k = v_k$ , divided by  $|C_{i,D}|$  (# of tuples of  $C_i$  in  $D$ )

$$p(X | C_i) = \prod_k p(x_k | C_i) = p(x_1 | C_i) \cdot p(x_2 | C_i) \cdots p(x_n | C_i)$$

- If feature  $x_k$  is continuous-valued,  $p(x_k = v_k | C_i)$  is usually computed based on Gaussian distribution with a mean  $\mu$  and standard deviation  $\sigma$

$$p(x_k = v_k | C_i) = N(x_k | \mu_{C_i}, \sigma_{C_i}) = \frac{1}{\sqrt{2\pi}\sigma_{C_i}} e^{-\frac{(x - \mu_{C_i})^2}{2\sigma^2}}$$

## Naïve Bayes Classifier: Training Dataset

Class:

C1:buys\_computer = 'yes'

C2:buys\_computer = 'no'

Data to be classified:

X = (age <=30, Income = medium,

Student = yes, Credit\_rating = Fair)

age	income	student	credit_rating	buys_computer
<=30	high	no	fair	no
<=30	high	no	excellent	no
31...40	high	no	fair	yes
>40	medium	no	fair	yes
>40	low	yes	fair	yes
>40	low	yes	excellent	no
31...40	low	yes	excellent	yes
<=30	medium	no	fair	no
<=30	low	yes	fair	yes
>40	medium	yes	fair	yes
<=30	medium	yes	excellent	yes
31...40	medium	no	excellent	yes
31...40	high	yes	fair	yes
>40	medium	no	excellent	no

- ตัวอย่างในการศึกษา

- แบ่งออกเป็น X และ Y

## Naïve Bayes Classifier: An Example

- $P(C_i)$ :  $P(\text{buys\_computer} = \text{"yes"}) = 9/14 = 0.643$   
 $P(\text{buys\_computer} = \text{"no"}) = 5/14 = 0.357$
  - Compute  $P(X|C_i)$  for each class
    - $P(\text{age} = \text{"<=30"} | \text{buys\_computer} = \text{"yes"}) = 2/9 = 0.222$
    - $P(\text{age} = \text{"<= 30"} | \text{buys\_computer} = \text{"no"}) = 3/5 = 0.6$
    - $P(\text{income} = \text{"medium"} | \text{buys\_computer} = \text{"yes"}) = 4/9 = 0.444$
    - $P(\text{income} = \text{"medium"} | \text{buys\_computer} = \text{"no"}) = 2/5 = 0.4$
    - $P(\text{student} = \text{"yes"} | \text{buys\_computer} = \text{"yes"}) = 6/9 = 0.667$
    - $P(\text{student} = \text{"yes"} | \text{buys\_computer} = \text{"no"}) = 1/5 = 0.2$
    - $P(\text{credit\_rating} = \text{"fair"} | \text{buys\_computer} = \text{"yes"}) = 6/9 = 0.667$
    - $P(\text{credit\_rating} = \text{"fair"} | \text{buys\_computer} = \text{"no"}) = 2/5 = 0.4$
  - **X = (age <= 30 , income = medium, student = yes, credit\_rating = fair)**
    - $P(X|C_i)$  :  $P(X | \text{buys\_computer} = \text{"yes"}) = 0.222 \times 0.444 \times 0.667 \times 0.667 = 0.044$
    - $P(X|C_i)$  :  $P(X | \text{buys\_computer} = \text{"no"}) = 0.6 \times 0.4 \times 0.2 \times 0.4 = 0.019$
    - $P(X|C_i) * P(C_i)$  :  $P(X | \text{buys\_computer} = \text{"yes"}) * P(\text{buys\_computer} = \text{"yes"}) = 0.028$
    - $P(X|C_i) * P(C_i)$  :  $P(X | \text{buys\_computer} = \text{"no"}) * P(\text{buys\_computer} = \text{"no"}) = 0.007$
- Therefore, X belongs to class ("buys\_computer = yes")

age	income	student	credit_rating	buys_computer
<=30	high	no	fair	no
<=30	high	no	excellent	no
31...40	high	no	fair	yes
>40	medium	no	fair	yes
>40	low	yes	fair	yes
>40	low	yes	excellent	no
31...40	low	yes	excellent	yes
<=30	medium	no	fair	no
<=30	low	yes	fair	yes
>40	medium	yes	fair	yes
<=30	medium	yes	excellent	yes
31...40	medium	no	excellent	yes
31...40	high	yes	fair	yes
>40	medium	no	excellent	no

- ตัวอย่างในการหาค่าต่าง ๆ