

Isomorphisms of Graphs



Definition:

Let G and G' be graphs with vertex sets $V(G)$ and $V(G')$ and edge sets $E(G)$ and $E(G')$, respectively. **G is isomorphic to G'** if, and only if, there exist one-to-one correspondences $g: V(G) \rightarrow V(G')$ and $h: E(G) \rightarrow E(G')$ that preserve the edge-endpoint functions of G and G' in the sense that for each $v \in V(G)$ and $e \in E(G)$,

$$v \text{ is an endpoint of } e \Leftrightarrow g(v) \text{ is an endpoint of } h(e).$$





Example:

Show that the following two graphs are isomorphic.



Example:

Show that the following two graphs are isomorphic.



Example:

Find all non-isomorphic graphs that have two vertices and two edges. In other words, find a collection of representative graphs with two vertices and two edges such that every graph with two vertices and two edges is isomorphic to one in the collection.

Example:


Find all non-isomorphic graphs that have two vertices and two edges. In other words, find a collection of representative graphs with two vertices and two edges such that every graph with two vertices and two edges is isomorphic to one in the collection.



(a)



(b)



(c)



(d)

Definition:

A property P is called an **invariant for graph isomorphism** if, and only if, given any graphs G and G' , if G has property P and G' is isomorphic to G , then G' has property P .

Theorem


Each of the following properties is an invariant for graph isomorphism, where n , m , and k are all nonnegative integers:

1. has n vertices
2. has m edges
3. has a vertex of degree k
4. has m vertices of degree k
5. has a circuit of length k
6. has a simple circuit of length k
7. has m simple circuits of length k
8. is connected
9. has an Euler circuit
10. has a Hamiltonian circuit.


Example:

Show that the following pairs of graphs are not isomorphic by finding an isomorphic invariant that they do not share.

a.




G




G'

b.



H





H'

Definition:




If G and G' are simple graphs, then **G is isomorphic to G'** if, and only if, there exists a one-to-one correspondence g from the vertex set $V(G)$ of G to the vertex set $V(G')$ of G' that preserves the edge-endpoint functions of G and G' in the sense that for all vertices u and v of G ,

$$\{u, v\} \text{ is an edge in } G \Leftrightarrow \{g(u), g(v)\} \text{ is an edge in } G'.$$

Example: Are the two graphs shown below isomorphic? If so, define an isomorphism.



Example: Are the two graphs shown below isomorphic? If so, define an isomorphism.



Edges of G	Edges of G'
$\{a, b\}$	$\{y, w\} = \{g(a), g(b)\}$
$\{a, c\}$	$\{y, x\} = \{g(a), g(c)\}$
$\{a, d\}$	$\{y, z\} = \{g(a), g(d)\}$
$\{c, d\}$	$\{x, z\} = \{g(c), g(d)\}$