Akoizo Original Practice - 2027 Rotation Form 2

Division C | Astronomy | 50-minute written practice

Original Akoizo curriculum practice. This is not a historical tournament paper. Written build, CAD, lab and communication exercises supplement real practical preparation.

Name: _____ Date: _____

1. (2 points) Conventional H-R temperature increases toward…

Unit: Stellar properties, fusion, and the H-R diagram.
Worked-case data: A star has radius 2 Rsun and temperature 2 Tsun. Calculate L/Lsun.

A. The left

B. The right

C. The top only

D. The bottom only

Answer: _____

2. (2 points) Five magnitudes fainter means flux is…

Unit: Astrophysics equations and quantitative synthesis.
Worked-case data: A source has m = 10 and M = 5, with negligible extinction. Find distance and compare its flux with the same source at 10 pc.

A. 100 times larger

B. Unchanged

C. 100 times smaller

D. 5 times smaller

Answer: _____

3. (2 points) A reflection nebula is seen primarily through…

Unit: Galaxies, stellar populations, and the interstellar medium.
Worked-case data: A region is dark in visible light but bright in infrared and contains embedded young sources. Explain without declaring it empty.

A. A mandatory supernova

B. No illumination

C. Scattered light

D. Only nuclear fusion in every dust grain

Answer: _____

4. (2 points) A cluster turnoff is primarily an age clue because…

Unit: Cloud collapse and mass-dependent stellar evolution.
Worked-case data: A toy main-sequence model uses L □ M^3·^5 and lifetime □ M/L. Compare a 2 Msun star with the Sun.

A. Faint stars always form last

B. Every cluster is dust-free

C. Massive stars evolve off the main sequence sooner

D. All stars have identical masses

Answer: _____

5. (4 points) Convert 50 mas to arcseconds.

Unit: Observational astronomy and the electromagnetic spectrum.
Worked-case data: A star has parallax 50 mas and approximate blackbody temperature 5800 K.

Answer: _____
_____
_____

6. (4 points) Calculate the radius ratio in the example.

Unit: Advanced graph and image interpretation.
Worked-case data: Two stars have equal effective temperature but luminosities 1 and 100 Lsun. Compare radii.

Answer: _____
_____
_____

7. (4 points) Why can material around a black hole shine?

Unit: Supernovae, white dwarfs, neutron stars, and black holes.
Worked-case data: Use 3 km per solar mass as a rounded Schwarzschild-radius scale for a 10 Msun nonrotating black hole.

Answer: _____
_____
_____

8. (4 points) Why not apply main-sequence mass-luminosity scaling to a giant?

Unit: Stellar properties, fusion, and the H-R diagram.
Worked-case data: A star has radius 2 Rsun and temperature 2 Tsun. Calculate L/Lsun.

Answer: _____
_____
_____

9. (4 points) What assumption does F = L/(4πd^2) use?

Unit: Astrophysics equations and quantitative synthesis.
Worked-case data: A source has m = 10 and M = 5, with negligible extinction. Find distance and compare its flux with the same source at 10 pc.

Answer: _____
_____
_____

10. (4 points) What should be matched across wavelength images?

Unit: Galaxies, stellar populations, and the interstellar medium.
Worked-case data: A region is dark in visible light but bright in infrared and contains embedded young sources. Explain without declaring it empty.

Answer: _____
_____
_____

11. (2 points) A synthetic star has parallax 0.005 arcseconds. Use d(pc)=1/p to find distance.

Answer: _____
_____
_____

12. (2 points) A standard-candle exercise gives apparent magnitude 12, absolute magnitude 2 and negligible extinction. Use m-M=5log10(d/10 pc). Find d.

Answer: _____
_____
_____

13. (2 points) A star has twice the reference radius and the same temperature. Find its luminosity ratio using L proportional to R^2 T^4.

Answer: _____
_____
_____

14. (2 points) Use Wien's constant 2.90e-3 m K for a 5,800 K star. Find peak wavelength.

Answer: _____
_____
_____

15. (2 points) An artificial galaxy recession speed is 140 km/s. Use H0=70 km/s/Mpc and the nearby linear Hubble law to find distance.

Answer: _____
_____
_____

16. (2 points) A normalized binary exercise has separation 2 AU and period 2 years. Use total mass in solar masses=a^3/P^2. Find total mass.

Answer: _____
_____
_____

17. (2 points) NASA identifies M82 X-2 as an ultraluminous pulsar. What does a coherent rotation-linked pulse train imply about the compact object, and why is luminosity alone insufficient?

Answer: _____
_____
_____

18. (2 points) SN 2014J is a Type Ia supernova in M82. Explain why standardized Type Ia luminosities can constrain distance and why dust must be considered.

Answer: _____
_____
_____

19. (2 points) GW170817 was observed in gravitational waves and electromagnetic radiation. Explain the value of this multi-messenger evidence for a neutron-star merger.

Answer: _____
_____
_____

20. (2 points) Synthetic starburst-galaxy case 2: ultraviolet emission and infrared dust emission both increase. Explain why using only visible light can miss part of the recent star formation.

Answer: _____
_____
_____