Scioly Summer Study Session - Cherrie_Lan - Repaired practice edition - Answer key

Division C | Anatomy And Physiology | Repair appendix

This appendix contains authored replacements and supplied written alternatives. It does not reconstruct unpublished historical material. For the listed item IDs, use this appendix instead of the archived wording and key.

ANSWER KEY - Review only after submitting.

MC-12. (4 points) Authored replacement for source item Question MC-12: Left foot infection drains via which duct to which vein?

Unit: Central Immune Organs and Lymph Flow.
Worked-case data: A small skin abrasion introduces material into tissue fluid of the left foot. Trace a plausible route toward immune surveillance and return to blood.
Separate investigation observations:
Origin: Antigen enters interstitial fluid around a skin abrasion.
Transport: Afferent lymph carries antigen-bearing cells from the tissue.
Cell encounter: Mature naive lymphocytes meet presented antigen in organized lymphoid tissue.

answer: Thoracic duct to left subclavian vein. (4 points).

SA-4. (2 points) Authored replacement for source item Question SA-4: Aspirated peanut most likely lodges in which bronchus and why?

Unit: Airways: From Nose to Bronchioles.
Worked-case data: Tube A has the baseline radius. Tube B has 80% of that radius. Both have the same length, viscosity, and driving pressure.

A. Right, wider and more vertical

B. Left, more horizontal

C. Trachea only

D. Left, narrower

Correct: A