



Abstract #212418

Comparing Loop Diuretic Efficacy and Safety in Patients with Acute Myeloid Leukemia and Differentiation Syndrome

Sanjana Nethagani^{1*}, Paul Fudacz^{2*}, Mark Furner^{2*} and Umer Rizwan, MD³

¹West Virginia University, Morgantown, WV; ²West Virginia School of Osteopathic Medicine, Parkersburg, WV; ³Camden Clark Medical Center/WVU Hospitals/Internal Medicine, Hospital, Parkersburg, WV

Background: Differentiation syndrome (DS) is a group of life-threatening complications associated with acute myeloid leukemia (AML) treatment. Loop diuretics are used to manage fluid overload and electrolyte imbalances in DS. The optimal choice of loop diuretics is unclear. Previous studies showed bumetanide and furosemide have similar efficacy and safety in patients with congestive heart failure. Recent research suggest torsemide may be non-inferior to furosemide in critically ill patients with acute kidney injury.

Methods: We performed a retrospective analysis of 5,670 patients diagnosed with AML and differentiation syndrome from a large federated network database. Patients were treated with furosemide (n=2,431), bumetanide (n=466), or torsemide (n=206), alongside glucocorticoids and allopurinol. Propensity score matching was utilized to control for confounding variables, including age, sex, and comorbidities such as chronic kidney disease, diabetes mellitus (types 1 and 2) and hypertension. Patients were matched by AML subtype, including acute myeloid leukemia and acute promyelocytic leukemia.

Results: No significant differences were seen in overall mortality or the incidence of acute renal failure among the three cohorts. Mean glomerular filtration rate, serum creatinine, blood urea nitrogen, serum uric acid, and serum potassium, did not differ significantly between the loop diuretic groups. However, a lower risk of hemodialysis utilization was observed in patients receiving furosemide compared to bumetanide (risk ratio 0.584, 95% CI 0.414-0.825).

Conclusion: Our findings suggest that furosemide, bumetanide, and torsemide have comparable efficacy and safety profiles in patients with AML and differentiation syndrome. Notably, furosemide was associated with a lower rate of hemodialysis

utilization compared to bumetanide, with no significant difference in renal function or mortality. This may be due to bumetanide's higher potency, rapid absorption, and shorter half-life, potentially increasing the risk of electrolyte imbalances and acute kidney injury. Conversely, furosemide may facilitate greater cumulative sodium excretion in patients with chronic renal insufficiency.

Abstract ID#:

212418

Password:

273230

Title:

Comparing Loop Diuretic Efficacy and Safety in Patients with Acute Myeloid Leukemia and Differentiation Syndrome

Submitter's E-mail Address:

ur0001@hsc.wvu.edu

Category:

Clinical/Epidemiological Research

Keywords:

Research, Clinical Practice (Health Services and Quality), Clinical Research, Health outcomes research, Adverse Events

First Time Submitting:

No

Scheduling conflicts due to religious observation:

I do not have any scheduling conflicts due to religious observation.

Is this abstract a Trial in Progress:

No

Registered Clinical Trial:

No

OffLabel Disclosure:

No

Compliance with the Declaration of Helsinki for Studies Involving Human Subjects:

N/A

Interim Analysis of Clinical Trial:

No

Update Analyses:

No

Research Funding:

Does not apply

ASH Funding:**Is the first author/presenter of this abstract a hematologist in training?:**

Yes

Review Category Selection:

615A. Acute Myeloid Leukemias: Clinical and Epidemiological

First Presenter

Presenter**Corresponding Presenter**

Sanjana Nethagani

West Virginia University

Morgantown, WV

Email: nethaganisanjana@gmail.com -- Will not be published**In the past 24 months, have you had any financial relationships with an ineligible company as defined above?** NoSigned on 08/02/2024 by *Sanjana Nethagani*Second Author

Paul Fudacz

West Virginia School of Osteopathic Medicine

800 Garfield Ave

Parkersburg, WV

Email: pfudacz@osteo.wvsom.edu -- Will not be published**In the past 24 months, have you had any financial relationships with an ineligible company as defined above?** NoSigned on 08/02/2024 by *Paul Fudacz*Third Author

Mark Furner

West Virginia School of Osteopathic Medicine

800 Garfield Ave

Parkersburg, WV

Email: markfurnerii@gmail.com -- Will not be published**In the past 24 months, have you had any financial relationships with an ineligible company as defined above?** NoSigned on 08/02/2024 by *Mark Furner*Fourth Author

Umer Rizwan, MD

Hospital

120 Ninth St

Apartment B

Camden Clark Medical Center/WVU Hospitals/Internal Medicine

Parkersburg, WV 26101

Email: ur0001@hsc.wvu.edu -- Will not be published**In the past 24 months, have you had any financial relationships with an ineligible company as defined above?** NoSigned on 08/02/2024 by *Umer Rizwan, MD*

