AI1103-Assignment 1

Name: Umesh Kalvakuntla Roll No.: CS20BTECH11024

Download all python codes from

https://github.com/Umesh-k26/Assignment/blob/ main/Assignment1/assignment.pdf

and latex-tikz codes from

https://github.com/Umesh-k26/Assignment/blob/ main/Assignment1/assignment.tex

QUESTION

A family has two children. What is the probability :. The probability that both the children are boys that both the children are boys given that at least given that at least one of them is a boy is $\frac{1}{3}$. one of them is a boy?

SOLUTION

Given, a family has two children. Let's denote girl by 'G' and boy by 'B'. Sample space of the outcomes is given by:

$$S = [(B, B), (G, B), (B, G), (G, G)]$$

Let X denote the random varible representing number of boys.

$$X = \{0, 1, 2\}$$

$$P(X = 0) = \frac{1}{4}$$

$$P(X = 1) = \frac{2}{4}$$

$$P(X = 2) = \frac{1}{4}$$

$$P(X \ge 1) = \frac{3}{4}$$

To find $P(X = 2 | X \ge 1)$.

$$P(X = 2 | X \ge 1) = \frac{P(X \ge 1) \cap P(X = 2)}{P(X \ge 1)}$$
$$= \frac{P(X = 2)}{P(X \ge 1)}$$
$$= \frac{\frac{1}{4}}{\frac{3}{4}}$$
$$= \frac{1}{3}$$