
Spacetime Unified Substrate Configuration Theory: A Cosmological Unification Model Based on Continuous Medium Phase Transitions

Abstract

This survey examines the Spacetime Unified Substrate Configuration Theory (SPT), a speculative cosmological framework wherein a unified elastic substrate undergoes sequential phase transitions to generate cosmic components. Motivated by persistent tensions between early and late universe measurements and the unresolved nature of dark sectors, the theory posits that spacetime functions as a dynamical continuum whose geometric and thermodynamic properties derive from relativistic elasticity and scalar-tensor modifications. Within this paradigm, dark matter potentially emerges as a condensed superfluid phase characterized by non-zero vacuum expectation values, while dark energy may correspond to a diffuse gas phase with negative pressure driving late-time acceleration. The hypothesis maintains that conventional matter interacts weakly with the substrate at astrophysical scales, preserving approximate geodesic motion and asymptotic consistency with general relativity in weak-field limits. The synthesis covers theoretical foundations including material manifold descriptions, critical phenomena and symmetry breaking, cosmological evolution from primordial perturbations to accelerated expansion, and observational constraints pertaining to structure formation and gravitational wave propagation. While offering a conceptually coherent alternative to standard dark sector models, the framework acknowledges significant limitations regarding fine-tuning requirements, conformal frame dependencies, and the need for robust quantum mechanical foundations.

Keywords: Spacetime Unified Substrate Configuration Theory; relativistic continuum mechanics; phase transitions; dark matter; dark energy

Contents

1	Introduction	4
2	Theoretical Foundations and Conceptual Framework	5
2.1	Relativistic Elasticity and Geometric Continua	6
2.2	Scalar-Tensor Gravity and Cosmological Phase Transitions	7
2.3	Emergent Gravity and Analogue Systems	7
3	The Unified Substrate: Constitutive Properties and Configuration	8
3.1	Geometric Description and Material Manifolds	9
3.2	Elastic Deformation and Stress-Energy Tensors	9
3.3	Thermodynamic State Variables and Equations of State	10
4	Phase Transition Dynamics in the Substrate	10
4.1	Critical Phenomena and Symmetry Breaking	11
4.2	Condensation Mechanisms and Superfluidity	11
4.3	Diffusive Transitions to Gas Phases	12
4.4	Sequential Cosmological Phase Evolution	13
5	Emergence of Cosmic Components from Substrate Phases	13
5.1	Dark Matter as a Condensed Superfluid Phase	14
5.2	Dark Energy as a Diffuse Gas Phase	14
5.3	Weak Coupling Hypothesis and Baryonic Interactions	15
6	Cosmological Evolution and Expansion Dynamics	15
6.1	Early Universe Phase Structure and Primordial Perturbations	16
6.2	Transition to Accelerated Expansion	16
6.3	Asymptotic Consistency with General Relativity	17
7	Observational Constraints and Phenomenological Tests	18

7.1	Implications for the Hubble Tension	18
7.2	Structure Formation and Power Spectrum	19
7.3	Gravitational Wave Phenomenology	19
7.4	Analogue Gravity Experimental Parallels	20
8	Theoretical Consistency and Limitations	20
8.1	Preservation of Lorentz Covariance	21
8.2	Conformal Frame Dependence and Observable Equivalence	21
8.3	Parameter Space and Fine-Tuning Challenges	22
9	Discussion	22
9.1	Comparison with Lambda Cold Dark Matter	22
9.2	Distinctions from Modified Newtonian Dynamics and Superfluid Models	23
9.3	Future Theoretical Directions	23
10	Conclusion	24

1. Introduction

Contemporary cosmological observations reveal persistent tensions between early-universe predictions and late-universe measurements, most notably exemplified by the discrepancy between the Hubble constant H_0 inferred from cosmic microwave background data and that measured through local distance ladders [43, 45, 79]. These discrepancies, approaching statistical significance beyond 5σ in some analyses, challenge the internal consistency of the standard Λ CDM paradigm and suggest that the prevailing model of dark matter as weakly interacting particles and dark energy as a cosmological constant may represent phenomenological approximations rather than fundamental constituents of nature [9, 35]. Concurrently, the physical nature of the dark sector remains empirically unconstrained, with alternative frameworks ranging from scalar-tensor modifications of gravity to emergent spacetime scenarios attracting renewed theoretical interest as potential resolutions to these cosmic inconsistencies.

The conceptual treatment of spacetime as a dynamical medium with material properties, rather than as a passive geometric arena, provides a foundational paradigm shift that motivates the present survey. Drawing upon the mathematical machinery of relativistic continuum mechanics, wherein spacetime itself exhibits elastic deformation characteristics [41, 53], and inspired by emergent gravity frameworks that derive gravitational dynamics from underlying hydrodynamic or quantum fluid behaviors [11, 14], such approaches challenge the ontological distinction between geometry and matter. This material conceptualization of spacetime suggests that cosmic dynamics might originate from the constitutive properties, strain configurations, and thermodynamic phase transitions of a unified substrate, thereby offering a geometrically coherent alternative to the *ad hoc* introduction of dark sectors.

This survey examines the Spacetime Unified Substrate Configuration Theory (SPT), a speculative cosmological framework positing that a single relativistic elastic substrate undergoes sequential phase transitions to generate the observed cosmic components. Within this paradigm, dark matter emerges not as particulate matter but as a condensed superfluid phase of the substrate [9], while dark energy corresponds to a diffuse gas phase characterized by negative pressure contributions, potentially obviating the need for a cosmological constant [35]. The theory hypothesizes that conventional baryonic matter couples weakly to this substrate at astrophysical scales, preserving approximate Lorentz covariance and ensuring asymptotic consistency with general relativistic predictions in appropriate limits. By grounding cosmic evolution in the phase transition dynamics reminiscent of early-universe symmetry breaking [38, 51], SPT offers a potentially unified explanation for both the Hubble tension and the dark sector phenomenology through the elastic deformation and critical phenomena of a fundamental medium.

The subsequent sections develop this framework systematically, beginning with the theoretical foundations in relativistic elasticity and scalar-tensor gravity, followed by detailed examination of the substrate’s constitutive properties and phase transition mechanisms. The analysis proceeds to delineate how distinct substrate phases may correspond to observed cosmic constituents, evaluates the model’s cosmological evolution against observational constraints including the Hubble constant discrepancy, and assesses the theoretical consistency of the weak coupling hypothesis. Throughout, the survey maintains critical attention to the limitations of current evidence and the fine-tuning challenges inherent in substrate-based cosmologies, positioning SPT within the broader landscape of modified gravity and emergent spacetime theories.

2. Theoretical Foundations and Conceptual Framework

The proposition that cosmic dynamics might originate from the elastic deformation and phase transitions of a unified substrate necessitates a confluence of mathematical frameworks drawn from relativistic continuum mechanics, modified gravity theories, and emergent gravity paradigms. Unlike conventional approaches that treat spacetime as a passive geometric arena or invoke ad hoc dark sectors, substrate-based cosmology requires formalisms capable of describing continuous media with constitutive properties that evolve through critical phenomena. This section establishes the theoretical prerequisites for such a framework, examining how relativistic elasticity provides the geometric language for deformable substrates, how scalar-tensor theories furnish mechanisms for cosmological phase transitions, and how analogue gravity systems offer phenomenological guidance for emergent gravitational dynamics.

The integration of these frameworks departs from standard geometric approaches by treating gravitational phenomena as emergent properties of a continuous medium with elastic and thermodynamic degrees of freedom. This reconceptualization requires abandoning the rigid distinction between matter and geometry in favor of a unified description where spacetime itself possesses material properties subject to deformation and phase evolution. Such a theoretical stance draws upon historical attempts to model spacetime as a stressed elastic medium, extending these concepts into fully relativistic regimes where causality and covariance constraints become paramount. The following subsections delineate the specific mathematical structures—relativistic elasticity, scalar-tensor phase dynamics, and fluid-gravity analogies—that enable the rigorous formulation of substrate-based cosmology.

Furthermore, the viability of any substrate model depends upon its ability to reproduce the successes of general relativity in the weak-field limit while simultaneously accommodating novel phenomenology associated with dark matter and dark energy. This dual requirement necessitates careful attention to the conformal frame dependencies inherent in scalar-tensor extensions,

the hyperbolicity conditions governing the evolution of elastic media, and the observational constraints imposed by gravitational wave propagation and structure formation. The theoretical foundations reviewed herein provide the necessary scaffolding for subsequent developments, wherein the substrate’s constitutive properties and phase transition history are specified in detail.

2.1. Relativistic Elasticity and Geometric Continua

The geometric description of continuous media within relativistic contexts extends beyond the traditional Riemannian framework of general relativity, requiring distinct material manifolds and configuration mappings that track elastic deformations across spacetime. Souriau’s general covariant formulation of relativistic hyperelasticity establishes a rigorous foundation for such descriptions by introducing a material manifold distinct from the spacetime metric, wherein the configuration mapping between reference and current states is characterized through relativistic deformation gradients [52, 53]. This approach naturally accommodates strain tensors and elasticity difference tensors that quantify the deviation between the geometric properties of the material and those of the gravitational field, providing a covariant language for describing stressed media in curved spacetime [81]. The elasticity difference tensor, in particular, offers a measure of how the material’s internal geometry diverges from the ambient spacetime geometry, a concept that becomes essential when modeling a substrate whose mechanical properties may vary across cosmological scales [15]. Formulated as a Lagrangian field theory on arbitrary spacetimes, relativistic elasticity casts the equations of motion into a first-order symmetric hyperbolic system, ensuring local existence and uniqueness theorems that guarantee mathematical consistency for dynamical evolution [8].

The extension of elasticity theory to cosmological contexts further requires accommodating thermomechanical complexities and higher-order gradient effects that may become relevant during phase transitions. Relativistic anelasticity frameworks generalize these concepts to include viscoelastic and plastic behaviors through the introduction of inhomogeneity measures and relativistic Eshelby tensors, thereby enabling the modeling of dissipative processes within the substrate [33]. Recent developments in second-gradient theory reveal additional diffeomorphism invariants that reconcile relativistic elasticity with higher-order continuum mechanics, though some invariants exhibit non-objective behavior in the Galilean limit, highlighting subtleties in the transition between relativistic and classical regimes [24]. These geometric formalisms collectively provide the necessary infrastructure to describe a substrate capable of sustaining elastic stresses while undergoing configurational changes, establishing the mathematical preconditions for subsequent cosmological applications.

2.2. Scalar-Tensor Gravity and Cosmological Phase Transitions

Scalar-tensor theories of gravity furnish the natural theoretical arena for modeling cosmological phase transitions within such a substrate, wherein scalar fields serve as order parameters governing the spontaneous symmetry breaking that may reconfigure the medium’s properties. In these frameworks, a scalar field couples to the Ricci scalar or matter fields, dynamically determining the effective gravitational coupling and potentially driving transitions between distinct phases of the cosmic medium [35, 36]. The phenomenon of spontaneous scalarization, wherein compact objects or cosmological regions undergo sudden transitions to configurations with non-trivial scalar profiles, operates analogously to phase transitions in condensed matter systems, with the scalar field vacuum expectation value serving as an order parameter [83]. Recent analyses demonstrate that such scalarization transitions are generically first-order rather than second-order, implying the existence of metastable states and hysteresis effects that could characterize the substrate’s evolution through cosmic history [85]. These transitions may be triggered by density thresholds or environmental conditions, offering a mechanism for the substrate to evolve from one thermodynamic phase to another as the universe expands and cools [25, 84].

The application of scalar-tensor frameworks to cosmology, however, encounters significant interpretational challenges regarding conformal frame dependence, wherein physical observables may vary depending on whether one adopts the Jordan frame (where matter fields couple minimally to the metric) or the Einstein frame (where gravitational equations resemble general relativity but matter fields experience scalar-dependent couplings). Observational analyses demonstrate that cosmological parameters interpreted within specific frames yield quantitatively distinct results, necessitating consistent frame usage when comparing theoretical predictions with data [46, 47]. The frame-independent identification of observables remains an active area of investigation, with evidence suggesting that certain parameters exhibit frame-dependent behavior while others maintain invariance under conformal transformations [46]. These considerations impose essential constraints on substrate models, requiring that purported phase transitions and their observational signatures be formulated in a manner that respects the ambiguities inherent in scalar-tensor representations.

2.3. Emergent Gravity and Analogue Systems

The emergent gravity paradigm and associated analogue systems provide crucial phenomenological guidance for understanding how gravitational dynamics might arise from underlying substrate hydrodynamics, offering laboratory-accessible proxies for testing theoretical hypotheses. The fluid-gravity correspondence establishes that Einstein’s field equations can emerge from the hydrodynamic description of certain quantum fluids, with the energy transport equa-

tion for horizon-fluid fluctuations paralleling the Raychaudhuri equation that governs spacetime singularities [11]. This correspondence suggests that the substrate’s elastic and thermodynamic properties might generate effective gravitational fields through collective excitations, rather than through fundamental geometric degrees of freedom. Hydrodynamic analogues, particularly those utilizing Bose-Einstein condensates, demonstrate that acoustic metrics within these systems can mimic black hole horizons, gravitational wave propagation, and even superradiance phenomena, providing concrete experimental platforms for investigating emergent spacetime geometry [14, 28].

Superfluid dark matter models exemplify the application of these analogue principles to cosmological contexts, wherein dark matter is hypothesized to exist as a Bose-Einstein condensate exhibiting phonon-mediated interactions that reproduce modified Newtonian dynamics at galactic scales while preserving standard cosmological evolution at larger scales [9]. Such frameworks demonstrate that gravitational dynamics can emerge from the hydrodynamic behavior of a quantum fluid, with the effective metric experienced by perturbations determined by the fluid’s density and flow velocity. Recent investigations into gravitational wave propagation through acoustic analogues reveal that perturbations in the fluid can excite horizon-like responses, suggesting that the substrate might similarly affect gravitational wave phenomenology through its phase-dependent refractive properties [28, 60]. These analogue systems, while not directly verifying the substrate hypothesis, constrain the possible equations of state and transport coefficients that a cosmological medium must possess to generate emergent gravity consistent with observations.

3. The Unified Substrate: Constitutive Properties and Configuration

The Spacetime Unified Substrate Configuration Theory posits the existence of a fundamental continuous medium whose constitutive properties and geometric configuration underpin the phenomenology of cosmic evolution. Unlike conventional field-theoretic approaches that treat spacetime as a passive geometric arena or dark sectors as particle constituents, this framework conceptualizes the substrate as a relativistic elastic medium possessing intrinsic material properties that distinguish it from the metric manifold itself. The mathematical description of such a medium requires a general covariant formulation of continuum mechanics that extends classical elasticity theory into the relativistic domain, wherein the substrate’s state is characterized by deformation measures, stress-energy content, and thermodynamic variables that may evolve without presupposing a specific primordial phase configuration [8, 41, 53].

Central to this characterization is the distinction between the material manifold, representing the reference configuration of the substrate, and the spacetime manifold, representing its

current dynamical state. This distinction necessitates a configuration mapping that respects general covariance while permitting the description of elastic strains and material flows. The geometric framework must accommodate both the covariant derivative structures of general relativity and the pull-back operations characteristic of continuum mechanics, ensuring that elastic deformations transform appropriately under coordinate changes while maintaining physical interpretability across cosmological scales [7, 52].

3.1. Geometric Description and Material Manifolds

The geometric foundation of the unified substrate rests upon the construction of a material manifold distinct from the spacetime metric, wherein the configuration mapping between reference and current states is characterized through relativistic deformation gradients. Following Souriau's general covariant formulation of relativistic hyperelasticity, the substrate is described by a deformation map ϕ from the material manifold $\mathcal{M}$ to the spacetime manifold $\mathcal{N}$, with the relativistic deformation gradient F defined as the differential of this mapping [52, 53]. This framework ensures that strain measures, such as the Green-St. Venant strain tensor $E = \frac{1}{2}(F^T F - \eta)$ where η denotes the material metric, transform covariantly under diffeomorphisms while preserving the causal structure of spacetime [8, 82].

The material manifold construction imposes specific constraints on the admissible configurations to ensure compatibility with relativistic causality and the initial-value formulation of the field equations. The requirement that the deformation gradient respect the Lorentzian signature of spacetime leads to the introduction of elasticity difference tensors that characterize the deviation between the material connection and the spacetime connection [70, 81]. These geometric objects facilitate the translation between continuum mechanical descriptions and gravitational physics, allowing the substrate's elastic properties to couple to the Einstein field equations through appropriately defined stress-energy measures that generalize classical elasticity to curved spacetime backgrounds [41, 56].

3.2. Elastic Deformation and Stress-Energy Tensors

The stress-energy content of the unified substrate derives from hyperelastic potentials expressed as functions of deformation invariants that remain scalar under general coordinate transformations. In the relativistic framework, the stress-energy tensor $T^{\mu\nu}$ emerges from the variational derivative of the elastic action with respect to the spacetime metric, incorporating contributions from the strain energy density and the kinetic energy of material displacements [15, 16]. The constitutive relations must respect the principle of material frame indifference adapted to general relativity, ensuring that the elastic response depends only on the strain invariants and not

on the absolute orientation of the material elements in spacetime [24, 82].

A distinctive feature of this formulation is the elasticity difference tensor, which quantifies the geometric incompatibility between the flat material connection and the curved spacetime connection induced by gravitational fields. This tensor appears in the modified Einstein field equations when the substrate’s elastic energy contributes to the total stress-energy budget, potentially altering the effective gravitational coupling at different scales [56, 81]. The Lagrangian approach to relativistic elasticity, grounded in DeWitt’s gauge-invariant action principle, provides a systematic methodology for deriving these modifications while ensuring energy-momentum conservation and hyperbolicity of the evolution equations necessary for well-posed initial value problems [8, 15].

3.3. Thermodynamic State Variables and Equations of State

The thermodynamic characterization of the unified substrate requires the specification of state variables including energy density ρ , pressure p , and entropy density s , which govern the medium’s response to expansion and compression across cosmological epochs. Unlike perfect fluid descriptions commonly employed in standard cosmology, the substrate’s equation of state must account for elastic contributions to the pressure tensor, leading to anisotropic stress components when the medium undergoes shear deformations [33, 59]. The fundamental relation between these thermodynamic quantities and the elastic potentials remains valid across different phases of the substrate, though the specific functional forms of the equations of state may vary as the medium undergoes subsequent transitions [34, 35].

The entropy contribution to the substrate’s stress-energy content plays a crucial role in determining the effective cosmological parameters, particularly when considering the dissipative processes associated with anelastic deformations. Relativistic anelasticity theory provides the framework for incorporating viscosity and plasticity into the substrate description, allowing for the modeling of irreversible processes that may generate entropy during cosmic evolution [33, 56]. These thermodynamic considerations establish the necessary foundation for understanding how the substrate’s material properties influence the expansion dynamics and structure formation, without invoking specific phase transition mechanisms that would characterize the cosmic history of the medium [58, 71].

4. Phase Transition Dynamics in the Substrate

The macroscopic behavior of the unified substrate is governed by temperature-dependent and density-dependent phase transitions that reconfigure its elastic and thermodynamic properties throughout cosmic history. Drawing upon frameworks from scalar-tensor gravity and relativistic

continuum mechanics, these transitions can be understood as spontaneous symmetry-breaking events in the substrate’s order parameter, analogous to phase transitions in condensed matter systems. The following subsections examine the critical phenomena driving these transitions, the specific mechanisms of condensation and vaporization within the substrate, and their chronological sequencing across cosmological epochs.

4.1. Critical Phenomena and Symmetry Breaking

Phase transitions in the substrate manifest as spontaneous symmetry-breaking processes wherein the order parameter acquires a non-zero vacuum expectation value, fundamentally altering the medium’s constitutive relations. Within scalar-tensor gravity frameworks, such transitions correspond to scalarization phenomena where the gravitational coupling becomes dynamically dependent on the scalar field configuration [29, 35]. The Landau-Ginzburg formalism provides a powerful descriptive tool for these transitions, wherein the free energy functional of the substrate expands in terms of the order parameter ψ as $F[\psi] = F_0 + a(T - T_c)\psi^2 + b\psi^4 + \dots$, with coefficients determining whether the transition proceeds continuously or discontinuously [83].

Recent analyses of scalarization mechanisms indicate that first-order phase transitions may dominate over second-order transitions in certain parameter regimes, introducing metastability and hysteresis effects in the substrate’s evolution [85]. During such transitions, topological defects—including domain walls, cosmic strings, and monopoles—may form as the substrate’s symmetry breaks, with the defect density determined by the correlation length and quench rate [48, 51]. These defects carry localized stress-energy contributions that could seed large-scale structure formation or primordial black holes, depending on the transition’s energy scale and the substrate’s elastic moduli [32, 57].

In the early universe context, radiatively induced symmetry breaking in grand unified theories provides a template for understanding substrate phase transitions, wherein thermal corrections to the effective potential drive the system from symmetric to broken phases [3, 38]. The gravitational phase transition hypothesis—wherein vacuum polarization effects induce a pure gravitational phase transition—further suggests that the substrate may undergo critical behavior independent of explicit scalar field potentials, potentially explaining the origin of cosmic structure through purely geometric mechanisms [74].

4.2. Condensation Mechanisms and Superfluidity

The transition to a condensed phase in the substrate corresponds to the formation of a Bose-Einstein condensate-like state characterized by a macroscopic occupation of the ground state and vanishing viscosity. Within the framework of superfluid dark matter models, this condensation

occurs when the temperature drops below a critical value T_c , inducing a phase change from a normal fluid to a superfluid with non-zero vacuum expectation value [9, 10]. The Landau two-fluid model adapted to relativistic contexts describes this state through the co-existence of superfluid and normal components, with the superfluid fraction exhibiting phonon-mediated interactions that modify gravitational dynamics at galactic scales [49, 50].

From the perspective of relativistic elasticity, the condensation transition modifies the substrate’s material manifold, introducing a strain tensor configuration that satisfies hyperelastic constitutive relations [8, 53]. The elastic difference tensor and deformation gradients associated with this phase transition generate effective stress-energy contributions that appear as additional matter components in the Einstein field equations [41, 81]. The transition kinetics are constrained by the requirement that the resulting superfluid phase maintain general covariance while exhibiting the hydrodynamic properties of an irrotational, inviscid fluid [33, 82].

Emergent gravity perspectives suggest that the condensation transition may correspond to the formation of a Bose-Einstein condensate of soft gravitons or quantum ether constituents, wherein the substrate’s collective behavior generates an effective metric perturbation interpreted as dark matter [17]. This corpuscular description aligns with the observation that superfluid dark matter models naturally interpolate between modified Newtonian dynamics at galactic scales and cold dark matter behavior on cosmological scales [20, 21].

4.3. Diffusive Transitions to Gas Phases

Conversely, the substrate may undergo diffusive transitions toward a dilute, weakly interacting gas phase characterized by negative pressure contributions that drive accelerated expansion. Environmental cosmic acceleration mechanisms in scalar-tensor theories demonstrate that density-triggered phase transitions can generate effective dark energy behavior by removing constant contributions to the Friedmann equation while preserving dynamic terms associated with the scalar field [25, 84]. The effective equation of state parameter w_{eff} in this phase may cross the phantom divide ($w < -1$) depending on the transition kinetics and the conformal frame chosen [67].

Observational constraints suggest that such transitions from matter-dominated to accelerated expansion occur at redshifts $z \approx 0.3$, consistent with scalar-tensor models where the scalar field drives the cosmological evolution through a non-standard kinetic Lagrangian [75, 76]. The diffuse gas phase maintains thermodynamic equilibrium through entropy production associated with the phase boundary, with the latent heat of transition potentially affecting the cosmic temperature evolution differently than standard recombination or decoupling processes.

Within the substrate framework, this gas phase represents a configuration where elastic moduli become negligible compared to thermal pressure, and the medium behaves as a nearly perfect fluid with equation of state $p = w\rho$ where $w \approx -1$ asymptotically. The transition to this phase may proceed through spinodal decomposition or nucleation processes, depending on whether the substrate’s free energy landscape supports metastable states between the condensed and gaseous configurations [4, 39].

4.4. Sequential Cosmological Phase Evolution

The cosmological history of the substrate comprises a sequence of phase transitions beginning with high-symmetry initial states and progressing through condensation and vaporization episodes. Dynamical systems analyses of scalar-tensor gravity demonstrate that such theories naturally accommodate radiation-dominated, matter-dominated, and exponentially accelerated epochs through sequential changes in the effective gravitational coupling [40]. The timing of these transitions determines the epoch-dependent equation of state and the expansion rate during critical periods such as nucleosynthesis and recombination.

Interrupted or aborted first-order phase transitions during reheating may produce macroscopic density perturbations that collapse into primordial black holes, providing a mechanism for matter genesis and baryogenesis through subsequent Hawking radiation [1, 2, 5]. These aborted transitions occur when bubble nucleation rates decrease as the universe cools, leaving remnant density fluctuations that accrete surrounding substrate material [65]. The resulting black hole population could contribute to dark matter density while generating baryon asymmetry through CP-violating emissions during evaporation [63, 64, 78].

The sequential evolution must satisfy constraints from big bang nucleosynthesis and cosmic microwave background observations, requiring that phase transitions altering the effective gravitational constant or expansion rate occur sufficiently early or sufficiently late to avoid disrupting light-element abundances [34, 36]. The substrate’s passage through distinct thermodynamic states—from potential initial high-temperature symmetric phases through superfluid condensation to late-time gaseous expansion—establishes a cosmological timeline wherein the effective degrees of freedom and elastic properties of spacetime evolve continuously yet critically at specific transition redshifts.

5. Emergence of Cosmic Components from Substrate Phases

The Spacetime Unified Substrate Configuration Theory posits that the cosmic dark sector does not comprise distinct fundamental fields but rather emerges as distinct thermodynamic configurations of a unified elastic medium. In this framework, the observed phenomenology of dark

matter and dark energy arises sequentially from spontaneous symmetry breaking and phase transitions within the substrate, wherein the same underlying material manifold manifests contrasting mechanical properties depending on its local thermodynamic state [38, 39, 48]. The condensation of the substrate into a coherent quantum state at galactic scales and its dilution into a rarefied gas at cosmological distances provide a unified conceptual basis for understanding the distinct gravitational effects attributed to these components [9, 25].

5.1. Dark Matter as a Condensed Superfluid Phase

The hypothesis that dark matter corresponds to a condensed superfluid phase of the unified substrate draws upon recent advances in relativistic superfluidity and emergent gravity phenomenology. Within this paradigm, the substrate undergoes a spontaneous symmetry breaking transition that establishes a non-zero vacuum expectation value, resulting in a Bose-Einstein condensate-like state characterized by long-range quantum coherence [9, 10]. This condensed phase exhibits vanishing viscosity and supports phonon-mediated interactions that modify gravitational dynamics at galactic scales, potentially obviating the need for particulate dark matter candidates [49, 50]. The superfluid description provides a natural explanation for the observed flatness of galactic rotation curves through the generation of a specific entanglement between the substrate’s elastic deformation and baryonic mass distributions [42].

The relativistic elastic properties of the substrate furnish the geometric framework necessary to describe this condensed state without invoking external dark matter fields. Drawing upon Souriau’s relativistic hyperelasticity [53], the superfluid phase can be characterized by specific deformation invariants that couple to spacetime curvature through modified stress-energy tensors. Gravitational wave propagation through this superfluid medium exhibits characteristic deviations from vacuum general relativity, including modified dispersion relations and effective refractive indices that serve as observational discriminants [20, 21]. These signatures suggest that the superfluid substrate phase behaves dynamically distinct from collisionless cold dark matter, particularly in systems undergoing mergers or exhibiting strong gravitational fields [20].

5.2. Dark Energy as a Diffuse Gas Phase

Conversely, the late-time accelerated expansion of the universe may originate from the substrate’s transition to a diffuse, weakly interacting gas phase characterized by negative pressure contributions to the cosmological stress-energy budget. This gaseous configuration emerges from a density-triggered phase transition in the late universe, wherein dynamic terms in the scalar-tensor sector generate effective cosmic acceleration without invoking a cosmological constant [25, 84]. The equation of state of this diffuse phase asymptotically approaches $w \approx -1$, effec-

tively mimicking the observed dark energy component while retaining the capacity for dynamical evolution across cosmic time [75, 76].

The thermodynamic properties of this gas phase are governed by the substrate’s elastic modulus and the specific entropy of the phase transition, parameters that determine the onset and duration of the accelerated expansion epoch. Self-accelerating solutions within scalar-tensor frameworks demonstrate that the transition from matter-dominated to accelerated expansion occurs naturally when the substrate’s effective pressure becomes sufficiently negative [6]. Notably, the DBI scalar field formalism suggests that this gaseous phase may remain approximately constant or slowly varying, providing a unified description where the same substrate component responsible for dark matter condensation at high densities transitions to dark energy-like behavior at cosmological scales [18, 19].

5.3. Weak Coupling Hypothesis and Baryonic Interactions

A critical component of the substrate theory is the hypothesis that conventional baryonic matter couples only weakly to the underlying elastic medium at astrophysical scales, ensuring that standard model particles follow approximately geodesic trajectories consistent with general relativity. This weak coupling mechanism operates analogously to screening mechanisms in extended scalar-tensor theories, where the interaction strength between matter and the substrate scalar field becomes suppressed in high-density environments [29, 34]. The negligible direct interaction preserves local Lorentz invariance and ensures consistency with solar system constraints on post-Newtonian parameters, while allowing the substrate to evolve independently according to its own elastic and thermodynamic equations of state [23].

The specific nature of this weak coupling remains an active area of theoretical investigation, as the microscopic interaction Lagrangian between baryonic fields and the substrate has yet to be fully specified within the relativistic elasticity framework. Current scalar-tensor constraints suggest that any coupling must respect stringent bounds from nucleosynthesis and cosmic microwave background observations, limiting the range of permissible coupling functions [23, 34]. While this hypothesis successfully decouples the substrate’s phase transition dynamics from baryonic physics in the weak-field regime, the precise mechanism by which matter fields source the substrate deformation—and the potential for strong-field deviations—requires further elaboration within the continuum mechanical description [29].

6. Cosmological Evolution and Expansion Dynamics

The Spacetime Unified Substrate Configuration Theory (SPT) posits that cosmic history manifests as a sequence of configurational changes in the unified elastic substrate, wherein distinct

cosmological epochs correspond to dominant phases of the medium. This framework reinterprets standard cosmological evolution not as the dynamics of disparate fluid components within a fixed spacetime geometry, but as the thermodynamic evolution of the substrate itself, punctuated by critical transitions that restructure its stress-energy content. The following subsections examine the implications of this paradigm for early universe phenomenology, the onset of accelerated expansion, and the recovery of general relativistic behavior at appropriate scales.

6.1. Early Universe Phase Structure and Primordial Perturbations

Early universe dynamics within SPT are characterized by the formation of topological defects and primordial black holes (PBHs) resulting from first-order or interrupted phase transitions in the substrate. The theory suggests that during reheating, aborted phase transitions—wherein nucleated bubbles of the condensed phase shrink and vanish as temperature decreases—generate macroscopic density perturbations that subsequently collapse into PBHs [1, 2]. These substrate-induced black holes may provide mechanisms for baryogenesis through CP-violating Hawking radiation emitted after the electroweak phase transition, potentially explaining the observed matter-antimatter asymmetry without requiring conventional sphaleron processes [63, 77, 78]. Furthermore, symmetry-breaking transitions in the early substrate may produce stable topological defects such as domain walls or cosmic strings, whose density and distribution depend on the order parameter dynamics during the transition [38, 48, 51].

The gravitational influence of these defects on structure formation presents both constraints and predictive signatures for the model, particularly given that defect-based cosmologies must reconcile observed cosmic microwave background anisotropies with inflationary predictions [32]. Additionally, gravitational phase transitions driven by vacuum-polarization effects in the early universe may offer alternative pathways for generating large-scale structure, though such mechanisms require specific parameter ranges within the gravitational Lagrangian to remain viable [74]. The formation of primordial black holes from bubble wall interactions during first-order transitions provides a specific testable prediction, linking the nucleation rate and bubble dynamics directly to the resulting black hole mass distribution and abundance [5].

6.2. Transition to Accelerated Expansion

The transition from matter-dominated to accelerated expansion in SPT emerges dynamically through a density-triggered phase transition in the dark sector, wherein the substrate transitions to a diffuse gas phase characterized by negative pressure contributions. Scalar-tensor gravity frameworks provide the mathematical machinery for modeling such transitions, wherein the scalar field governing gravitational coupling undergoes spontaneous scalarization that drives

cosmic acceleration without invoking a cosmological constant [75, 76]. Recent extensions demonstrate that first-order phase transitions in scalar-tensor theories—rather than second-order transitions—may dominate the late-time dynamics, generating metastable configurations that affect the effective equation of state [83, 85]. Environmental cosmic acceleration mechanisms, wherein constant potential contributions effectively degravitate leaving only dynamic terms to drive expansion, offer a specific realization consistent with late-time cosmological observations [25, 84].

The effective equation of state parameter w_{eff} may evolve dynamically across the phantom divide ($w < -1$) depending on the kinetics of the substrate’s gas-phase transition, providing phenomenological flexibility to accommodate diverse expansion histories [67]. Self-accelerating solutions within scalar-tensor frameworks demonstrate that cosmic expansion can be driven by the scalar field’s kinetic structure while maintaining compatibility with background evolution constraints [6]. Dynamical system analyses of scalar-tensor representations reveal that such theories permit sequences of cosmological phases—radiation-dominated, matter-dominated, and exponentially accelerated epochs—that align with Planck satellite measurements when appropriately parameterized [40].

6.3. Asymptotic Consistency with General Relativity

Ensuring consistency with local gravity constraints requires that SPT recover general relativistic predictions in the weak-field limit and within solar system scales, where post-Newtonian parameters tightly constrain deviations from Einstein gravity. Scalar-tensor theories generically predict variations in the gravitational constant G and deviations in post-Newtonian parameters such as γ and β , necessitating careful tuning of the conformal coupling to satisfy observational bounds from lunar laser ranging and binary pulsar systems [34, 36]. The weak coupling hypothesis posited in SPT—wherein baryonic matter interacts negligibly with the substrate at astrophysical scales—may naturally suppress fifth-force effects, preserving geodesic motion for standard matter while allowing the substrate to evolve independently.

However, the frame dependence of cosmological observables in scalar-tensor gravity necessitates consistent interpretation within specific conformal frames, as the Jordan and Einstein frames yield differing effective parameters that must be reconciled when comparing cosmological predictions with local measurements [46, 47]. Screening mechanisms or specific choices of coupling functions may ensure that the theory satisfies local gravity tests while allowing significant deviations at cosmological scales [69]. The recovery of general relativistic behavior in the strong-field regime and during the matter-dominated era provides essential constraints on the allowed parameter space of substrate elastic moduli and phase transition critical points [68].

7. Observational Constraints and Phenomenological Tests

The Spacetime Unified Substrate Configuration Theory generates specific empirical signatures that can be evaluated against current cosmological observations, particularly concerning the Hubble tension, large-scale structure formation, gravitational wave propagation, and laboratory analogues of substrate dynamics. Unlike phenomenological dark sector models that treat cosmological parameters as fixed or slowly varying background quantities, the substrate framework predicts redshift-dependent couplings and phase-dependent mechanical properties that may leave distinctive imprints across multiple observational channels. The following subsections examine the compatibility of these predictions with existing data while identifying potential discriminatory tests that could validate or falsify the substrate hypothesis.

7.1. Implications for the Hubble Tension

The persistent discrepancy between early-universe CMB constraints and late-universe distance ladder measurements of the Hubble constant represents a critical challenge that scalar-tensor substrate models may address through redshift-dependent effective gravitational couplings. Current observations reveal a statistically significant tension exceeding 5σ between Planck CMB predictions and local supernova-calibrated measurements, with the latter yielding values around $73 - 74 \text{ km s}^{-1} \text{ Mpc}^{-1}$ while CMB data prefer approximately $67 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [44, 45, 72]. Within the substrate framework, sequential phase transitions in the unified medium could generate an evolving effective Newton constant or modified expansion dynamics that manifest as apparent variations in H_0 across cosmic time, potentially reconciling these measurements without invoking systematic errors in either dataset [22, 79].

The specific mechanism involves density-triggered transitions in the substrate’s scalar configuration that alter the Friedmann equation’s effective energy density contributions during different cosmological epochs. Recent analyses suggest that the Hubble tension may be narrowed to discrepancies between Planck data and the first two rungs of the distance ladder, indicating that intermediate-redshift systematics alone cannot explain the divergence [44, 45]. If the substrate undergoes a late-time phase transition associated with the onset of dark energy dominance, the resulting environmental cosmic acceleration could modify distance-redshift relations in a manner that mimics local variations in H_0 while preserving early-universe physics [25, 84]. However, current scalar-tensor implementations capable of addressing the tension often introduce additional uncertainties regarding conformal frame dependencies and the specific coupling functions governing the substrate’s equation of state [46, 47, 73].

7.2. Structure Formation and Power Spectrum

The identification of dark matter with a condensed superfluid phase of the substrate fundamentally alters the predicted spectrum of primordial density perturbations and subsequent structure formation compared to standard cold dark matter paradigms. Superfluid dark matter models characterized by phonon-mediated interactions exhibit distinct sound speeds and Jeans lengths that suppress small-scale power, potentially resolving persistent discrepancies between CDM predictions and observed galactic structures while maintaining consistency with large-scale clustering [9, 10, 42]. The effective equation of state of the condensed substrate depends critically on the quantum pressure and self-interaction strength of the underlying scalar field, parameters that directly influence the cutoff scale in the matter power spectrum and the formation of galactic cores [9].

Topological defects emerging from symmetry-breaking phase transitions in the early substrate may additionally contribute to structure formation through distinct mechanisms from standard inflationary perturbations. Domain walls, cosmic strings, and monopoles generated during substrate condensation could seed density inhomogeneities that evolve into large-scale structure, though numerical analyses of CMB anisotropies indicate that purely defect-based models struggle to reproduce the observed acoustic peak structure without inflationary contributions [32, 51]. The substrate theory potentially occupies an intermediate position wherein defect formation accompanies but does not dominate primordial perturbations, with the superfluid phase later modulating structure growth through its modified gravitational coupling to baryonic matter [49, 50].

7.3. Gravitational Wave Phenomenology

Gravitational wave propagation through the substrate medium provides a novel probe of the theory’s phase structure and mechanical properties, with superfluid dark matter phases predicting measurable deviations from general relativistic wave dynamics. In Bose-Einstein condensate dark matter scenarios analogous to the substrate’s condensed phase, gravitational waves experience an effective refractive index modification that alters their propagation speed relative to the vacuum case, potentially producing frequency-dependent time delays or polarization mixing detectable by pulsar timing arrays and interferometric observatories [20, 21]. The magnitude of these effects scales with the superfluid density and phonon coupling parameters, offering a direct test of the substrate’s elastic properties at astrophysical scales [21].

Beyond superfluid phase effects, first-order phase transitions in the substrate may generate stochastic gravitational wave backgrounds through bubble nucleation and collision dynamics, analogous to early universe phase transitions in standard cosmology. Such transition-induced

gravitational radiation would exhibit characteristic spectral signatures distinct from astrophysical foregrounds, potentially detectable by future space-based interferometers [1, 5]. The damping of gravitational waves due to substrate viscosity or energy dissipation during propagation through the diffuse gas phase (associated with dark energy) could further distinguish the model from Λ CDM predictions, though current observational constraints on the graviton speed and damping remain compatible with general relativity at the percent level [27, 28].

7.4. Analogue Gravity Experimental Parallels

Laboratory analogue systems provide essential empirical constraints on the substrate hypothesis by realizing hydrodynamic and quantum fluid configurations that mimic the geometric and thermodynamic properties of the hypothesized spacetime medium. Bose-Einstein condensate experiments and hydrodynamic flows can generate acoustic metrics with horizons and ergoregions, enabling experimental investigation of Hawking radiation, superradiance, and emergent gravity phenomena that parallel predicted substrate behaviors [13, 14, 54]. These analogues demonstrate that effective spacetime geometries can emerge from underlying fluid dynamics, supporting the conceptual foundation of substrate-based cosmology while providing quantitative bounds on the transport coefficients and equation of state parameters relevant to the unified medium [28, 55].

Recent advances in analogue gravity have extended beyond linear perturbations to include nonlinear wave dynamics and spinning black hole analogues, offering increasingly sophisticated platforms for testing substrate phase transitions. Experimental realizations of acoustic Kerr metrics through Berry curvature effects in anomalous Hall systems and rotating fluid configurations establish that horizon formation and ergoregion physics depend sensitively on the underlying medium's equation of state and viscosity [30, 37, 62]. While current analogue experiments cannot replicate the full relativistic regime of cosmic substrate dynamics, observations of Hawking temperature measurements and entanglement entropy in these systems constrain the possible microphysical properties of any emergent spacetime medium, including the elastic moduli and critical exponents governing phase transitions in the substrate [11, 27, 61].

8. Theoretical Consistency and Limitations

The internal consistency of the Spacetime Unified Substrate Configuration Theory hinges upon its compatibility with fundamental symmetry principles, its behavior under conformal transformations, and the naturalness of its parameter space. While the framework offers a unified description of cosmic components through elastic phase transitions, it must satisfy stringent theoretical constraints to remain physically viable across relativistic and cosmological scales.

8.1. Preservation of Lorentz Covariance

Local Lorentz invariance must be maintained despite the substrate’s elastic properties and material constitution. The general covariant formulation of relativistic hyperelasticity, as developed by Souriau and extended in modern continuum mechanics, provides the necessary geometric machinery to ensure that constitutive relations and strain measures transform appropriately under local Lorentz transformations [8, 53]. The elasticity difference tensor and deformation gradients must respect the causal structure of spacetime, ensuring that observable physics remains independent of the choice of local inertial frame and that the substrate does not introduce preferred directions that would violate relativistic covariance.

The hyperbolic nature of the relativistic elastic field equations ensures that causality is preserved and that perturbations propagate within the light cone [82]. This mathematical property guarantees that the stress-energy content derived from elastic potentials does not permit superluminal signaling, thereby safeguarding local Lorentz covariance. However, the specific form of the elastic potential and the resulting equations of state must be carefully constructed to avoid instabilities or acausal behavior that could emerge from higher-order gradient corrections or non-linear elastic effects at extreme energy densities.

8.2. Conformal Frame Dependence and Observable Equivalence

The scalar-tensor representation of the substrate theory admits distinct conformal frames—specifically the Jordan and Einstein frames—that yield different effective gravitational couplings and cosmological dynamics. Observables interpreted within specific frames produce quantitatively distinct predictions for fundamental parameters, including the Hubble constant and the effective equation of state, underscoring the necessity of consistent frame usage when analyzing cosmological evolution [46, 47]. The frame-dependent nature of these interpretations complicates the reconciliation of early and late universe measurements, as identical physical configurations may appear discrepant when viewed through different conformal representations.

The question of whether cosmological observables are genuinely frame-independent remains contested within the broader scalar-tensor literature; while some geometric quantities transform trivially between frames, others exhibit significant dependence that affects the inferred expansion history and structure formation [34, 35]. This conformal inequality necessitates that all observational comparisons be conducted within a single, consistently defined frame to avoid physically contradictory conclusions. The substrate theory must therefore specify its physical frame unambiguously or demonstrate that its phase transition dynamics yield frame-invariant predictions for measurable quantities.

8.3. Parameter Space and Fine-Tuning Challenges

The sequential phase transitions hypothesized to generate dark matter and dark energy components depend sensitively on critical temperatures, nucleation rates, and elastic moduli that may require substantial fine-tuning to match observed cosmological epochs. The Landau theory of phase transitions, applied to the substrate’s symmetry-breaking mechanisms, suggests that the specific critical points and bubble nucleation dynamics must align precisely with constraints from primordial nucleosynthesis and recombination to avoid disrupting standard cosmic evolution [38, 51]. This sensitivity raises concerns regarding the naturalness of the initial conditions and the theoretical justification for the specific values of transition parameters that enable the substrate to mimic observed dark sector behavior.

In scalar-tensor extensions of the framework, the coupling functions and potentials that drive spontaneous scalarization admit significant arbitrariness, with viable cosmological models occupying narrow regions of parameter space that satisfy both local gravity tests and large-scale structure formation [35, 83]. The dependence of the cosmic acceleration transition on these finely-tuned parameters, particularly the critical density thresholds for phase transitions and the elastic response coefficients, introduces a degree of theoretical contrivance that challenges the model’s predictive power. Without additional symmetry principles or dynamical mechanisms to fix these parameters, the substrate theory may suffer from the same fine-tuning criticisms that afflict other modified gravity and dark energy models.

9. Discussion

The Spacetime Unified Substrate Configuration Theory (SPT) positions itself as a continuum-mechanical alternative to particle-based dark sector models, requiring careful comparative analysis against established cosmological paradigms to assess its theoretical viability and distinctiveness.

9.1. Comparison with Lambda Cold Dark Matter

The standard Λ CDM model posits dark matter as collisionless cold particles and dark energy as a cosmological constant, representing fundamentally distinct sectors without dynamical connection [43, 79]. In contrast, SPT unifies these components through phase transitions of a single elastic substrate, potentially offering a more fundamental origin wherein both phenomena emerge from the same underlying medium [29, 35]. However, this unification introduces additional theoretical complexity through the requirement of specifying elastic moduli, phase transition kinetics, and substrate thermodynamics that Λ CDM avoids by construction [26, 34].

While Λ CDM successfully reproduces cosmic microwave background anisotropies and large-scale structure formation through empirical parameter fitting [69], SPT must demonstrate equivalent observational consistency through the derived equations of state from substrate elasticity [47, 67]. The conformal frame dependence inherent in scalar-tensor extensions of gravity presents particular challenges for substrate models, as cosmological observables interpreted in different frames yield divergent parameter constraints [46, 47], necessitating careful frame selection to maintain consistency with post-Newtonian solar system tests [34].

9.2. Distinctions from Modified Newtonian Dynamics and Superfluid Models

Existing superfluid dark matter theories provide a partial unification by treating dark matter as a Bose-Einstein condensate that generates MOND-like phenomenology at galactic scales while recovering Λ CDM behavior cosmologically [9, 10]. However, these models typically retain dark energy as a separate cosmological constant or quintessence field [18, 19], whereas SPT uniquely derives both dark matter and dark energy from different phases of the identical substrate medium. Furthermore, while MONDian gravity emerges dynamically in superfluid models through specific phonon-mediated interactions [49, 50], SPT’s weak coupling hypothesis posits a fundamentally different interaction mechanism wherein baryonic matter couples negligibly to the substrate at astrophysical scales, preserving geodesic motion without invoking the specific acceleration scales characteristic of MOND [80].

The tensor-vector-scalar frameworks that attempt to relativistically extend MOND face persistent challenges in reconciling local gravity constraints with cosmic evolution [12, 66], issues that SPT potentially circumvents through its elastic continuum formulation. Unlike scalar-tensor theories that require specific screening mechanisms to hide fifth forces [17], the substrate’s weak coupling to baryonic matter operates through a distinct material interaction hypothesis rather than dynamical screening. Observational discrimination between SPT and superfluid dark matter may ultimately depend on gravitational wave propagation characteristics, as superfluid phases predict specific speed deviations for gravitational waves [20, 21], whereas the elastic substrate may exhibit frequency-dependent dispersion relations tied to its constitutive parameters.

9.3. Future Theoretical Directions

Several foundational questions remain unresolved within the substrate framework, particularly regarding the quantum mechanical treatment of the unified medium. While current formulations rely on classical relativistic elasticity [8, 53], a complete theory requires the quantization of substrate fluctuations and the derivation of an effective field-theoretic Lagrangian that reduces

to the continuous medium description in the hydrodynamic limit [24, 59]. The nature of the weak coupling mechanism between baryonic matter and the substrate demands specification at the level of quantum field interactions, potentially involving Yukawa-type couplings or topological constraints that remain to be articulated [23, 31].

Future developments must address the parameter space constraints identified in scalar-tensor analogs, where fine-tuning of phase transition critical points threatens theoretical naturalness [83, 85]. Numerical simulations of cosmological structure formation within the substrate framework could provide discriminatory power regarding the sound speed and Jeans length of the condensed phase [42], while laboratory analogue systems utilizing Bose-Einstein condensates [14, 28] or hydrodynamic flows [13, 54] may constrain the elastic properties hypothesized for the cosmological substrate. The reconciliation of first-order phase transition dynamics with observed cosmic evolution [1, 2] represents a critical direction for establishing the model’s consistency with early universe nucleosynthesis and recombination epochs.

10. Conclusion

This survey has examined the Spacetime Unified Substrate Configuration Theory as a novel approach to cosmological unification, wherein the dynamical behavior of a relativistic elastic substrate replaces conventional dark sector phenomenology. By interpreting dark matter as a condensed superfluid phase and dark energy as a diffuse gas phase arising from sequential symmetry-breaking transitions, the framework offers a mechanically coherent alternative to the standard cosmological model’s disparate components. The theory’s emphasis on continuous medium mechanics provides a mathematically structured vocabulary for describing cosmic evolution, while the weak coupling hypothesis ensures asymptotic consistency with general relativity in well-tested regimes.

Despite its conceptual elegance, the substrate model remains highly speculative, requiring substantial theoretical refinement to address fundamental questions regarding the quantum field-theoretic foundation of the elastic medium, the specific mechanism governing baryonic coupling suppression, and the fine-tuning of phase transition critical points across cosmological history. Future investigations must prioritize the development of predictive phenomenology distinguishable from cold dark matter and cosmological constant paradigms, particularly regarding structure formation power spectra, gravitational wave propagation characteristics, and potential resolutions to the Hubble tension. Laboratory analogue systems and next-generation observational facilities may ultimately determine whether this substrate unification represents a physical reality or a mathematical metaphor for deeper gravitational principles.

References

- [1] Wen-Yuan Ai, Lucien Heurtier, and Tae Hyun Jung. Primordial black holes from an aborted phase transition. 2024. URL <https://arxiv.org/pdf/2409.02175>.
- [2] Wen-Yuan Ai, Lucien Heurtier, and Tae Hyun Jung. Primordial black holes from an interrupted phase transition. Physical Review D, 113(2), January 2026. ISSN 2470-0029. doi: 10.1103/9yzd-r6kc. URL <http://dx.doi.org/10.1103/9yzd-r6kc>.
- [3] Andreas Albrecht and Paul J. Steinhardt. Cosmology for grand unified theories with radiatively induced symmetry breaking. Physical Review Letters, 48(17):1220–1223, April 1982. ISSN 0031-9007. doi: 10.1103/physrevlett.48.1220. URL <http://dx.doi.org/10.1103/PhysRevLett.48.1220>.
- [4] Luca Amendola, Carlo Baccigalupi, Marcelo Gleiser, and Franco Occhionero. First order phase transitions in cosmology. 1999. URL <https://linkinghub.elsevier.com/retrieve/pii/S1384107699000196>.
- [5] Michael J. Baker, Moritz Breitbach, Joachim Kopp, and Lukas Mittnacht. Primordial black holes from first-order cosmological phase transitions. 2021. URL <https://arxiv.org/pdf/2105.07481>.
- [6] Gabriela Barenboim and Joseph Lykken. Self-accelerating solutions of scalar-tensor gravity. 2007. URL <https://arxiv.org/pdf/0711.3653>.
- [7] Robert Beig. Nonrelativistic and relativistic continuum mechanics. 2004. URL <https://arxiv.org/pdf/gr-qc/0403073>.
- [8] Robert Beig and Bernd G. Schmidt. Relativistic elasticity. 2002. URL <https://arxiv.org/pdf/gr-qc/0211054>.
- [9] Lasha Berezhiani and Justin Khoury. Theory of dark matter superfluidity. Physical Review D, 92(10), November 2015. ISSN 1550-2368. doi: 10.1103/physrevd.92.103510. URL <http://dx.doi.org/10.1103/PhysRevD.92.103510>.
- [10] Lasha Berezhiani and Justin Khoury. Theory of dark matter superfluidity. 2015. URL <https://arxiv.org/pdf/1507.01019>.
- [11] Swastik Bhattacharya and S. Shankaranarayanan. How emergent is gravity? International Journal of Modern Physics D, 24(12):1544005, October 2015. ISSN 1793-6594. doi: 10.1142/s0218271815440058. URL <http://dx.doi.org/10.1142/S0218271815440058>.

- [12] Luc Blanchet and Constantinos Skordis. Khronon-tensor theory reproducing mond and the cosmological model. 2025. URL <https://arxiv.org/pdf/2507.00912v1>.
- [13] Samuel L. Braunstein, Mir Faizal, Lawrence M. Krauss, Francesco Marino, and Naveed A. Shah. Analogue simulations of quantum gravity with fluids. 2023. URL <https://www.nature.com/articles/s42254-023-00630-y>?
- [14] Samuel L. Braunstein, Mir Faizal, Lawrence M. Krauss, Francesco Marino, and Naveed A. Shah. Analogue simulations of quantum gravity with fluids. 2024. URL <https://arxiv.org/pdf/2402.16136>.
- [15] J. David Brown. Elasticity theory in general relativity. 2020. URL <https://arxiv.org/pdf/2004.03641>.
- [16] J David Brown. Elasticity theory in general relativity. Classical and Quantum Gravity, 38(8):085017, March 2021. ISSN 1361-6382. doi: 10.1088/1361-6382/abe1ff. URL <http://dx.doi.org/10.1088/1361-6382/abe1ff>.
- [17] M. Cadoni, R. Casadio, A. Giusti, and M. Taveri. Emergence of a dark force in corpuscular gravity. Physical Review D, 97(4), February 2018. ISSN 2470-0029. doi: 10.1103/physrevd.97.044047. URL <http://dx.doi.org/10.1103/PhysRevD.97.044047>.
- [18] Rong-Gen Cai and Shao-Jiang Wang. Dark matter superfluid and dbi dark energy. 2015. URL <https://arxiv.org/pdf/1511.00627>.
- [19] Rong-Gen Cai and Shao-Jiang Wang. Dark matter superfluid and dbi dark energy. Physical Review D, 93(2), January 2016. ISSN 2470-0029. doi: 10.1103/physrevd.93.023515. URL <http://dx.doi.org/10.1103/PhysRevD.93.023515>.
- [20] Rong-Gen Cai, Tong-Bo Liu, and Shao-Jiang Wang. Gravitational wave as probe of superfluid dark matter. 2017. URL <https://arxiv.org/pdf/1710.02425>.
- [21] Rong-Gen Cai, Tong-Bo Liu, and Shao-Jiang Wang. Gravitational wave as probe of superfluid dark matter. Physical Review D, 97(2), January 2018. ISSN 2470-0029. doi: 10.1103/physrevd.97.023027. URL <http://dx.doi.org/10.1103/PhysRevD.97.023027>.
- [22] Salvatore Capozziello, Giuseppe Sarracino, and Alessandro D.A.M. Spallicci. Questioning the h_0 tension via the look-back time. 2023. URL <https://arxiv.org/pdf/2302.13671>.
- [23] R. Catena, N. Fornengo, A. Masiero, M. Pietroni, and F. Rosati. Dark matter relic abundance and scalar-tensor dark energy. 2004. URL <https://arxiv.org/pdf/astro-ph/0403614>.

- [24] Mina Chapon, Lionel Darondeau, Rodrigue Desmorat, Clément Ecker, and Boris Kolev. Relativistic second gradient theory of continuous media. 2025. URL <https://arxiv.org/pdf/2507.13030v1>.
- [25] Ø. Christiansen, F. Hassani, and D.F. Mota. Environmental cosmic acceleration from a phase transition in the dark sector. *Journal of Cosmology and Astroparticle Physics*, 2025 (01):043, January 2025. ISSN 1475-7516. doi: 10.1088/1475-7516/2025/01/043. URL <http://dx.doi.org/10.1088/1475-7516/2025/01/043>.
- [26] E.V. Chubaryan, R.M. Avakyan, G.G. Harutyunyan, and A.S. Piloyan. Role of scalar fields in cosmological models. 2009. URL <https://linkinghub.elsevier.com/retrieve/pii/S0273117709005262>.
- [27] Chiara Coviello. Gravitational waves and black hole perturbations in acoustic analogues. 2024. URL <https://arxiv.org/pdf/2409.15403>.
- [28] Chiara Coviello, Maria Luisa Chiofalo, Dario Grasso, Stefano Liberati, Massimo Mannarelli, and Silvia Trabucco. Gravitational waves and black hole perturbations in acoustic analogues. 2024. URL <https://arxiv.org/pdf/2410.00264>.
- [29] Marco Crisostomi, Kazuya Koyama, and Gianmassimo Tasinato. Extended scalar-tensor theories of gravity. 2016. URL <https://arxiv.org/pdf/1602.03119>.
- [30] Surojit Dalui, Arpan Krishna Mitra, Deeshani Mitra, and Subir Ghosh. Spinning black hole in a fluid. 2024. doi: 10.1103/PhysRevD.109.064055. URL <https://journals.aps.org/prd/abstract/10.1103/PhysRevD.109.064055>
- [31] V. I. Denissov and B. V. Mehta. The massless scalar field as an alternative to dark matter. *Astronomical & Astrophysical Transactions*, 14(3):165–170, December 1997. ISSN 1476-3540. doi: 10.1080/10556799708202986. URL <http://dx.doi.org/10.1080/10556799708202986>.
- [32] R. Durrer, M. Kunz, and A. Melchiorri. Cosmic structure formation with topological defects. 2002. URL <https://linkinghub.elsevier.com/retrieve/pii/S0370157302000145>.
- [33] M. Epstein, D. A. Burton, R. W. Tucker, D.A. Burton, and R.W. Tucker. Relativistic anelasticity. 2005. URL <https://arxiv.org/pdf/gr-qc/0510113>.
- [34] Gilles Esposito-Farese. Scalar-tensor theories and cosmology. 2003. URL <https://arxiv.org/pdf/gr-qc/0306018>.

- [35] Valerio Faraoni. *Cosmology in Scalar-Tensor Gravity*. Springer Netherlands, 2004. ISBN 9781402019890. doi: 10.1007/978-1-4020-1989-0. URL <http://dx.doi.org/10.1007/978-1-4020-1989-0>.
- [36] Yasunori Fujii and Kei ichi Maeda. *The scalar-tensor theory of gravitation*. 2003. URL <https://www.cambridge.org/core/product/identifier/9780511535093/type/book>.
- [37] Rohith Ghosh, Souvik Ghose, Biplab Raychaudhuri, Apashanka Das, and Tapas K. Das. Emergent gravity from nonlinear perturbation of spherical accretion with variable adiabatic index. 2026. URL <https://arxiv.org/pdf/2605.04158v1>.
- [38] Marcelo Gleiser. Phase transitions in the universe. *Contemporary Physics*, 39(4):239–253, 1998. ISSN 1366-5812. doi: 10.1080/001075198181937. URL <http://dx.doi.org/10.1080/001075198181937>.
- [39] Marcelo Gleiser. Phase transitions in the universe. 1998. URL <https://arxiv.org/pdf/hep-ph/9803291>.
- [40] Tiago B. Gonçalves, João Luís Rosa, and Francisco S. N. Lobo. Dynamical reconstruction of the Λ CDM model in scalar-tensor gravity. *Physical Review D*, 109(8), April 2024. ISSN 2470-0029. doi: 10.1103/physrevd.109.084008. URL <http://dx.doi.org/10.1103/PhysRevD.109.084008>.
- [41] Walter C. Hernandez. Elasticity theory in general relativity. *Physical Review D*, 1(4): 1013–1018, February 1970. ISSN 0556-2821. doi: 10.1103/physrevd.1.1013. URL <http://dx.doi.org/10.1103/PhysRevD.1.1013>.
- [42] Sabine Hossenfelder and Tobias Mistele. The milky way’s rotation curve with superfluid dark matter. 2020. URL <https://arxiv.org/pdf/2003.07324>.
- [43] Jian-Ping Hu and Fa-Yin Wang. Hubble tension: The evidence of new physics. 2023. URL <https://www.mdpi.com/2218-1997/9/2/94>.
- [44] Lu Huang, Rong-Gen Cai, Shao-Jiang Wang, Jian-Qi Liu, and Yan-Hong Yao. Narrowing down the hubble tension to the first two rungs of distance ladders. *Science China Physics*,

- Mechanics & Astronomy*, 68(8), May 2025. ISSN 1869-1927. doi: 10.1007/s11433-025-2641-2. URL <http://dx.doi.org/10.1007/s11433-025-2641-2>.
- [45] Lu Huang, Rong-Gen Cai, Shao-Jiang Wang, Jian-Qi Liu, and Yan-Hong Yao. Narrowing down the hubble tension to the first two rungs of distance ladders. 2025. doi: 10.1007/s11433-025-2641-2. URL <https://link.springer.com/article/10.1007/s11433-025-2641-2>.
 - [46] Young-Hwan Hyun, Yoonbai Kim, and Seokcheon Lee. Conformal frame dependence on cosmological observations in scalar-tensor theories of gravity. 2017. URL <https://arxiv.org/pdf/1712.09353>.
 - [47] Young-Hwan Hyun, Yoonbai Kim, and Seokcheon Lee. Conformal frame dependence on cosmological observations in scalar-tensor theories of gravity. 2019. doi: 10.3938/jkps.74.1101. URL <http://link.springer.com/10.3938/jkps.74.1101>.
 - [48] Michael Kachelriess. Phase transitions and topological defects. 2018. doi: 10.1093/oso/9780198802877.001.0001/oso-9780198802877-chapter-16. URL <https://oxford.universitypressscholarship.com/view/10.1093/oso/9780198802877.001.0001/oso-9780198802877-chapter-16>.
 - [49] Justin Khoury. Another path for the emergence of modified galactic dynamics from dark matter superfluidity. 2016. URL <https://arxiv.org/pdf/1602.05961>.
 - [50] Justin Khoury. Another path for the emergence of modified galactic dynamics from dark matter superfluidity. *Physical Review D*, 93(10), May 2016. ISSN 2470-0029. doi: 10.1103/physrevd.93.103533. URL <http://dx.doi.org/10.1103/PhysRevD.93.103533>.
 - [51] T. W. B. Kibble. *Phase Transitions in the Early Universe and Defect Formation*, page 1–26. Springer US, 1995. ISBN 9781461518839. doi: 10.1007/978-1-4615-1883-9_1. URL http://dx.doi.org/10.1007/978-1-4615-1883-9_1.
 - [52] Boris Kolev and Rodrigue Desmorat. Souriau’s relativistic general covariant formulation of hyperelasticity revisited. 2023. URL <https://arxiv.org/pdf/2301.09437>.
 - [53] Boris Kolev, Rodrigue Desmorat, B. Kolev, and R. Desmorat. Souriau’s general covariant formulation of relativistic hyperelasticity revisited. 2023. URL https://www.sciencedirect.com/science/article/pii/S0022509623002673?dgcid=rss_sd_all&.
 - [54] Lawrence M. Krauss, Francesco Marino, Samuel L. Braunstein, Mir Faizal, and Naveed A. Shah. Listening to quantum gravity? *International Journal of Modern Physics D*, 33(15), August 2023. ISSN 1793-6594. doi: 10.1142/s0218271824410062. URL <http://dx.doi.org/10.1142/S0218271824410062>.

- [55] Lawrence M. Krauss, Francesco Marino, Samuel L. Braunstein, Mir Faizal, and Naveed A. Shah. Listening to quantum gravity? 2024. URL <https://arxiv.org/pdf/2405.19385v1>.
- [56] Eun-Ho Lee. Relativistic constitutive modeling of inelastic deformation of continua moving in space-time. 2024. URL https://www.sciencedirect.com/science/article/pii/S1007570424000078?dgcid=rss_sd_all&.
- [57] S. Lola and G.G. Ross. Structure formation from unstable domain walls. 1993. URL <https://linkinghub.elsevier.com/retrieve/pii/055032139390176P>.
- [58] Enrico Massa. Relative continuum mechanics in general relativity: I. —kinematical foundations. *Annali di Matematica Pura ed Applicata*, 117(1):311–337, December 1978. ISSN 1618-1891. doi: 10.1007/bf02417897. URL <http://dx.doi.org/10.1007/BF02417897>.
- [59] Gérard A. Maugin. *Relativistic Continuum Mechanics: A 20th Century Adventure*, page 267–280. Springer Netherlands, 2013. ISBN 9789400763531. doi: 10.1007/978-94-007-6353-1_15. URL http://dx.doi.org/10.1007/978-94-007-6353-1_15.
- [60] Jianwei Mei. A hydrodynamical description of gravitational waves. *The European Physical Journal C*, 83(1), January 2023. ISSN 1434-6052. doi: 10.1140/epjc/s10052-022-11160-9. URL <http://dx.doi.org/10.1140/epjc/s10052-022-11160-9>.
- [61] Mordehai Milgrom. Massive particles in acoustic space-times: Emergent inertia and passive gravity. *Physical Review D*, 73(8), April 2006. ISSN 1550-2368. doi: 10.1103/physrevd.73.084005. URL <http://dx.doi.org/10.1103/PhysRevD.73.084005>.
- [62] Deeshani Mitra, Surojit Dalui, Subir Ghosh, and Arpan Krishna Mitra. Acoustic kerr metric in analogue gravity. 2023. URL <https://arxiv.org/pdf/2306.12201>.
- [63] Yukinori Nagatani. Black hole baryogenesis. 1998. URL <https://arxiv.org/pdf/hep-ph/9805455>.
- [64] Yukinori Nagatani. Electroweak domain wall by hawking radiation: Baryogenesis and dark matter from several hundred kg black holes. 2001. URL <https://arxiv.org/pdf/hep-ph/0104160>.
- [65] Mitesh Patel and George M. Fuller. Primordial black holes, phase transitions, and the fate of the universe. 2000. URL <https://arxiv.org/pdf/astro-ph/0003062>.
- [66] R. H. Sanders and R.H. Sanders. A tensor-vector-scalar framework for modified dynamics and cosmic dark matter. 2005. URL <https://arxiv.org/pdf/astro-ph/0502222>.

- [67] Emmanuel N. Saridakis and Minas Tsoukalas. Cosmology in new gravitational scalar-tensor theories. 2016. URL <https://arxiv.org/pdf/1601.06734>.
- [68] Meir Shimon. A globally unevolving universe. 2014. URL <https://arxiv.org/pdf/1412.1078>.
- [69] Diego Sáez-Gómez. Scalar-tensor theories and current cosmology. 2008. URL <https://arxiv.org/pdf/0812.1980>.
- [70] A. Tartaglia. Four dimensional elasticity and general relativity. 1995. URL <https://arxiv.org/pdf/gr-qc/9509043>.
- [71] Angelo Tartaglia, Ninfa Radicella, Jean-Michel Alimi, and André Fuözfa. From elastic continua to space-time. In AIP Conference Proceedings, page 1156–1163. AIP, 2010. doi: 10.1063/1.3462613. URL <http://dx.doi.org/10.1063/1.3462613>.
- [72] S. Taubenberger, S. H. Suyu, E. Komatsu, I. Jee, S. Birrer, V. Bonvin, F. Courbin, C. E. Rusu, A. J. Shajib, and K. C. Wong. The hubble constant determined through an inverse distance ladder including quasar time delays and type ia supernovae. 2019. doi: 10.1051/0004-6361/201935980. URL <https://www.aanda.org/10.1051/0004-6361/201935980>.
- [73] Rahul Kumar Thakur, Shashikant Gupta, Rahul Nigam, and PK Thiruvikraman. Investigating the hubble tension through hubble parameter data. Research in Astronomy and Astrophysics, 23(6):065017, May 2023. ISSN 2397-6209. doi: 10.1088/1674-4527/acd0e8. URL <http://dx.doi.org/10.1088/1674-4527/acd0e8>.
- [74] Igor’ I. Tkachev. Gravitational phase transition: An origin of the large-scale structure in the universe? 1992. doi: 10.1103/PhysRevD.45.R4367. URL <https://link.aps.org/doi/10.1103/PhysRevD.45.R4367>.
- [75] SANIL UNNIKRISHNAN and T. R. SESHADRI. Cosmic acceleration in a model of scalar-tensor gravitation. International Journal of Modern Physics D, 17(11):2007–2015, October 2008. ISSN 1793-6594. doi: 10.1142/s0218271808013674. URL <http://dx.doi.org/10.1142/S0218271808013674>.
- [76] Sanil Unnikrishnan, T. R. Seshadri, and T.R.Seshadri. Cosmic acceleration in a model of scalar-tensor gravitation. 2005. URL <https://arxiv.org/pdf/gr-qc/0511035>.
- [77] Niraj Upadhyay, Patrick Das Gupta, and R. P. Saxena. Baryogenesis from primordial black holes after the electroweak phase transition. Physical Review D, 60(6), August 1999. ISSN 1089-4918. doi: 10.1103/physrevd.60.063513. URL <http://dx.doi.org/10.1103/PhysRevD.60.063513>.

- [78] Niraj Upadhyay, Patrick Das Gupta, and R. P. Saxena. Baryogenesis from primordial blackholes after electroweak phase transition. 1999. URL <https://arxiv.org/pdf/astro-ph/9903253>.
- [79] Eleonora Di Valentino, Olga Mena, Supriya Pan, Luca Visinelli, Weiqiang Yang, Alessandro Melchiorri, David F. Mota, Adam G. Riess, and Joseph Silk. In the realm of the hubble tension – a review of solutions. 2021. URL <https://arxiv.org/pdf/2103.01183>.
- [80] Milovan Vasilic. Searching for mond in scalar-tensor theories of gravity. 2018. URL <https://arxiv.org/pdf/1803.01666>.
- [81] E. G. L. R. Vaz and Irene Brito. Analysing the elasticity difference tensor of general relativity. General Relativity and Gravitation, 40(9):1947–1966, February 2008. ISSN 1572-9532. doi: 10.1007/s10714-008-0615-7. URL <http://dx.doi.org/10.1007/s10714-008-0615-7>.
- [82] Michael Wernig-Pichler. Relativistic elastodynamics. 2006. URL <https://arxiv.org/pdf/gr-qc/0605025>.
- [83] Murat Özınan, Kıvanç İ. Ünlütürk, and Fethi M. Ramazanoğlu. Underlying mechanisms of phase transitions in scalar-tensor theories. 2026. URL <https://arxiv.org/pdf/2604.24867v1>.
- [84] Øyvind Christiansen, Farbod Hassani, and David F. Mota. Environmental cosmic acceleration from a phase transition in the dark sector. 2024. URL <https://arxiv.org/pdf/2405.00668>.
- [85] Kıvanç İ. Ünlütürk, Semih Tuna, Oğuzhan K. Yamak, and Fethi M. Ramazanoğlu. Nature of phase transitions and metastability in scalar-tensor theories. 2025. URL <https://arxiv.org/pdf/2502.01781v1>.