

# Network models: random graphs

Yury Dvorkin

February 11, 2020



### Empirical network features:

- ▶ Power-law (heavy-tailed) degree distribution
- ▶ Small average distance (graph diameter)
- ▶ Large clustering coefficient (transitivity)
- ▶ Giant connected component, hierarchical structure, etc

### Generative models:

- ▶ Random graph model (Erdos & Renyi, 1959)
- ▶ "Small world" model (Watts & Strogatz, 1998)
- ▶ Preferential attachment model (Barabasi & Albert, 1999)

Well-known property:

If  $a \circ b$  &  $b \circ c$ , then  $a \circ c$ ,

where  $\circ$  is a transitive operator. Example:

If  $a = b$  &  $b = c$ , then  $a = c$

Graph extension is for edges:

If nodes  $a \circ b$  & nodes  $b \circ c$ , then nodes  $a \circ c$ ,

where  $\circ$  is a transitive operator (e.g. connection) Graph example of transitivity:

- ▶ Full: a friend of my friend is my friend
- ▶ Partial: a friend of my friend is not necessarily my friend

Well-known property:

If  $a \circ b$  &  $b \circ c$ , then  $a \circ c$ ,

where  $\circ$  is a transitive operator. Example:

If  $a = b$  &  $b = c$ , then  $a = c$

Graph extension is for edges:

If nodes  $a \circ b$  & nodes  $b \circ c$ , then nodes  $a \circ c$ ,

where  $\circ$  is a transitive operator (e.g. connection) Graph example of transitivity:

- ▶ Full: a friend of my friend is my friend
- ▶ Partial: a friend of my friend is not necessarily my friend

Well-known property:

If  $a \circ b$  &  $b \circ c$ , then  $a \circ c$ ,

where  $\circ$  is a transitive operator. Example:

If  $a = b$  &  $b = c$ , then  $a = c$

Graph extension is for edges:

If nodes  $a \circ b$  & nodes  $b \circ c$ , then nodes  $a \circ c$ ,

where  $\circ$  is a transitive operator (e.g. connection) Graph example of transitivity:

- ▶ Full: a friend of my friend is my friend
- ▶ Partial: a friend of my friend is not necessarily my friend

Well-known property:

If  $a \circ b$  &  $b \circ c$ , then  $a \circ c$ ,

where  $\circ$  is a transitive operator. Example:

If  $a = b$  &  $b = c$ , then  $a = c$

Graph extension is for edges:

If nodes  $a \circ b$  & nodes  $b \circ c$ , then nodes  $a \circ c$ ,

where  $\circ$  is a transitive operator (e.g. connection) Graph example of transitivity:

- ▶ Full: a friend of my friend is my friend
- ▶ Partial: a friend of my friend is not necessarily my friend

How can transitivity be measured?

- ▶ Only relative to all two-edge paths within a given graph
- ▶ Using *closed* two-edge paths: closed path forms a loop of three nodes
- ▶ Clustering coefficient:

$$C = \frac{\text{N of closed two-edge paths}}{\text{N of all two-edge paths}}$$

- ▶ Interpretations:
  - ▶  $C = 0$ : full transitivity
  - ▶  $C = 1$ : radial/tree topology

How can transitivity be measured?

- ▶ Only relative to all two-edge paths within a given graph
- ▶ Using *closed* two-edge paths: closed path forms a loop of three nodes
- ▶ Clustering coefficient:

$$C = \frac{\text{N of closed two-edge paths}}{\text{N of all two-edge paths}}$$

- ▶ Interpretations:
  - ▶  $C = 0$ : full transitivity
  - ▶  $C = 1$ : radial/tree topology



How can transitivity be measured?

- ▶ Only relative to all two-edge paths within a given graph
- ▶ Using *closed* two-edge paths: closed path forms a loop of three nodes
- ▶ Clustering coefficient:

$$C = \frac{\text{N of closed two-edge paths}}{\text{N of all two-edge paths}}$$

- ▶ Interpretations:
  - ▶  $C = 0$ : full transitivity
  - ▶  $C = 1$ : radial/tree topology

### Why generative models?

- ▶ Move from analyzing statistical properties to analyzing network behavior?
- ▶ Need to simulate behavior given a property (e.g. power laws)
- ▶ Hence, random graphs as a starting point

Simple example: Given  $n$  nodes and  $m$  edges, but edges are random



- ▶ Can we select a subgraph?
- ▶ Can we represent a graph as an ensemble of subgraphs?
- ▶ How to obtain a probabilistic representation of  $G$  in terms of subgraphs?
  - ▶ Properties of the random graph = *statistical* properties of the ensemble of subgraphs

Simple example: Given  $n$  nodes and  $m$  edges, but edges are random



- ▶ Can we select a subgraph?
- ▶ Can we represent a graph as an ensemble of subgraphs?
- ▶ How to obtain a probabilistic representation of  $G$  in terms of subgraphs?
  - ▶ Properties of the random graph = *statistical* properties of the ensemble of subgraphs

Simple example: Given  $n$  nodes and  $m$  edges, but edges are random



- ▶ Can we select a subgraph?
- ▶ Can we represent a graph as an ensemble of subgraphs?
- ▶ How to obtain a probabilistic representation of  $G$  in terms of subgraphs?
  - ▶ Properties of the random graph = *statistical* properties of the ensemble of subgraphs

Simple example: Given  $n$  nodes and  $m$  edges, but edges are random



- ▶ Can we select a subgraph?
- ▶ Can we represent a graph as an ensemble of subgraphs?
- ▶ How to obtain a probabilistic representation of  $G$  in terms of subgraphs?
  - ▶ Properties of the random graph = *statistical* properties of the ensemble of subgraphs

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$ , Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
  - ▶ Probability distribution:

$$P(G) = \frac{1}{\binom{\frac{n}{2}}{m}}$$

- ▶ Diameter of the graph:

$$\langle d \rangle = \sum_G P(G) d(G), \text{ where } d(G) \text{ is given}$$

- ▶ Assume that instead of  $m$ , we parameterize the probability of edge  $p$

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$ , Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
  - ▶ Probability distribution:

$$P(G) = \frac{1}{\binom{\frac{n}{2}}{m}}$$

- ▶ Diameter of the graph:

$$\langle d \rangle = \sum_G P(G) d(G), \text{ where } d(G) \text{ is given}$$

- ▶ Assume that instead of  $m$ , we parameterize the probability of edge  $p$



Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$ , Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
  - ▶ Probability distribution:

$$P(G) = \frac{1}{\binom{\frac{n}{2}}{m}}$$

- ▶ Diameter of the graph:

$$\langle d \rangle = \sum_G P(G) d(G), \text{ where } d(G) \text{ is given}$$

- ▶ Assume that instead of  $m$ , we parameterize the probability of edge  $p$

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$ , Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
  - ▶ Probability distribution:

$$P(G) = \frac{1}{\binom{\frac{n}{2}}{m}}$$

- ▶ Diameter of the graph:

$$\langle d \rangle = \sum_G P(G) d(G), \text{ where } d(G) \text{ is given}$$

- ▶ Assume that instead of  $m$ , we parameterize the probability of edge  $p$

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$ , Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
  - ▶ Probability distribution with parameterized  $m$ :

$$P(G) = \frac{1}{\binom{\frac{n}{2}}{m}}$$

- ▶ Probability distribution with parameterized  $p$ : (**Erdos-Renyi random graph**):

$$P(G) = p^m (1 - p)^{\binom{n}{2} - m}$$

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$ , Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
  - ▶ Probability distribution with parameterized  $m$ :

$$P(G) = \frac{1}{\binom{\frac{n}{2}}{m}}$$

- ▶ Probability distribution with parameterized  $p$ : (**Erdos-Renyi random graph**):

$$P(G) = p^m (1 - p)^{\binom{n}{2} - m}$$

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$ , Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
  - ▶ Probability distribution with parameterized  $m$ :

$$P(G) = \frac{1}{\binom{\frac{n}{2}}{m}}$$

- ▶ Probability distribution with parameterized  $p$ : (**Erdos-Renyi random graph**):

$$P(G) = p^m(1 - p)^{\binom{n}{2} - m}$$

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$  Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,q}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
- ▶ What properties should a random graph have?
- ▶ Number of edges ( $m$ ) and of degrees ( $k$ ), as well as density ( $\rho$ )

$$\langle m \rangle = pN = p \frac{n(n-1)}{2}$$

$$\langle k \rangle = \frac{1}{n} \sum_i k_i = \frac{2\langle m \rangle}{n} = p(n-1) \approx pn$$

$$\rho = \frac{\langle m \rangle}{n(n-1)/2} = p$$

Graph  $G\{E, V\}$ , nodes  $n = |V|$ , edges  $m = |E|$  Erdos and Renyi, 1959. Random graph models

- ▶  $G_{n,m}$ , a randomly selected graph from the set of  $G_N^m$  graphs;  $N = \frac{n(n-1)}{2}$ , with  $n$  nodes and  $m$  edges
- ▶ Random graph = ensemble of subgraphs
- ▶  $G_{n,p}$  each pair out of  $N = \frac{n(n-1)}{2}$  pairs of nodes is connected with probability  $p$ ,  $m$  - random number
- ▶ What properties should a random graph have?
- ▶ Number of edges ( $m$ ) and of degrees ( $k$ ), as well as density ( $\rho$ )

$$\langle m \rangle = pN = p \frac{n(n-1)}{2}$$

$$\langle k \rangle = \frac{1}{n} \sum_i k_i = \frac{2\langle m \rangle}{n} = p(n-1) \approx pn$$

$$\rho = \frac{\langle m \rangle}{n(n-1)/2} = p$$

- ▶ Probability that  $i$ -th node has a degree is connected to  $k$  other nodes is  $p^k(1 - p)^{n-1-k}$
- ▶ Numbers of way to choose  $k$  out of  $n$  nodes is  $\frac{n-1}{k}$
- ▶ What is the probability that node  $i$  is connected to  $k$  nodes?

$$p_k \frac{n-1}{k} p^k (1-p)^{n-1-k}$$

(binominal distribution, i.e a Bernoulli distribution)



- ▶ Probability that  $i$ -th node has a degree is connected to  $k$  other nodes is  $p^k(1 - p)^{n-1-k}$
- ▶ Numbers of way to choose  $k$  out of  $n$  nodes is  $\frac{n-1}{k}$
- ▶ What is the probability that node  $i$  is connected to  $k$  nodes?

$$p_k \frac{n-1}{k} p^k (1-p)^{n-1-k}$$

(binominal distribution, i.e a Bernoulli distribution)

- Probability that i-th node has a degree  $k_i = k$

$$P(k_i = k) = P(k) = C_{n-1}^k p^k (1-p)^{n-1-k}$$

(Bernoulli distribution)

$p^k$  probability that connects to  $k$  nodes (has  $k$ -edges)

$(1-p)^{n-1-k}$  probability that does not connect to any other node

$C_{n-1}^k$  number of ways to select  $k$  nodes out of all to connect

- Interesting property of  $\ln[(1-p)^{n-1-k}]$ :

$$\ln[(1-p)^{n-1-k}] = n-1-k \ln[(1-p)]$$

Recall that  $p = \frac{c}{n-1}$  and hence  $\ln[(1-p)] = \ln[1 - \frac{c}{n-1}]$

Use the Taylor's approximation:

$$\ln[(1-p)^{n-1-k}] \approx -(n-1-k) \frac{c}{n-1}$$

- Probability that i-th node has a degree  $k_i = k$

$$P(k_i = k) = P(k) = C_{n-1}^k p^k (1-p)^{n-1-k}$$

(Bernoulli distribution)

$p^k$  probability that connects to  $k$  nodes (has  $k$ -edges)

$(1-p)^{n-1-k}$  probability that does not connect to any other node

$C_{n-1}^k$  number of ways to select  $k$  nodes out of all to connect

- Interesting property of  $\ln[(1-p)^{n-1-k}]$ :

$$\ln[(1-p)^{n-1-k}] = n-1-k \ln[(1-p)]$$

Recall that  $p = \frac{c}{n-1}$  and hence  $\ln[(1-p)] = \ln[1 - \frac{c}{n-1}]$

Use the Taylor's approximation:

$$\ln[(1-p)^{n-1-k}] \approx -(n-1-k) \frac{c}{n-1}$$

- Probability that  $i$ -th node has a degree  $k_i = k$

$$P(k_i = k) = P(k) = C_{n-1}^k p^k (1-p)^{n-1-k}$$

(Bernoulli distribution)

$p^k$  probability that connects to  $k$  nodes (has  $k$ -edges)

$(1-p)^{n-1-k}$  probability that does not connect to any other node

$C_{n-1}^k$  number of ways to select  $k$  nodes out of all to connect

- Interesting property of  $\ln[(1-p)^{n-1-k}]$ :

$$\ln[(1-p)^{n-1-k}] = n-1 - k \ln[(1-p)]$$

Recall that  $p = \frac{c}{n-1}$  and hence  $\ln[(1-p)] = \ln[1 - \frac{c}{n-1}]$

Use the Taylor's approximation:

$$\ln[(1-p)^{n-1-k}] \approx -(n-1-k) \frac{c}{n-1}$$

How will

$$\ln[(1-p)^{n-1-k}] \approx -(n-1-k)\frac{c}{n-1}$$

behave if  $n \rightarrow \infty$ ?

- ▶ No general rule!
- ▶ But if we fix  $k$ , the exact equality follows

$$\ln[(1-p)^{n-1-k}] = -(n-1-k)\frac{c}{n-1} = -c$$

Now we have a new assumption ( $n \rightarrow \infty$ ?). How will it affect  $C_{n-1}^k$ ?

$$\text{▶ } C_{n-1}^k = \frac{(n-1)!}{(n-1-k)!k!} \approx \{n \rightarrow \infty\} \approx \frac{(n-1)^k}{k!}$$

How will

$$\ln[(1-p)^{n-1-k}] \approx -(n-1-k)\frac{c}{n-1}$$

behave if  $n \rightarrow \infty$ ?

- ▶ No general rule!
- ▶ But if we fix  $k$ , the exact equality follows

$$\ln[(1-p)^{n-1-k}] = -(n-1-k)\frac{c}{n-1} = -c$$

Now we have a new assumption ( $n \rightarrow \infty$ ?). How will it affect  $C_{n-1}^k$ ?

$$\text{▶ } C_{n-1}^k = \frac{(n-1)!}{(n-1-k)!k!} \approx \{n \rightarrow \infty\} \approx \frac{(n-1)^k}{k!}$$

How will

$$\ln[(1-p)^{n-1-k}] \approx -(n-1-k)\frac{c}{n-1}$$

behave if  $n \rightarrow \infty$ ?

- ▶ No general rule!
- ▶ But if we fix  $k$ , the exact equality follows

$$\ln[(1-p)^{n-1-k}] = -(n-1-k)\frac{c}{n-1} = -c$$

Now we have a new assumption ( $n \rightarrow \infty$ ?). How will it affect  $C_{n-1}^k$ ?

$$\text{▶ } C_{n-1}^k = \frac{(n-1)!}{(n-1-k)!k!} \approx \{n \rightarrow \infty\} \approx \frac{(n-1)^k}{k!}$$

- Probability that  $i$ -th node has a degree  $k_i = k$

$$P(k_i = k) = P(k) = C_{n-1}^k p^k (1-p)^{n-1-k}$$

(Bernoulli distribution)

$p^k$  probability that connects to  $k$  nodes (has  $k$ -edges)

$(1-p)^{n-1-k}$  probability that does not connect to any other node

$C_{n-1}^k$  number of ways to select  $k$  nodes out of all to connect

- Limiting case of Bernoulli distribution, when  $n \rightarrow \infty$  at fixed  $\langle k \rangle = pn = \lambda$

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!}$$

(Poisson distribution)



- Probability that  $i$ -th node has a degree  $k_i = k$

$$P(k_i = k) = P(k) = C_{n-1}^k p^k (1-p)^{n-1-k}$$

(Bernoulli distribution)

$p^k$  probability that connects to  $k$  nodes (has  $k$ -edges)


$(1-p)^{n-1-k}$  probability that does not connect to any other node

$C_{n-1}^k$  number of ways to select  $k$  nodes out of all to connect

- Limiting case of Bernoulli distribution, when  $n \rightarrow \infty$  at fixed  $\langle k \rangle = pn = \lambda$

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!}$$

(Poisson distribution)



$$P(k_i = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \lambda = pn$$

- For  $n \rightarrow \infty$  at fixed  $\langle k \rangle = pn = \lambda$ , we have the following Poisson distribution:

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!}$$

- Useful for the clustering coefficient:

$$C = \frac{\text{N of closed two-edge paths}}{\text{N of all two-edge paths}}$$

- Recall that  $p = \frac{c}{n-1}$ , which leads to

$$C = \frac{c}{n-1}$$

- Important: when  $n \rightarrow \infty$  we have  $C \rightarrow 0$  (unlike in many real networks)

- For  $n \rightarrow \infty$  at fixed  $\langle k \rangle = pn = \lambda$ , we have the following Poisson distribution:

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!}$$

- Useful for the clustering coefficient:

$$C = \frac{\text{N of closed two-edge paths}}{\text{N of all two-edge paths}}$$

- Recall that  $p = \frac{c}{n-1}$ , which leads to

$$C = \frac{c}{n-1}$$

- Important: when  $n \rightarrow \infty$  we have  $C \rightarrow 0$  (unlike in many real networks)

Consider  $G_{n,p}$  as a function of  $p$ :

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!},$$

where  $\langle k \rangle = pn = \lambda$

- ▶  $p = 0$ , empty graph (means no edges and you have  $n$  separate components)
  - ▶ Each component has cardinality of 1.
- ▶  $p = 1$ , complete (full) graph (means nodes are interconnected and you have one giant component)
  - ▶ The giant component has cardinality of  $n$ .
- ▶ There exist critical  $p_c$ , structural changes from  $p < p_c$  to  $p > p_c$
- ▶ Gigantic connected component appears at  $p > p_c$

Consider  $G_{n,p}$  as a function of  $p$ :

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!},$$

where  $\langle k \rangle = pn = \lambda$

- ▶  $p = 0$ , empty graph (means no edges and you have  $n$  separate components)
  - ▶ Each component has cardinality of 1.
- ▶  $p = 1$ , complete (full) graph (means nodes are interconnected and you have one giant component)
  - ▶ The giant component has cardinality of  $n$ .
- ▶ There exist critical  $p_c$ , structural changes from  $p < p_c$  to  $p > p_c$
- ▶ Gigantic connected component appears at  $p > p_c$

Consider  $G_{n,p}$  as a function of  $p$ :

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!},$$

where  $\langle k \rangle = pn = \lambda$

- ▶  $p = 0$ , empty graph (means no edges and you have  $n$  separate components)
  - ▶ Each component has cardinality of 1.
- ▶  $p = 1$ , complete (full) graph (means nodes are interconnected and you have one giant component)
  - ▶ The giant component has cardinality of  $n$ .
- ▶ There exist critical  $p_c$ , structural changes from  $p < p_c$  to  $p > p_c$
- ▶ Gigantic connected component appears at  $p > p_c$

Consider  $G_{n,p}$  as a function of  $p$ :

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!},$$

where  $\langle k \rangle = pn = \lambda$

- ▶  $p = 0$ , empty graph (means no edges and you have  $n$  separate components)
  - ▶ Each component has cardinality of 1.
- ▶  $p = 1$ , complete (full) graph (means nodes are interconnected and you have one giant component)
  - ▶ The giant component has cardinality of  $n$ .
- ▶ There exist critical  $p_c$ , structural changes from  $p < p_c$  to  $p > p_c$
- ▶ Gigantic connected component appears at  $p > p_c$



Consider  $G_{n,p}$  as a function of  $p$ :

$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!},$$

where  $\langle k \rangle = pn = \lambda$

- ▶  $p = 0$ , empty graph (means no edges and you have  $n$  separate components)
  - ▶ Each component has cardinality of 1.
- ▶  $p = 1$ , complete (full) graph (means nodes are interconnected and you have one giant component)
  - ▶ The giant component has cardinality of  $n$ .
- ▶ There exist critical  $p_c$ , structural changes from  $p < p_c$  to  $p > p_c$ 
  - ▶ Gigantic connected component appears at  $p > p_c$

Consider  $G_{n,p}$  as a function of  $p$ :


$$P(k) = \frac{\langle k \rangle^k e^{-\langle k \rangle}}{k!} = \frac{\lambda^k e^{-\lambda}}{k!},$$

where  $\langle k \rangle = pn = \lambda$


- ▶  $p = 0$ , empty graph (means no edges and you have  $n$  separate components)
  - ▶ Each component has cardinality of 1.
- ▶  $p = 1$ , complete (full) graph (means nodes are interconnected and you have one giant component)
  - ▶ The giant component has cardinality of  $n$ .
- ▶ There exist critical  $p_c$ , structural changes from  $p < p_c$  to  $p > p_c$
- ▶ Gigantic connected component appears at  $p > p_c$

A more formal definition of the giant component:


- ▶ Component of the network
- ▶ This component grows in size in proportion to  $n$



$$p < p_c$$




$$p = p_c$$



$$p > p_c$$

**Any real life examples come to mind?**



$$p \gg p_c$$

Let  $u$  be a fraction of nodes that do not belong to GCC. The probability that a node does not belong to GCC

$$u = P(k=0) + P(k=1) \cdot u + P(k=2) \cdot u^2 + P(k=3) \cdot u^3 \dots =$$

$$= \sum_{k=0}^{\infty} P(k) u^k = \sum_{k=0}^{\infty} \frac{\lambda^k e^{-\lambda}}{k!} u^k = e^{-\lambda} e^{\lambda u} = e^{\lambda(u-1)}$$

(Recall that  $\langle k \rangle = pn = \lambda$ )

Let  $s$  be a fraction of nodes belonging to GCC (size of GCC)

$$s = 1 - u$$

$$1 - s = u = e^{\lambda(u-1)} = e^{-\lambda s}$$

when  $\lambda \rightarrow \infty, s \rightarrow 1$

when  $\lambda \rightarrow 0, s \rightarrow 0$

Let  $u$  be a fraction of nodes that do not belong to GCC. The probability that a node does not belong to GCC

$$u = P(k=0) + P(k=1) \cdot u + P(k=2) \cdot u^2 + P(k=3) \cdot u^3 \dots =$$

$$= \sum_{k=0}^{\infty} P(k) u^k = \sum_{k=0}^{\infty} \frac{\lambda^k e^{-\lambda}}{k!} u^k = e^{-\lambda} e^{\lambda u} = e^{\lambda(u-1)}$$

(Recall that  $\langle k \rangle = pn = \lambda$ )

Let  $s$  be a fraction of nodes belonging to GCC (size of GCC)

$$s = 1 - u$$

$$1 - s = u = e^{\lambda(u-1)} = e^{-\lambda s}$$

when  $\lambda \rightarrow \infty, s \rightarrow 1$

when  $\lambda \rightarrow 0, s \rightarrow 0$

Let  $u$  be a fraction of nodes that do not belong to GCC. The probability that a node does not belong to GCC

$$u = P(k=0) + P(k=1) \cdot u + P(k=2) \cdot u^2 + P(k=3) \cdot u^3 \dots =$$

$$= \sum_{k=0}^{\infty} P(k) u^k = \sum_{k=0}^{\infty} \frac{\lambda^k e^{-\lambda}}{k!} u^k = e^{-\lambda} e^{\lambda u} = e^{\lambda(u-1)}$$

(Recall that  $\langle k \rangle = pn = \lambda$ )

Let  $s$  be a fraction of nodes belonging to GCC (size of GCC)

$$s = 1 - u$$

$$1 - s = u = e^{\lambda(u-1)} = e^{-\lambda s}$$

when  $\lambda \rightarrow \infty, s \rightarrow 1$

when  $\lambda \rightarrow 0, s \rightarrow 0$



Let  $u$  be a fraction of nodes that do not belong to GCC. The probability that a node does not belong to GCC

$$u = P(k=0) + P(k=1) \cdot u + P(k=2) \cdot u^2 + P(k=3) \cdot u^3 \dots =$$

$$= \sum_{k=0}^{\infty} P(k) u^k = \sum_{k=0}^{\infty} \frac{\lambda^k e^{-\lambda}}{k!} u^k = e^{-\lambda} e^{\lambda u} = e^{\lambda(u-1)}$$

(Recall that  $\langle k \rangle = pn = \lambda$ )

Let  $s$  be a fraction of nodes belonging to GCC (size of GCC)

$$s = 1 - u$$

$$1 - s = u = e^{\lambda(u-1)} = e^{-\lambda s}$$

when  $\lambda \rightarrow \infty, s \rightarrow 1$

when  $\lambda \rightarrow 0, s \rightarrow 0$

Let  $u$  be a fraction of nodes that do not belong to GCC. The probability that a node does not belong to GCC

$$u = P(k=0) + P(k=1) \cdot u + P(k=2) \cdot u^2 + P(k=3) \cdot u^3 \dots =$$

$$= \sum_{k=0}^{\infty} P(k) u^k = \sum_{k=0}^{\infty} \frac{\lambda^k e^{-\lambda}}{k!} u^k = e^{-\lambda} e^{\lambda u} = e^{\lambda(u-1)}$$

(Recall that  $\langle k \rangle = pn = \lambda$ )

Let  $s$  be a fraction of nodes belonging to GCC (size of GCC)


$$s = 1 - u$$

$$1 - s = u = e^{\lambda(u-1)} = e^{-\lambda s}$$

when  $\lambda \rightarrow \infty, s \rightarrow 1$

when  $\lambda \rightarrow 0, s \rightarrow 0$

$$s = 1 - e^{-\lambda s}$$



non-zero solution exists when (at  $s = 0$ ):


$$\lambda e^{-\lambda s} > 1$$

critical value:

$$\lambda_c = 1$$

$$\lambda_c = \langle k \rangle = p_c n = 1, p_c = \frac{1}{n}$$

$$s = 1 - e^{-\lambda s}$$



non-zero solution exists when (at  $s = 0$ ):


$$\lambda e^{-\lambda s} > 1$$

critical value:

$$\lambda_c = 1$$

$$\lambda_c = \langle k \rangle = p_c n = 1, p_c = \frac{1}{n}$$

$$s = 1 - e^{-\lambda s}$$





non-zero solution exists when (at  $s = 0$ ):

$$\lambda e^{-\lambda s} > 1$$

critical value:

$$\lambda_c = 1$$

$$\lambda_c = \langle k \rangle = p_c n = 1, p_c = \frac{1}{n}$$




$$\langle k \rangle = pn$$

Graph  $G(n, p)$ , for  $n \rightarrow \infty$ , critical value  $p_c = \frac{1}{n}$

- ▶ when  $p < p_c$  , ( $\langle k \rangle < 1$ ) there is no components with more than  $O(\ln n)$  nodes, largest component is a tree
- ▶ when  $p = p_c$  , ( $\langle k \rangle = 1$ ) the largest component has  $O(n^{2/3})$  nodes
- ▶ when  $p > p_c$  , ( $\langle k \rangle > 1$ ) gigantic component has all  $O(n)$  nodes

Critical value:  $\langle k \rangle = p_c n = 1$ - on average one neighbor for a node


Another interpretation: *core vs periphery*

Clauset, 2014



Graph  $G(n, p)$

Threshold probabilities when different subgraphs of  $k$ -nodes and  $l$ -edges appear in a random graph  $p_c \sim n^{-k/l}$



When  $p > p_c$  :

- ▶  $p_c \sim n^{-k/(k-1)}$ , having a tree with  $k$  nodes
- ▶  $p_c \sim n^{-1}$ , having a cycle with  $k$  nodes
- ▶  $p_c \sim n^{-2/(k-1)}$ , complete subgraph with  $k$  nodes

Barabasi, 2002


- Clustering coefficient (probability that two neighbors link to each other):

$$C_{ik} = \frac{\text{\#of links between NN}}{\text{\#max number of links NN}} = \frac{pk(k-1)/2}{k(k-1)/2} = p$$

$$C = p = \frac{\langle k \rangle}{n}$$

- when  $n \rightarrow \infty$ ,  $C \rightarrow 0$

- $G(n, p)$  is locally tree-like (GCC) (no loops; low clustering coefficient)



- on average, the number of nodes  $d$  steps away from a node

$$n = 1 + \langle k \rangle + \langle k \rangle^2 + \dots + \langle k \rangle^D = \frac{\langle k \rangle^{D+1} - 1}{\langle k \rangle - 1} \approx \langle k \rangle^D$$

- in GCC, around  $p_c$ ,  $\langle k \rangle^D \sim n$ ,

$$D \sim \frac{\ln n}{\ln \langle k \rangle}$$

$G(n,p)$  model has many problems:

- ▶ Node degree distribution - Poisson, not a power law distribution:

$$P(k) = \frac{\lambda^k e^{-\lambda}}{k!}, \lambda = pn$$

- ▶ Clustering coefficient - small (no transitivity, unlike in many real networks ):

$$C = p$$

- ▶ Graph diameter - small (no correlation among connected nodes):

$$D \sim \ln n$$

- ▶ Key idea: replace a degree distribution with a degree sequence
- ▶ In other words: specify an exact degree of each node, rather than a degree distribution
- ▶ Random graph with  $n$  nodes with a given degree sequence:  $D = \{k_1, k_2, k_3 \cdots k_n\}$ , and  $m = 1/2 \sum_i k_i$  edges is fixed
- ▶ Configuration model is somewhere between  $G(n, m)$  and  $G(n, p)$ , but more accurate than the latter

- ▶ Key idea: replace a degree distribution with a degree sequence
- ▶ In other words: specify an exact degree of each node, rather than a degree distribution
- ▶ Random graph with  $n$  nodes with a given degree sequence:  $D = \{k_1, k_2, k_3 \cdots k_n\}$ , and  $m = 1/2 \sum_i k_i$  edges is fixed
- ▶ Configuration model is somewhere between  $G(n, m)$  and  $G(n, p)$ , but more accurate than the latter

- ▶ Key idea: replace a degree distribution with a degree sequence
- ▶ In other words: specify an exact degree of each node, rather than a degree distribution
- ▶ Random graph with  $n$  nodes with a given degree sequence:  $D = \{k_1, k_2, k_3 \cdots k_n\}$ , and  $m = 1/2 \sum_i k_i$  edges
- ▶ Construct subgraphs by randomly matching two stubs and connecting them by an edge.



- ▶ Can contain self loops and multiple edges
- ▶ Probability that two nodes  $i$  and  $j$  are connected

$$p_{ij} = \frac{k_i k_j}{2m - 1}$$

- ▶ Will be a simple graph for special "graphical degree sequence"

- Probability that two nodes  $i$  and  $j$  are connected


$$p_{ij} = \frac{k_i k_j}{2m - 1}$$

- When we have a large number of edges, i.e.  $m \rightarrow \infty$ :


$$p_{ij} = \frac{k_i k_j}{2m}$$



Can be used as a "null model" for comparative network analysis



karate club



configuration model

Clauset, 2014

- ▶ On random graphs I, P. Erdos and A. Renyi, Publicationes Mathematicae 6, 290-297 (1959).
- ▶ On the evolution of random graphs, P. Erdos and A. Renyi, Publication of the Mathematical Institute of the Hungarian Academy of Sciences, 17-61 (1960)