

A multi-sorted construction in finitely decidable varieties

Matthew Smedberg

Vanderbilt University
Department of Mathematics

5 October 2013

Decidability and Finite Decidability

Finitely decidable
varieties

M. Smedberg

Finitely decidable
modular varieties

Strongly Abelian
varieties

Unification

Further questions

We say that a variety \mathcal{V} is said to be *decidable* if $\text{Th}(\mathcal{V})$ is a computable set of sentences, and likewise *finitely decidable* if $\text{Th}_{\text{fin}}(\mathcal{V}) = \text{Th}(\mathcal{V}_{\text{fin}})$ is a computable set.

Locally finite *decidable* varieties were studied by McKenzie and Valeriote in the 80s; they showed that every such variety is a varietal product of

- ▶ a strongly abelian subvariety,
- ▶ an affine subvariety, and
- ▶ a discriminator subvariety.

Since every finitely generated discriminator variety is decidable, and an explicit characterization exists for decidability in strongly abelian varieties (more on this in a moment), it only remained to determine which affine varieties are decidable.

Open Problem 1

For which finite rings \mathbf{R} is the variety of all \mathbf{R} -modules decidable? Finitely decidable?

No decomposition for FD varieties

Finitely decidable
varieties

M. Smedberg

In 1997, Idziak characterized locally finite FD varieties having modular congruence lattices (which, by TCT and a nice result of Valeriote and Willard, are exactly those varieties omitting type 1), up to Problem 1.

Finitely decidable
modular varieties

Strongly Abelian
varieties

Unification

Further questions

Idziak's proof essentially gives a recipe for taking a FD variety \mathcal{V} admitting only type 3, another FD variety \mathcal{W} of modules, and producing a FD supervariety $\mathcal{V} \star \mathcal{W}$ such that

- ▶ if $\mathbf{A} \in \mathcal{V} \star \mathcal{W}$ omits type 2, then $\mathbf{A} \in \mathcal{V}$;
- ▶ if $\mathbf{A}/\text{Rad}(\mathbf{A})$ (which must lie in \mathcal{V}) is subdirectly irreducible, every $\text{Rad}(\mathbf{A})$ -block is a module from \mathcal{W} ;
- ▶ there exist algebras in $\mathcal{V} \star \mathcal{W}$ whose congruence lattice is the ordered sum of a nontrivial solvable interval and a nontrivial chain of boolean type covers.

Going in the reverse direction, Idziak gave sufficient conditions that guarantee, for modular variety \mathcal{V} and $\mathbf{A} \in \mathcal{V}$, that the poset of meet-irreducible $\tau \in \text{Con}(\mathbf{A})$ with boolean upper cover forms a tree \mathbb{T} with root \top_A ; that each $\tau/[\tau, \tau]$ -block is a module over a ring uniformly definable from the variety; and that \mathbf{A} could be isomorphically recovered from the family

$$\langle \mathbf{A}/[\tau, \tau] : \tau \in \mathbb{T} \rangle$$

in a uniform way. It follows that deciding the theory of \mathcal{V} is no harder than deciding \mathcal{V}_3 and the theory of modules over the indicated ring.

Question

Can we do something similar in the case where a variety \mathcal{V} omits type 2: that is, find a variety \mathcal{W} of strongly abelian algebras such that we can associate blocks of strongly abelian congruences in \mathcal{V} to algebras in \mathcal{W} well enough to determine the structure of algebras in \mathcal{V} ?

Back to strongly abelian algebras

Finitely decidable
varieties

M. Smedberg

Finitely decidable
modular varieties

Strongly Abelian
varieties

Unification

Further questions

Lemma

Let \mathbf{A} be a finite, strongly abelian algebra, and let k be the largest number such that \mathbf{A} has an idempotent term operation $t(x_1, \dots, x_k)$ that depends on all its variables. Then we can find a term $d(x_1, \dots, x_k)$ and a decomposition

$$A = A_1 \times \cdots \times A_k$$

such that

$$d(a_1, \dots, a_k) = d \begin{pmatrix} a_1^1 & a_2^1 & \cdots & a_k^1 \\ a_1^2 & a_2^2 & \cdots & a_k^2 \\ \vdots & \vdots & \ddots & \vdots \\ a_1^k & a_2^k & \cdots & a_k^k \end{pmatrix} = \begin{pmatrix} a_1^1 \\ a_2^2 \\ \vdots \\ a_k^k \end{pmatrix}$$

Suppose $d(x_1, \dots, x_k)$ acts as a decomposition term throughout the strongly abelian variety \mathcal{V} . Valeriote and McKenzie define a k -sorted companion variety $\mathcal{V}[d]$ by turning each n -ary fundamental operation of \mathcal{V} into k operations of arity $k \cdot n$.

Theorem 1

- ▶ \mathcal{V} is essentially k -ary iff $\mathcal{V}[d]$ is essentially unary.
- ▶ If $\mathcal{V}[d]$ is not essentially unary, then $\mathcal{V}[d]$ and \mathcal{V} are both undecidable and finitely undecidable.

Question

Can we do something similar in the case where we cannot decompose an algebra into a product of strongly abelian and totally nonabelian factors?

Unifying the constructions

Finitely decidable
varieties

M. Smedberg

Finitely decidable
modular varieties

Strongly Abelian
varieties

Unification

Further questions

Let \mathbf{A} be a finite algebra in a variety \mathcal{V} which satisfies all the known necessary conditions for finite decidability, and suppose $\text{typ}\{\mathcal{V}\} = \{1, 3\}$. Then the meet-irreducible congruences $\tau \in \text{Con}(\mathbf{A})$ with boolean upper cover are arranged in a tree \mathbb{T} with $\top_{\mathbf{A}}$ at the root; for each $\tau \in \mathbb{T}$, the least $\zeta \in \text{Con}(\mathbf{A})$ such that τ is solvable over ζ is in fact $[\tau, \tau]$; and \mathbf{A} is a subdirect product of the $\mathbf{A}/[\tau, \tau]$.

Theorem 2 (S.)

Replacing \mathbf{A} by one of the $\mathbf{A}/[\tau, \tau]$, we know that $\text{Rad}(\mathbf{A})$ is a strongly abelian congruence, and meet-irreducible with boolean monolith. Say $\text{Rad}(\mathbf{A})$ has m blocks. For each block B_i , let \mathbf{A}_i be the algebra induced on B_i by all terms (not polynomials) which respect it. \mathbf{A}_i is a strongly abelian algebra, with k_i -ary decomposition term d_i .

Then if any term of \mathbf{A}_i depends on more than k_i variables, then \mathcal{V} is hereditarily finitely undecidable.

Proof by construction

Finitely decidable
varieties

M. Smedberg

To prove it, we define a multi-sorted first-order language L^b .
The sorts are

Finitely decidable
modular varieties

Strongly Abelian
varieties

Unification

Further questions

$$\{\langle i, j \rangle : 1 \leq i \leq m, 1 \leq j \leq k_i\}$$

and for each basic operation $f(x_1, \dots, x_n)$ of **A** and any
blocks

$$f : B_{i_1} \times \dots \times B_{i_n} \rightarrow B_{i_0}$$

we include k_{i_0} basic operations

$$f_{i_1, \dots, i_n}^j \left(\begin{array}{cccc} \langle i_1, 1 \rangle & \langle i_2, 1 \rangle & \cdots & \langle i_n, 1 \rangle \\ \langle i_1, 2 \rangle & \langle i_2, 2 \rangle & \cdots & \langle i_n, 2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle i_1, k_{i_1} \rangle & \langle i_2, k_{i_2} \rangle & \cdots & \langle i_n, k_{i_n} \rangle \end{array} \right) \rightarrow \langle i_0, j \rangle$$

Then if d_i decomposes B_i into $B_{i,1} \times B_{i,2} \times \cdots \times B_{i,k_i}$, we define an L^b -structure \mathbf{A}^b , interpreting sort $\langle i, j \rangle$ by $B_{i,j}$, and evaluating the functions in the only reasonable way.

Lemma

\mathbf{A}^b is strongly abelian.

(Recall that \mathbf{A} was not even abelian!)

Lemma

Call $\mathbf{C} \in \mathcal{V}$ flattable if this construction sends it to a sorted algebra $\mathbf{C}^b \in \mathcal{V}^b = \text{HSP}(\mathbf{A}^b)$. Let $\mathbf{C}, \mathbf{C}_1, \mathbf{C}_2$ be flattable.

Then

- ▶ \mathbf{C}^b is strongly abelian, and $\mathbf{C}/\text{Rad}(\mathbf{C})$ is canonically isomorphic to $\mathbf{A}/\text{Rad}(\mathbf{A})$. Call the canonical projection $\pi_{\mathbf{C}}$.
- ▶ Subalgebras of \mathbf{C}^b are precisely the flat images of subalgebras of \mathbf{C} which have nonempty intersection with every $\text{Rad}(\mathbf{C})$ -block.
- ▶ Homomorphic images of \mathbf{C}^b are precisely the flat images of homomorphic images of \mathbf{C} by solvable congruences.
- ▶ The product $\mathbf{C}_1^b \times \mathbf{C}_2^b$ is the flat image of

$$\left\{ \begin{bmatrix} x \\ y \end{bmatrix} : \pi_{\mathbf{C}_1}(x) = \pi_{\mathbf{C}_2}(y) \right\} \leq \mathbf{C}_1 \times \mathbf{C}_2$$

The variety generated by \mathbf{A}^b , in other words, interprets into \mathcal{V} . Notice that the trivial algebra in this variety is obtained by flatting $\mathbf{A}/\text{Rad}(\mathbf{A})$.

All that remains is now to show that Valeriote's proof of Theorem 1 goes through.

It does.

“When proving something is undecidable, use local structure. When proving something is decidable, you have to use global structure.”

– Pawel Idziak

Open Problem 2

Are the algebras $\mathbf{A}/[\tau, \tau]$ sufficiently regular to be manageable, while simultaneously encoding enough information to recover \mathbf{A} in some uniform, first-order way?

Open Problem 3

What about varieties that admit both type 1 and 2? Even in the abelian case, we're pretty much stumped so far.