

Residually Finite Varieties

Keith A. Kearnes
University of Colorado

Residual character

The *residual character* of a variety \mathcal{V} is the least $\chi_{\mathcal{V}}$ such that every subdirectly irreducible algebra in \mathcal{V} has cardinality $< \chi_{\mathcal{V}}$, if exists. Else $\chi_{\mathcal{V}} = \infty$.

Residual character

The *residual character* of a variety \mathcal{V} is the least $\chi_{\mathcal{V}}$ such that every subdirectly irreducible algebra in \mathcal{V} has cardinality $< \chi_{\mathcal{V}}$, if exists. Else $\chi_{\mathcal{V}} = \infty$. \mathcal{V} is *residually finite* if $\chi_{\mathcal{V}} \leq \omega$.

Residual character

The *residual character* of a variety \mathcal{V} is the least $\chi_{\mathcal{V}}$ such that every subdirectly irreducible algebra in \mathcal{V} has cardinality $< \chi_{\mathcal{V}}$, if exists. Else $\chi_{\mathcal{V}} = \infty$. \mathcal{V} is *residually finite* if $\chi_{\mathcal{V}} \leq \omega$.

Theorem (McKenzie-Shelah, 1974)

If \mathcal{V} has finite type, then $\chi_{\mathcal{V}} \in \{3, 4, \dots, \omega, \omega^+, (2^\omega)^+, \infty\}$.

Residual character

The *residual character* of a variety \mathcal{V} is the least $\chi_{\mathcal{V}}$ such that every subdirectly irreducible algebra in \mathcal{V} has cardinality $< \chi_{\mathcal{V}}$, if exists. Else $\chi_{\mathcal{V}} = \infty$. \mathcal{V} is *residually finite* if $\chi_{\mathcal{V}} \leq \omega$.

Theorem (McKenzie-Shelah, 1974)

If \mathcal{V} has finite type, then $\chi_{\mathcal{V}} \in \{3, 4, \dots, \omega, \omega^+, (2^\omega)^+, \infty\}$.

Residual character

The *residual character* of a variety \mathcal{V} is the least $\chi_{\mathcal{V}}$ such that every subdirectly irreducible algebra in \mathcal{V} has cardinality $< \chi_{\mathcal{V}}$, if exists. Else $\chi_{\mathcal{V}} = \infty$. \mathcal{V} is *residually finite* if $\chi_{\mathcal{V}} \leq \omega$.

Theorem (McKenzie-Shelah, 1974)

If \mathcal{V} has finite type, then $\chi_{\mathcal{V}} \in \{3, 4, \dots, \omega, \omega^+, (2^\omega)^+, \infty\}$.

It seems hard to realize $\chi_{\mathcal{V}} = \omega$ in finite type.

Residual character

The *residual character* of a variety \mathcal{V} is the least $\chi_{\mathcal{V}}$ such that every subdirectly irreducible algebra in \mathcal{V} has cardinality $< \chi_{\mathcal{V}}$, if exists. Else $\chi_{\mathcal{V}} = \infty$. \mathcal{V} is *residually finite* if $\chi_{\mathcal{V}} \leq \omega$.

Theorem (McKenzie-Shelah, 1974)

If \mathcal{V} has finite type, then $\chi_{\mathcal{V}} \in \{3, 4, \dots, \omega, \omega^+, (2^\omega)^+, \infty\}$.

It seems hard to realize $\chi_{\mathcal{V}} = \omega$ in finite type. A variety with this property has infinitely many finite SI's, but no infinite ones.

Residual character

The *residual character* of a variety \mathcal{V} is the least $\chi_{\mathcal{V}}$ such that every subdirectly irreducible algebra in \mathcal{V} has cardinality $< \chi_{\mathcal{V}}$, if exists. Else $\chi_{\mathcal{V}} = \infty$. \mathcal{V} is *residually finite* if $\chi_{\mathcal{V}} \leq \omega$.

Theorem (McKenzie-Shelah, 1974)

If \mathcal{V} has finite type, then $\chi_{\mathcal{V}} \in \{3, 4, \dots, \omega, \omega^+, (2^\omega)^+, \infty\}$.

It seems hard to realize $\chi_{\mathcal{V}} = \omega$ in finite type. A variety with this property has infinitely many finite SI's, but no infinite ones. The McKenzie-Shelah example is the variety of \mathbb{Z} -sets, presented with two unary operations satisfying $f(g(x)) = g(f(x)) = x$.

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) *a variety of groups (Olshanskii, 1969),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) a variety of groups (Olshanskii, 1969),*
- (2) a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) *a variety of groups (Olshanskii, 1969),*
- (2) *a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) *a variety of rings (McKenzie, 1982),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) a variety of groups (Olshanskii, 1969),*
- (2) a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) a variety of rings (McKenzie, 1982),*
- (4) an affine complete variety (Kaarli-Pixley, 1987),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) *a variety of groups (Olshanskii, 1969),*
- (2) *a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) *a variety of rings (McKenzie, 1982),*
- (4) *an affine complete variety (Kaarli-Pixley, 1987),*
- (5) *a variety of lattices (Freese-Nation, ?),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) a variety of groups (Olshanskii, 1969),*
- (2) a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) a variety of rings (McKenzie, 1982),*
- (4) an affine complete variety (Kaarli-Pixley, 1987),*
- (5) a variety of lattices (Freese-Nation, ?),*
- (6) a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) a variety of groups (Olshanskii, 1969),*
- (2) a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) a variety of rings (McKenzie, 1982),*
- (4) an affine complete variety (Kaarli-Pixley, 1987),*
- (5) a variety of lattices (Freese-Nation, ?),*
- (6) a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*
- (7) a variety that omits nontrivial abelian congruences (Kearnes-Willard, 1999),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) a variety of groups (Olshanskii, 1969),*
- (2) a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) a variety of rings (McKenzie, 1982),*
- (4) an affine complete variety (Kaarli-Pixley, 1987),*
- (5) a variety of lattices (Freese-Nation, ?),*
- (6) a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*
- (7) a variety that omits nontrivial abelian congruences (Kearnes-Willard, 1999),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) a variety of groups (Olshanskii, 1969),*
- (2) a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) a variety of rings (McKenzie, 1982),*
- (4) an affine complete variety (Kaarli-Pixley, 1987),*
- (5) a variety of lattices (Freese-Nation, ?),*
- (6) a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*
- (7) a variety that omits nontrivial abelian congruences (Kearnes-Willard, 1999), ($\text{typ}\{\mathcal{V}\} \subseteq \{3, 4, 5\}$)*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) *a variety of groups (Olshanskii, 1969),*
- (2) *a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) *a variety of rings (McKenzie, 1982),*
- (4) *an affine complete variety (Kaarli-Pixley, 1987),*
- (5) *a variety of lattices (Freese-Nation, ?),*
- (6) *a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*
- (7) *a variety that omits nontrivial abelian congruences (Kearnes-Willard, 1999), ($\text{typ}\{\mathcal{V}\} \subseteq \{3, 4, 5\}$)*
- (8) *a variety that omits nontrivial rectangular congruences (Kearnes-Willard, ?),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) *a variety of groups (Olshanskii, 1969),*
- (2) *a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) *a variety of rings (McKenzie, 1982),*
- (4) *an affine complete variety (Kaarli-Pixley, 1987),*
- (5) *a variety of lattices (Freese-Nation, ?),*
- (6) *a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*
- (7) *a variety that omits nontrivial abelian congruences (Kearnes-Willard, 1999), ($\text{typ}\{\mathcal{V}\} \subseteq \{3, 4, 5\}$)*
- (8) *a variety that omits nontrivial rectangular congruences (Kearnes-Willard, ?),*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) *a variety of groups (Olshanskii, 1969),*
- (2) *a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) *a variety of rings (McKenzie, 1982),*
- (4) *an affine complete variety (Kaarli-Pixley, 1987),*
- (5) *a variety of lattices (Freese-Nation, ?),*
- (6) *a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*
- (7) *a variety that omits nontrivial abelian congruences (Kearnes-Willard, 1999), ($\text{typ}\{\mathcal{V}\} \subseteq \{3, 4, 5\}$)*
- (8) *a variety that omits nontrivial rectangular congruences (Kearnes-Willard, ?), ($\text{typ}\{\mathcal{V}\} \subseteq \{2, 3, 4\}$)*

Evidence that $\chi_{\mathcal{V}} = \omega$ is hard to realize

Theorems

If \mathcal{V} has finite type, and is among the following, then $\chi_{\mathcal{V}} \neq \omega$:

- (1) *a variety of groups (Olshanskii, 1969),*
- (2) *a variety of semigroups (McKenzie, 1981), (Golubov-Sapir, 1982),*
- (3) *a variety of rings (McKenzie, 1982),*
- (4) *an affine complete variety (Kaarli-Pixley, 1987),*
- (5) *a variety of lattices (Freese-Nation, ?),*
- (6) *a variety of Lie algebras over a finite field of characteristic ≥ 5 (Premet-Semenov, 1990),*
- (7) *a variety that omits nontrivial abelian congruences (Kearnes-Willard, 1999), ($\text{typ}\{\mathcal{V}\} \subseteq \{3, 4, 5\}$)*
- (8) *a variety that omits nontrivial rectangular congruences (Kearnes-Willard, ?), ($\text{typ}\{\mathcal{V}\} \subseteq \{2, 3, 4\}$)*



Conjecture

If \mathcal{V} has finite type and satisfies $\text{typ}\{\mathcal{V}\} \subseteq \{2, 3, 4, 5\}$, then $\chi_{\mathcal{V}} \neq \omega$.

How to prove $\chi_\nu \neq \omega$

SI's: $A_1,$ $A_2,$ $A_3,$ $|A| \nearrow \omega$

Con's:






...

How to prove $\chi_\nu \neq \omega$

SI's: $A_1,$ $A_2,$ $A_3,$ $|A| \nearrow \omega$

Con's:






...

$$\nu = (0 : \mu)$$

How to prove $\chi_V \neq \omega$


SI's: $A_1,$ $A_2,$ $A_3,$ $|A| \nearrow \omega$

Con's:



...

A





$$\nu = (0 : \mu)$$


How to prove $\chi_V \neq \omega$

SI's: $A_1,$ $A_2,$ $A_3,$ $|A| \nearrow \omega$

Con's:



...



$$\nu = (0 : \mu)$$

$$S = \{s_1, s_2, \dots\}$$

= a noncentralizing set

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

Key fact: If $\chi_{\mathcal{V}} \leq \omega$, \mathcal{V} has finite type, and \mathcal{V} omits nontrivial rectangular congruences, then “[$\text{Cg}(w, x), \text{Cg}(y, z)$] = 0” is definable.

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

Key fact: If $\chi_{\mathcal{V}} \leq \omega$, \mathcal{V} has finite type, and \mathcal{V} omits nontrivial rectangular congruences, then “[$\text{Cg}(w, x), \text{Cg}(y, z)$] = 0” is definable.

Hence \exists formula $\Gamma(p, q, u, v)$ such that

(1) Γ is a principal congruence formula (if $A \models \Gamma(e, f, a, b)$, then $(e, f) \in \text{Cg}(a, b)$).

(2) If $[\text{Cg}(a, b), \text{Cg}(c, d)] \neq 0$, then $\exists \{e, f\} \in A^{(2)}$ such that $A \models \Gamma(e, f, a, b)$ and $A \models \Gamma(e, f, c, d)$.

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

Key fact: If $\chi_{\mathcal{V}} \leq \omega$, \mathcal{V} has finite type, and \mathcal{V} omits nontrivial rectangular congruences, then “ $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ ” is definable.

Hence \exists formula $\Gamma(p, q, u, v)$ such that

(1) Γ is a principal congruence formula (if $A \models \Gamma(e, f, a, b)$, then $(e, f) \in \text{Cg}(a, b)$).

(2) If $[\text{Cg}(a, b), \text{Cg}(c, d)] \neq 0$, then $\exists \{e, f\} \in A^{(2)}$ such that $A \models \Gamma(e, f, a, b)$ and $A \models \Gamma(e, f, c, d)$.

Example: (Rings, $\text{Cg}(a, b) = \text{Cg}(a - b, 0)$)

$$[\text{Cg}(a, 0), \text{Cg}(c, 0)] = 0 \iff (a)(c) = (0) = (c)(a)$$

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

Key fact: If $\chi_{\mathcal{V}} \leq \omega$, \mathcal{V} has finite type, and \mathcal{V} omits nontrivial rectangular congruences, then “ $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ ” is definable.

Hence \exists formula $\Gamma(p, q, u, v)$ such that

(1) Γ is a principal congruence formula (if $A \models \Gamma(e, f, a, b)$, then $(e, f) \in \text{Cg}(a, b)$).

(2) If $[\text{Cg}(a, b), \text{Cg}(c, d)] \neq 0$, then $\exists \{e, f\} \in A^{(2)}$ such that $A \models \Gamma(e, f, a, b)$ and $A \models \Gamma(e, f, c, d)$.

Example: (Rings, $\text{Cg}(a, b) = \text{Cg}(a - b, 0)$)

$$\begin{aligned} [\text{Cg}(a, 0), \text{Cg}(c, 0)] = 0 &\Leftrightarrow (a)(c) = (0) = (c)(a) \\ &\Leftrightarrow \forall r (arc = 0 \ \& \ cra = 0) \end{aligned}$$

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

Key fact: If $\chi_{\mathcal{V}} \leq \omega$, \mathcal{V} has finite type, and \mathcal{V} omits nontrivial rectangular congruences, then “ $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ ” is definable.

Hence \exists formula $\Gamma(p, q, u, v)$ such that

(1) Γ is a principal congruence formula (if $A \models \Gamma(e, f, a, b)$, then $(e, f) \in \text{Cg}(a, b)$).

(2) If $[\text{Cg}(a, b), \text{Cg}(c, d)] \neq 0$, then $\exists \{e, f\} \in A^{(2)}$ such that $A \models \Gamma(e, f, a, b)$ and $A \models \Gamma(e, f, c, d)$.

Example: (Rings, $\text{Cg}(a, b) = \text{Cg}(a - b, 0)$)

$$\begin{aligned} [\text{Cg}(a, 0), \text{Cg}(c, 0)] = 0 &\Leftrightarrow (a)(c) = (0) = (c)(a) \\ &\Leftrightarrow \forall r (arc = 0 \ \& \ cra = 0) \end{aligned}$$

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

Key fact: If $\chi_{\mathcal{V}} \leq \omega$, \mathcal{V} has finite type, and \mathcal{V} omits nontrivial rectangular congruences, then “ $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ ” is definable.

Hence \exists formula $\Gamma(p, q, u, v)$ such that

(1) Γ is a principal congruence formula (if $A \models \Gamma(e, f, a, b)$, then $(e, f) \in \text{Cg}(a, b)$).

(2) If $[\text{Cg}(a, b), \text{Cg}(c, d)] \neq 0$, then $\exists \{e, f\} \in A^{(2)}$ such that $A \models \Gamma(e, f, a, b)$ and $A \models \Gamma(e, f, c, d)$.

Example: (Rings, $\text{Cg}(a, b) = \text{Cg}(a - b, 0)$)

$$\begin{aligned} [\text{Cg}(a, 0), \text{Cg}(c, 0)] \neq 0 &\Leftrightarrow (a)(c) \neq (0) \text{ or } (c)(a) \neq (0) \\ &\Leftrightarrow \exists r (arc \neq 0 \text{ or } cra \neq 0) \end{aligned}$$

S ‘noncentralizing’ means: if $\{s, t\} \in S^{(2)}$ and $\{a, b\} \in A^{(2)}$, then $[\text{Cg}(s, t), \text{Cg}(a, b)] \geq [\text{Cg}(s, t), \mu] > 0$.

Key fact: If $\chi_{\mathcal{V}} \leq \omega$, \mathcal{V} has finite type, and \mathcal{V} omits nontrivial rectangular congruences, then “ $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ ” is definable.

Hence \exists formula $\Gamma(p, q, u, v)$ such that

(1) Γ is a principal congruence formula (if $A \models \Gamma(e, f, a, b)$, then $(e, f) \in \text{Cg}(a, b)$).

(2) If $[\text{Cg}(a, b), \text{Cg}(c, d)] \neq 0$, then $\exists \{e, f\} \in A^{(2)}$ such that $A \models \Gamma(e, f, a, b)$ and $A \models \Gamma(e, f, c, d)$.

Example: (Rings, $\text{Cg}(a, b) = \text{Cg}(a - b, 0)$)

$$\begin{aligned} [\text{Cg}(a, 0), \text{Cg}(c, 0)] \neq 0 &\Leftrightarrow (a)(c) \neq (0) \text{ or } (c)(a) \neq (0) \\ &\Leftrightarrow \exists r (arc \neq 0 \text{ or } cra \neq 0) \end{aligned}$$

$$\Gamma(e, 0, a, 0) = \exists s (e = sa \text{ or } e = as)$$

Bound the size of noncentralizing sets in SI's

For pcf Γ' , write $S \xrightarrow{\Gamma'} \{u, v\}$ to mean $A \models \Gamma'(u, v, s, t)$ for all $\{s, t\} \in S^{(2)}$.

Bound the size of noncentralizing sets in SI's

For pcf Γ' , write $S \xrightarrow{\Gamma'} \{u, v\}$ to mean $A \models \Gamma'(u, v, s, t)$ for all $\{s, t\} \in S^{(2)}$.

Claim. \exists pcf Γ_n such that for any noncentralizing $S \subseteq A \in \mathcal{V}_{SI}$, $|S| = n$,
 $\forall \{a, b\} \in A^{(2)} \exists \{u, v\} \in A^{(2)} (\{a, b\} \xrightarrow{\Gamma_n} \{u, v\} \xleftarrow{\Gamma_n} S)$:

Bound the size of noncentralizing sets in SI's

For pcf Γ' , write $S \xrightarrow{\Gamma'} \{u, v\}$ to mean $A \models \Gamma'(u, v, s, t)$ for all $\{s, t\} \in S^{(2)}$.

Claim. \exists pcf Γ_n such that for any noncentralizing $S \subseteq A \in \mathcal{V}_{SI}$, $|S| = n$,
 $\forall \{a, b\} \in A^{(2)} \exists \{u, v\} \in A^{(2)} (\{a, b\} \xrightarrow{\Gamma_n} \{u, v\} \xleftarrow{\Gamma_n} S)$:

$$\begin{array}{ccc} & \{s_1, t_1\} & \\ & \downarrow \Gamma & \\ \{a, b\} & \xrightarrow{\Gamma} & \{u_1, v_1\} \end{array}$$

Bound the size of noncentralizing sets in SI's

For pcf Γ' , write $S \xrightarrow{\Gamma'} \{u, v\}$ to mean $A \models \Gamma'(u, v, s, t)$ for all $\{s, t\} \in S^{(2)}$.

Claim. \exists pcf Γ_n such that for any noncentralizing $S \subseteq A \in \mathcal{V}_{SI}$, $|S| = n$,
 $\forall \{a, b\} \in A^{(2)} \exists \{u, v\} \in A^{(2)} (\{a, b\} \xrightarrow{\Gamma_n} \{u, v\} \xleftarrow{\Gamma_n} S)$:

$$\begin{array}{ccccc} & \{s_1, t_1\} & & \{s_2, t_2\} & \\ & \downarrow \Gamma & & \downarrow \Gamma & \\ \{a, b\} & \xrightarrow{\Gamma} & \{u_1, v_1\} & \xrightarrow{\Gamma} & \{u_2, v_2\} \end{array}$$

Bound the size of noncentralizing sets in SI's

For pcf Γ' , write $S \xrightarrow{\Gamma'} \{u, v\}$ to mean $A \models \Gamma'(u, v, s, t)$ for all $\{s, t\} \in S^{(2)}$.

Claim. \exists pcf Γ_n such that for any noncentralizing $S \subseteq A \in \mathcal{V}_{SI}$, $|S| = n$,
 $\forall \{a, b\} \in A^{(2)} \exists \{u, v\} \in A^{(2)} (\{a, b\} \xrightarrow{\Gamma_n} \{u, v\} \xleftarrow{\Gamma_n} S)$:

$$\begin{array}{ccccccc}
 & \{s_1, t_1\} & & \{s_2, t_2\} & & & \{s_m, t_m\} \\
 & \downarrow \Gamma & & \downarrow \Gamma & & & \downarrow \Gamma \\
 \{a, b\} & \xrightarrow{\Gamma} & \{u_1, v_1\} & \xrightarrow{\Gamma} & \{u_2, v_2\} & \xrightarrow{\Gamma} & \cdots \xrightarrow{\Gamma} \{u_m, v_m\}
 \end{array}$$

Bound the size of noncentralizing sets in SI's


For pcf Γ' , write $S \xrightarrow{\Gamma'} \{u, v\}$ to mean $A \models \Gamma'(u, v, s, t)$ for all $\{s, t\} \in S^{(2)}$.

Claim. \exists pcf Γ_n such that for any noncentralizing $S \subseteq A \in \mathcal{V}_{SI}$, $|S| = n$,
 $\forall \{a, b\} \in A^{(2)} \exists \{u, v\} \in A^{(2)} (\{a, b\} \xrightarrow{\Gamma_n} \{u, v\} \xleftarrow{\Gamma_n} S)$:

$$\begin{array}{ccccccc}
 & \{s_1, t_1\} & & \{s_2, t_2\} & & & \{s_m, t_m\} \\
 & \downarrow \Gamma & & \downarrow \Gamma & & & \downarrow \Gamma \\
 \{a, b\} & \xrightarrow{\Gamma} & \{u_1, v_1\} & \xrightarrow{\Gamma} & \{u_2, v_2\} & \xrightarrow{\Gamma} & \cdots \xrightarrow{\Gamma} \{u_m, v_m\} \\
 & & & & & & \parallel \\
 & & & & & & \{u, v\}
 \end{array}$$

Unbounded noncentralizing sets yield infinite SI's

If SI's have noncentralizing sets of unbounded size, then it is consistent with $\text{Th}(\mathcal{V})$ that some SI $A \in \mathcal{V}$ contains elements related as follows.



Unbounded noncentralizing sets yield infinite SI's

If SI's have noncentralizing sets of unbounded size, then it is consistent with $\text{Th}(\mathcal{V})$ that some SI $A \in \mathcal{V}$ contains elements related as follows.

$$\begin{array}{ccc} & S_4 & \\ & \downarrow \Gamma_4 & \\ \{u_5, v_5\} & \xrightarrow{\Gamma_4} & \{u_4, v_4\} \end{array}$$

Unbounded noncentralizing sets yield infinite SI's

If SI's have noncentralizing sets of unbounded size, then it is consistent with $\text{Th}(\mathcal{V})$ that some SI $A \in \mathcal{V}$ contains elements related as follows.

$$\begin{array}{ccccc} & & S_4 & & S_3 \\ & & \downarrow \Gamma_4 & & \downarrow \Gamma_3 \\ \{u_5, v_5\} & \xrightarrow{\Gamma_4} & \{u_4, v_4\} & \xrightarrow{\Gamma_3} & \{u_3, v_3\} \end{array}$$

Unbounded noncentralizing sets yield infinite SI's


If SI's have noncentralizing sets of unbounded size, then it is consistent with $\text{Th}(\mathcal{V})$ that some SI $A \in \mathcal{V}$ contains elements related as follows.

$$\begin{array}{ccccccc} & & S_4 & & S_3 & & S_2 \\ & & \downarrow \Gamma_4 & & \downarrow \Gamma_3 & & \downarrow \Gamma_2 \\ \{u_5, v_5\} & \xrightarrow{\Gamma_4} & \{u_4, v_4\} & \xrightarrow{\Gamma_3} & \{u_3, v_3\} & \xrightarrow{\Gamma_2} & \{u_2, v_2\} \end{array}$$

$|S_i| = i$ $u_2 \neq v_2$


Unbounded noncentralizing sets yield infinite SI's

If SI's have noncentralizing sets of unbounded size, then it is consistent with $\text{Th}(\mathcal{V})$ that some SI $A \in \mathcal{V}$ contains elements related as follows.



Unbounded noncentralizing sets yield infinite SI's

If SI's have noncentralizing sets of unbounded size, then it is consistent with $\text{Th}(\mathcal{V})$ that some SI $A \in \mathcal{V}$ contains elements related as follows.




Hence \mathcal{V} has an infinite SI.

Bound on $|S| = \text{bound on } [1 : \nu]$

Hypotheses

$\text{Con}(A)$, A SI




Bound on $|S|$ = bound on $[1 : \nu]$

Hypotheses

- $\chi\nu \leq \omega$
- $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$
definable

$\text{Con}(A)$, A SI




Bound on $|S| = \text{bound on } [1 : \nu]$

Hypotheses

- $\chi\nu \leq \omega$
- $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$
definable

$\text{Con}(A)$, A SI




\vdash bounded

Bound on $|S| = \text{bound on } [1 : \nu]$

Hypotheses

- $\chi\nu \leq \omega$
- $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$
definable
- $\langle 0, \mu \rangle$ not type 1

$\text{Con}(A)$, A SI




\vdash bounded

Bound on $|S|$ = bound on $[1 : \nu]$

Hypotheses

- $\chi\nu \leq \omega$
- $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$
definable
- $\langle 0, \mu \rangle$ not type **1**

$\text{Con}(A)$, A SI




Bound on $|S| = \text{bound on } [1 : \nu]$

Hypotheses

- $\chi\nu \leq \omega$
- $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$
definable
- $\langle 0, \mu \rangle$ not type **1**
- \mathcal{V} omits type **1**

$\text{Con}(A)$, A SI




Bound on $|S|$ = bound on $[1 : \nu]$

Hypotheses

- $\chi\nu \leq \omega$
- $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$
definable
- $\langle 0, \mu \rangle$ not type 1
- \mathcal{V} omits type 1

$\text{Con}(A)$, A SI




Bound on $|S|$ = bound on $[1 : \nu]$

Hypotheses

- $\chi_{\mathcal{V}} \leq \omega$
- $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ definable
- $\langle 0, \mu \rangle$ not type **1**
- \mathcal{V} omits type **1**

$\text{Con}(A)$, A SI



If $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ is definable and \mathcal{V} omits type **1**, then proving $\chi_{\mathcal{V}} \neq \omega$ can be reduced to the case of modules.

Varieties of modules

\mathcal{V} = variety of modules over a f.g. ring R . R is *minimal* if it is infinite, but every proper quotient is finite (like \mathbb{Z}).

Varieties of modules

\mathcal{V} = variety of modules over a f.g. ring R . R is *minimal* if it is infinite, but every proper quotient is finite (like \mathbb{Z}).

Lemmas

Varieties of modules

\mathcal{V} = variety of modules over a f.g. ring R . R is *minimal* if it is infinite, but every proper quotient is finite (like \mathbb{Z}).

Lemmas

(1) $\chi_{\mathcal{V}} < \omega$ iff $|R| < \omega$.

Varieties of modules

\mathcal{V} = variety of modules over a f.g. ring R . R is *minimal* if it is infinite, but every proper quotient is finite (like \mathbb{Z}).

Lemmas

- (1) $\chi_{\mathcal{V}} < \omega$ iff $|R| < \omega$.
- (2) *Every f.g. infinite ring has a minimal quotient.*

Varieties of modules

\mathcal{V} = variety of modules over a f.g. ring R . R is *minimal* if it is infinite, but every proper quotient is finite (like \mathbb{Z}).

Lemmas

- (1) $\chi_{\mathcal{V}} < \omega$ iff $|R| < \omega$.
- (2) *Every f.g. infinite ring has a minimal quotient.*
- (3) *Every injective module over a minimal ring is faithful.*

Varieties of modules

\mathcal{V} = variety of modules over a f.g. ring R . R is *minimal* if it is infinite, but every proper quotient is finite (like \mathbb{Z}).

Lemmas

- (1) $\chi_{\mathcal{V}} < \omega$ iff $|R| < \omega$.
- (2) *Every f.g. infinite ring has a minimal quotient.*
- (3) *Every injective module over a minimal ring is faithful.*
- (4) *Every variety of modules has an SI injective.*

Varieties of modules

\mathcal{V} = variety of modules over a f.g. ring R . R is *minimal* if it is infinite, but every proper quotient is finite (like \mathbb{Z}).

Lemmas

- (1) $\chi_{\mathcal{V}} < \omega$ iff $|R| < \omega$.
- (2) *Every f.g. infinite ring has a minimal quotient.*
- (3) *Every injective module over a minimal ring is faithful.*
- (4) *Every variety of modules has an SI injective.*

Corollary

*If $[\text{Cg}(w, x), \text{Cg}(y, z)] = 0$ is definable, and \mathcal{V} omits type **1**, then $\chi_{\mathcal{V}} \neq \omega$.*