

KERNELS OF EPIMORPHISMS OF FINITELY GENERATED FREE LATTICES

WILLIAM DEMEO, PETER MAYR, AND NIK RUŠKUC

1. INTRODUCTION

1.1. Bounded Homomorphisms. The notation and definitions we use follow those of Freese, Jezek, Nation [FJN95], although they have their origins in the earlier work of Ralph McKenzie (reference needed).

We define a pair of closure operators, denoted by superscripts $^{\wedge}$ and $^{\vee}$, on subsets of an arbitrary lattice $\mathbf{L} = \langle L, \vee, \wedge \rangle$ as follows: For each $A \subseteq L$, let

$$A^{\wedge} = \{ \bigwedge B : B \text{ is a finite subset of } A \}.$$

We adopt the following convention: if \mathbf{L} has a greatest element $1_{\mathbf{L}}$, then $\bigwedge \emptyset = 1_{\mathbf{L}}$, and we include this in A^{\wedge} for every $A \subseteq L$. (For lattices without a greatest element, $\bigwedge \emptyset$ is undefined.) The set A^{\vee} is defined dually.

If $\mathbf{K} = \langle K, \vee, \wedge \rangle$ is a lattice generated by a finite set X , then K is the union of a chain of subsets $H_0 \subseteq H_1 \subseteq \dots$ defined inductively by setting $H_0 := X^{\wedge}$ and $H_{k+1} := (H_k)^{\vee\wedge}$, for all $k \geq 0$. By induction, each $H_n = X^{\wedge(\vee\wedge)^n}$ is a finite meet-closed subset of K , and $\bigcup H_n = K$, since X generates \mathbf{K} .

Let $h: \mathbf{K} \rightarrow \mathbf{L}$ be a lattice epimorphism and define, for each $y \in L$ and $k < \omega$,

$$\beta_k(y) = \bigwedge \{ w \in H_k : h(w) \geq y \}.$$

On page 30 of [FJN95], immediately after Theorem 2.4, the authors make the following remark, which is a crucial ingredient of our proof:

"...[h is lower bounded] if and only if for each $a \leq h(1_{\mathbf{K}})$ there exists $N \in \omega$ such that $\beta_n(a) = \beta_N(a)$ for all $n \geq N$."

Fact. The following are equivalent:

- (1) h is not lower bounded;
- (2) $(\exists y_0 \in L)(\forall N)(\exists n > N) \beta_n(a) \neq \beta_N(a)$;
- (3) $(\exists y_0 \in L)(\exists N)(\forall n > N) \beta_n(a) \neq \beta_N(a)$.

wjd 2018-08-30: To do: verify this!

2. MAIN THEOREM

Let X be a finite set and $\mathbf{F} := \mathbf{F}(X)$ the free lattice generated by X .

Theorem 2.1. *Suppose $\mathbf{L} = \langle L, \wedge, \vee \rangle$ is a finite lattice and $h: \mathbf{F} \rightarrow \mathbf{L}$ a lattice epimorphism. If h is bounded then the kernel of h is a finitely generated sublattice of $\mathbf{F} \times \mathbf{F}$.*

Date: 2018-08-30.

Proof. Assume h is bounded. That is, the preimage of each $y \in L$ under h is bounded. For each $y \in L$, let $\alpha y = \bigvee h^{-1}\{y\}$ and $\beta y = \bigwedge h^{-1}\{y\}$ denote the greatest and least elements of $h^{-1}\{y\}$, respectively (both of which exist by the boundedness assumption). Observe that $h\alpha h = h$, and $h\beta h = h$. In fact, α and β are adjoint to h . Indeed, it is easy to see that

$$\begin{aligned} hx \leq y &\Leftrightarrow x \leq \alpha y, \\ y \leq hx &\Leftrightarrow \beta y \leq x. \end{aligned}$$

For each $y \in L$, let $X_y := X \cap h^{-1}\{y\}$, the set of generators that lie in the inverse image of y under h . Let G be the (finite) set of pairs in $\mathbf{F} \times \mathbf{F}$ defined as follows:

$$G = \bigcup_{y \in L} \{(x, \alpha y), (\alpha y, x), (x, \beta y), (\beta y, x), (\alpha y, \beta y), (\beta y, \alpha y) : x \in X_y\}.$$

We claim that G generates $\ker h$. To prove this, we first show, by induction on term complexity, that for every $y \in L$, for every $r \in h^{-1}\{y\}$, the pairs $(r, \alpha y)$ and $(r, \beta y)$ belong to the sublattice $\langle G \rangle \leq \mathbf{F} \times \mathbf{F}$ generated by G .

Case 0. If $r \in X$, then $(r, \alpha y)$ and $(r, \beta y)$ belong to G itself, so there's nothing to prove.

Case 1. Suppose $r = s \vee t$ and assume (the induction hypothesis) that $(s, \alpha h(s))$, $(s, \beta h(s))$, $(t, \alpha h(t))$, and $(t, \beta h(t))$ belong to $\langle G \rangle$. Then $y = h(r) = h(s \vee t) = h(s) \vee h(t)$, so

$$h(\alpha h(s) \vee \alpha h(t)) = h\alpha h(s) \vee h\alpha h(t) = h(s) \vee h(t) = y.$$

Similarly, $h(\beta h(s) \vee \beta h(t)) = h(s) \vee h(t) = y$. Therefore,

$$\beta y \leq \beta h(s) \vee \beta h(t) \leq \alpha h(s) \vee \alpha h(t) \leq \alpha y.$$

Also, $r \leq \alpha y$, so $r = \alpha y \wedge (s \vee t)$. Taken together, these observations yield

$$\begin{aligned} \begin{pmatrix} r \\ \beta y \end{pmatrix} &= \begin{pmatrix} \alpha y \wedge (s \vee t) \\ \beta y \end{pmatrix} = \begin{pmatrix} \alpha y \wedge (s \vee t) \\ \beta y \wedge (\beta h(s) \vee \beta h(t)) \end{pmatrix} \\ &= \begin{pmatrix} \alpha y \\ \beta y \end{pmatrix} \wedge \left[\begin{pmatrix} s \\ \beta h(s) \end{pmatrix} \vee \begin{pmatrix} t \\ \beta h(t) \end{pmatrix} \right], \end{aligned}$$

and each term in the last expression belongs to $\langle G \rangle$, so $(r, \beta y) \in \langle G \rangle$, as desired.

Similarly, $(r, \alpha y) \in \langle G \rangle$. Indeed, $\beta y \leq r$ implies $r = \beta y \vee s \vee t$, and $\beta h(s) \vee \beta h(t) \leq \alpha y$ implies $\alpha y = \alpha y \vee \beta h(s) \vee \beta h(t)$. Therefore,


$$\begin{pmatrix} r \\ \alpha y \end{pmatrix} = \begin{pmatrix} \beta y \vee s \vee t \\ \alpha y \vee \beta h(s) \vee \beta h(t) \end{pmatrix} = \begin{pmatrix} \beta y \\ \alpha y \end{pmatrix} \vee \begin{pmatrix} s \\ \beta h(s) \end{pmatrix} \vee \begin{pmatrix} t \\ \beta h(t) \end{pmatrix}.$$

Case 2. Suppose $r = s \wedge t$ and assume $(s, \alpha h(s))$, $(s, \beta h(s))$, $(t, \alpha h(t))$, and $(t, \beta h(t))$ belong to $\langle G \rangle$. Then $h(s \wedge t) = h(r) = y$, so $h(\alpha h(s) \wedge \alpha h(t)) = y = h(\beta h(s) \wedge \beta h(t))$, so

$$\beta y \leq \beta h(s) \wedge \beta h(t) \leq \alpha h(s) \wedge \alpha h(t) \leq \alpha y.$$

Also, $\beta y \leq r \leq \alpha y$ so $r = \alpha y \wedge s \wedge t$ and $r = \beta y \vee (s \wedge t)$. Taken together, these observations yield

$$\begin{pmatrix} r \\ \alpha y \end{pmatrix} = \begin{pmatrix} \beta y \vee (s \wedge t) \\ \alpha y \vee (\alpha h(s) \wedge \alpha h(t)) \end{pmatrix} = \begin{pmatrix} \beta y \\ \alpha y \end{pmatrix} \vee \left[\begin{pmatrix} s \\ \alpha h(s) \end{pmatrix} \wedge \begin{pmatrix} t \\ \alpha h(t) \end{pmatrix} \right],$$


 FIGURE 1. The free lattice over M_3 .

and each term in the last expression belongs to $\langle Y \rangle$.

Note, we could have used β 's instead:

$$\begin{pmatrix} r \\ \alpha y \end{pmatrix} = \begin{pmatrix} \beta y \vee (s \wedge t) \\ \alpha y \vee (\beta h(s) \wedge \beta h(t)) \end{pmatrix} = \begin{pmatrix} \beta y \\ \alpha y \end{pmatrix} \vee \left[\begin{pmatrix} s \\ \beta h(s) \end{pmatrix} \wedge \begin{pmatrix} t \\ \beta h(t) \end{pmatrix} \right].$$

Similarly,

$$\begin{pmatrix} r \\ \beta y \end{pmatrix} = \begin{pmatrix} \alpha y \wedge s \wedge t \\ \beta y \wedge \alpha h(s) \wedge \alpha h(t) \end{pmatrix} = \begin{pmatrix} \alpha y \\ \beta y \end{pmatrix} \wedge \begin{pmatrix} s \\ \alpha h(s) \end{pmatrix} \wedge \begin{pmatrix} t \\ \alpha h(t) \end{pmatrix}.$$

Again, we could have used β 's instead:

$$\begin{pmatrix} r \\ \beta y \end{pmatrix} = \begin{pmatrix} \alpha y \wedge s \wedge t \\ \beta y \wedge \beta h(s) \wedge \beta h(t) \end{pmatrix} = \begin{pmatrix} \alpha y \\ \beta y \end{pmatrix} \wedge \begin{pmatrix} s \\ \beta h(s) \end{pmatrix} \wedge \begin{pmatrix} t \\ \beta h(t) \end{pmatrix}.$$

In each case, we end up with an expression involving terms from $\langle G \rangle$, and this proves that $(r, \alpha y)$ and $(r, \beta y)$ belong to $\langle G \rangle$. \square

We conjecture the converse of Theorem 2.1. Suppose $\mathbf{L} = \langle L, \wedge, \vee \rangle$ is a finite lattice and $h: \mathbf{F} \rightarrow \mathbf{L}$ a lattice epimorphism. If the kernel of h is a finitely generated sublattice of $\mathbf{F} \times \mathbf{F}$, then h is bounded. If we could assume that whenever h is unbounded then there is a class of $\ker h$ containing both an infinite chain and a generator of \mathbf{F} , then there is a straightforward proof of the conjecture. (See the Appendix for details.) Unfortunately, as the next result shows, this assumption is not always valid.

Proposition 2.2. *Let $\mathbf{F} = \mathbf{F}(x, y, z)$, and let $\mathbf{L} = \mathbf{F}_{M_3}(3)$ (see Figure 1). Let $h: \mathbf{F} \rightarrow \mathbf{L}$ be an epimorphism. Then $K = \ker h$ is not finitely generated.*

Proof. Define the sequences $\{m_n\}$, $\{x_n\}$, $\{y_n\}$, $\{z_n\}$ ($n < \omega$) of elements of $\mathbf{F}(X)$ as follows:

Let $x_0 = x$, $y_0 = y$, $z_0 = z$, and for $n \geq 0$,

$$\begin{aligned} m_n &= (x_n \wedge y_n) \vee (x_n \wedge z_n) \vee (y_n \wedge z_n); \\ x_{n+1} &= x_n \vee m_n = x_n \vee (y_n \wedge z_n). \end{aligned}$$

Define y_{n+1} and z_{n+1} similarly.

Claim 2.3. If $\{s_n\}$ is any one of the four sequences just defined, then for $n > 0$, we have $s_{n+1} > s_n$ and $h(s_{n+1}) = h(s_n)$.

□

jd 2018-08-30: To
o: fill in proof!

3. EXAMPLES

Let $\mathbf{M}_3 = \langle \{0, a, b, c, 1\}, \wedge, \vee \rangle$, where $a \wedge b = a \wedge c = b \wedge c = 0$ and $a \vee b = a \vee c = b \vee c = 1$. Let $\mathbf{F} := \mathbf{F}(x, y, z)$ denote the free lattice generated by $\{x, y, z\}$.

Proposition 3.1. *Let $h: \mathbf{F} \rightarrow \mathbf{M}_3$ be the epimorphism that acts on the generators as follows: $x \mapsto a$, $y \mapsto b$, $z \mapsto c$. Then $\ker h$ is not finitely generated.*

Proof. Let $K := \ker h$, and for $u \in \{x, y, z\}$ let $C_u := u/K := \{v \in F : h(v) = h(u)\}$. Define sequences of elements in these classes by the following mutual recursions:

- for $i \in \mathbb{N}$,

$$m_{0,i} = (m_{x,i} \wedge m_{y,i}) \vee (m_{x,i} \wedge m_{z,i}) \vee (m_{y,i} \wedge m_{z,i});$$

- for $u \in \{x, y, z\}$,

$$\begin{aligned} m_{u,0} &= u, \\ m_{u,i+1} &= m_{u,i} \vee m_{0,i}. \end{aligned}$$

Notice that $m_{0,0} = (x \wedge y) \vee (x \wedge z) \vee (y \wedge z)$ and $m_{x,i+1} = m_{x,i} \vee (m_{y,i} \wedge m_{z,i})$.

Let X be a finite subset of K . We will prove there exists $(p, q) \in K \setminus \langle X \rangle$. Fix $u \in \{0, x, y, z\}$. Since X is finite, Lemma 3.3 implies that there exists $M \in \mathbb{N}$ such that for every $(p, q) \in X$ with $p, q \in C_u$, we have $p, q \leq m_{u,M}$.

Subclaim 1. For $(p, q) \in \langle X \rangle$ and $u \in \{x, y, z\}$, the following implication holds:

$$q \leq u \implies p \leq m_{u,M}. \quad (3.1)$$

We prove the subclaim by induction on the complexity of terms. Fix $(p, q) \in \langle X \rangle$. Then $p, q \in C_u$ for some $u \in \{x, y, z\}$.

- **Case 0.** Suppose $(p, q) \in X$. Then by definition of M we have $p, q \leq m_{u,M}$.
- **Case 1.** Suppose $(p, q) = (p_1, q_1) \wedge (p_2, q_2)$, where (p_i, q_i) satisfies (3.1) for $i = 1, 2$. If $q = q_1 \wedge q_2 \leq u$, then, since generators in the free lattice are meet-prime (see Theorem B.8 below), we have $q_1 \leq u$ or $q_2 \leq u$. Assume $q_1 \leq u$. Then, by the induction hypothesis, $p_1 \leq m_{u,M}$. Therefore, $p = p_1 \wedge p_2 \leq m_{u,M}$, as desired.

- **Case 2.** Suppose $(p, q) = (p_1, q_1) \vee (p_2, q_2)$, where (p_i, q_i) satisfies (3.1) for $i = 1, 2$. If $q = q_1 \vee q_2 \leq u$, then $q_i \leq u$ for $i = 1, 2$. It now follows from the induction hypothesis that $p_i \leq m_{u,M}$ for $i = 1, 2$, so $p = p_1 \vee p_2 \leq m_{u,M}$, as desired.

This completes the proof of Subclaim 1.

It follows from the subclaim just proved and Lemma 3.2 that $(m_{x,M+1}, x) \in K \setminus \langle X \rangle$, so the proof of the proposition is complete. \square

Lemma 3.2. *For each $u \in \{0, x, y, z\}$, the sequence $\{m_{u,n} : n \in \mathbb{N}\}$ is a strictly ascending chain; that is, $m_{u,0} < m_{u,1} < m_{u,2} < \dots$.*

Proof. We split the proof up into cases: either $u \in \{x, y, z\}$, or $u = 0$.

- **Case 1.** $u \in \{x, y, z\}$.

For simplicity, assume $u = x$ for the remainder of the proof of this case. (Of course, the same argument goes through when u is y or z .) Fix $n \in \mathbb{N}$. We prove $m_{x,n} < m_{x,n+1}$.

Subclaim 2. For all $n \in \mathbb{N}$,

- (1) $m_{x,n} \in C_x$,
- (2) $m_{x,n} \not\geq y$, and $m_{x,n} \not\geq z$.

Proof of Subclaim 2. The first item is obvious; for the second, if $m_{x,n} \geq y$, then $m_{x,n} \wedge y = y$, and then $0 = h(m_{x,n} \wedge y) = h(y) = b$. A similar contradiction is reached if we assume $m_{x,n} \geq z$, so the subclaim is proved.

Recall, $m_{x,n} = m_{x,n} \vee (m_{y,n} \wedge m_{z,n})$, so our desired conclusion, $m_{x,n} < m_{x,n+1}$, holds unless $m_{x,n} \geq m_{y,n} \wedge m_{z,n}$. So, by way of contradiction, suppose

$$m_{x,n} \geq m_{y,n} \wedge m_{z,n}. \quad (3.2)$$

Now, $m_{y,n} = y \vee (x \wedge z) \vee \dots$, so clearly $m_{y,n} \geq y$. Similarly, $m_{z,n} \geq z$. This, together with (3.2), implies $m_{x,n} \geq m_{y,n} \wedge m_{z,n} \geq y \wedge z$. But then Theorem B.8 below implies that either $m_{x,n} \geq y$ or $m_{x,n} \geq z$, which contradicts Subclaim 2.

- **Case 2.** $u = 0$.

We first prove that $m_{0,0} = (x \wedge y) \vee (x \wedge z) \vee (y \wedge z)$ is strictly below $m_{0,1} = (m_{x,1} \wedge m_{y,1}) \vee (m_{x,1} \wedge m_{z,1}) \vee (m_{y,1} \wedge m_{z,1})$.

By symmetry, it suffices to show $x \wedge y < m_{x,1} \wedge m_{y,1}$; that is, $x \wedge y < (x \vee (y \wedge z)) \wedge (y \vee (x \wedge z))$.

Clearly $x \wedge y \leq (x \vee (y \wedge z)) \wedge (y \vee (x \wedge z))$. Suppose $x \wedge y = (x \vee (y \wedge z)) \wedge (y \vee (x \wedge z))$. Then $(x \vee (y \wedge z)) \wedge (y \vee (x \wedge z)) \leq x$. By Theorem B.8, the latter holds iff $x \vee (y \wedge z) \leq x$ or $y \vee (x \wedge z) \leq x$. The first of these inequalities is clearly false, so it must be the case that $y \vee (x \wedge z) \leq x$. But then $y \leq x$, which is obviously false. We conclude that $x \wedge y < (x \vee (y \wedge z)) \wedge (y \vee (x \wedge z))$.

This proves $m_{0,0} < m_{0,1}$.

Now fix $n \in \mathbb{N}$ and assume $m_{0,n} < m_{0,n+1}$. We show $m_{0,n+1} < m_{0,n+2}$.

(**To do:** complete the proof in this case; i.e., for $u = 0$.)

—scratch work—

$$m_{0,n} := (m_{x,n} \wedge m_{y,n}) \vee (m_{x,n} \wedge m_{z,n}) \vee (m_{y,n} \wedge m_{z,n}),$$

$$m_{0,n+1} := (m_{x,n+1} \wedge m_{y,n+1}) \vee (m_{x,n+1} \wedge m_{z,n+1}) \vee (m_{y,n+1} \wedge m_{z,n+1}),$$

By the first Case above, $m_{u,n} < m_{u,n+1}$.

□

Lemma 3.3. *For all $u \in \{x, y, z\}$ and $p \in C_u \cup C_0$ there exists $n \in \mathbb{N}$ such that $p \leq m_{u,n}$.*

Proof. (By induction on the complexity of p .)

- **Case 0.** $p \in \{x, y, z\}$. Then $u = p = m_{p,0}$.

For the remaining cases assume $u = x$, without loss of generality.

- **Case 1.** $p = p_1 \vee p_2$.

If $p \in C_x \cup C_0$, then $p_i \in C_x \cup C_0$ for $i = 1, 2$, and the induction hypothesis yields i and j for which $p_1 \leq m_{x,i}$ and $p_2 \leq m_{x,j}$. Letting $n = \max\{i, j\}$, we have $p_1, p_2 \leq m_{x,n}$, from which $p = p_1 \vee p_2 \leq m_{x,n}$, as desired.

- **Case 2.** $p = p_1 \wedge p_2$.

If $p \in C_x$, then we may assume $p_1 \in C_x$ and $p_2 \in C_x \cup C_0$. By the induction hypothesis, there exists $n \in \mathbb{N}$ such that $p_1 \leq m_{x,n}$, whence $p \leq p_1 \leq m_{x,n}$. If $p \in C_0$, then each p_i belongs to $C_u \cup C_0$ for some $u \in \{x, y, z\}$. If $p_1 \in C_x \cup C_0$, then $p_1 \leq m_{x,n}$, as above and we're done. Similarly, if $p_2 \in C_x \cup C_0$. So assume $p_1 \in C_y \cup C_0$ and $p_2 \in C_z \cup C_0$. Then the induction hypothesis implies that there exist i and j such that $p_1 \leq m_{y,i}$ and $p_2 \leq m_{z,j}$. If $n = \max\{i, j\}$, then $p_1 \leq m_{y,n}$ and $p_2 \leq m_{z,n}$. Then, by the above definition of the sequences, we have $p_1 \wedge p_2 \leq m_{y,n} \wedge m_{z,n} \leq m_{0,n} \leq m_{x,n+1}$.

□

3.1. Other Examples. In each of the propositions in this section, X is a finite set and $\mathbf{F} = \mathbf{F}(X)$ is the free lattice generated by X . The symbol F denotes the universe of \mathbf{F} . The proof in each case is straightforward, but tedious; we omit proofs of the first two, and give a detailed proof of the third.

Prop. 3.4. *Let $X = \{x, y, z\}$, and let $\mathbf{L} = \mathbf{2}$ be the 2-element chain. Then the kernel of an epimorphism $h: \mathbf{F} \twoheadrightarrow \mathbf{L}$ is a finitely generated sublattice of $\mathbf{F} \times \mathbf{F}$.*

Prop. 3.5. *Let $X = \{x, y, z\}$ and let $\mathbf{L} = \mathbf{3}$ be the 3-element chain. Then the kernel of an epimorphism $h: \mathbf{F} \twoheadrightarrow \mathbf{L}$ is finitely generated.*

Prop. 3.6. *Let $n > 2$, $X = \{x_0, x_1, \dots, x_{n-1}\}$, and $\mathbf{L} = \mathbf{2} \times \mathbf{2}$. Let $h: \mathbf{F} \twoheadrightarrow \mathbf{L}$ be an epimorphism. Then $K = \ker h$ is finitely generated.*

Proof. Let the universe of $\mathbf{L} = \mathbf{2} \times \mathbf{2}$ be $\{0, a, b, 1\}$, where $a \vee b = 1$ and $a \wedge b = 0$. For each $y \in L$, denote by $X_y = X \cap h^{-1}\{y\}$ the set of generators mapped by h to y . Denote the least and greatest elements of $h^{-1}\{y\}$ (if they exist) by ℓ_y and g_y , respectively. For example, $\ell_a = \bigwedge h^{-1}\{a\} = \bigwedge \{x \in F : h(x) = a\}$, $g_b = \bigvee \{x \in F : h(x) = b\}$, etc. In the present example, the least and greatest elements exist in each case, as we now show.

Subclaim 3. $h^{-1}\{a\}$ has least and greatest elements, namely $\ell_a = \bigwedge (X_a \cup X_1)$ and $g_a = \bigvee (X_a \cup X_0)$. (Similarly, $h^{-1}\{b\}$ has least and greatest elements, ℓ_b and g_b .)

Proof of Subclaim 3. Let $M(a) := \bigwedge (X_a \cup X_1)$ and $J(a) := \bigvee (X_a \cup X_0)$ and note that these values exist in F , since the sets involved are finite. Also, Then $h(M(a)) = a = h(J(a))$. Fix $r \in h^{-1}\{a\}$.

- If $r \in X_a$, then $r \geq \bigwedge X_a \geq \bigwedge (X_a \cup X_1) = M(a)$.
- If $r = s \vee t$, where $h(s) = a$ and $h(t) \in \{a, 0\}$, then assume (the induction hypothesis) that $s \geq M(a)$, and we have $r = s \vee t \geq M(a)$.
- If $r = s \wedge t$, where $h(s) = a$ and $h(t) \in \{a, 1\}$, then assume (the induction hypothesis) that $s, t \geq M(a)$, and we have $s \wedge t \geq M(a)$. This proves that for each $r \in h^{-1}\{a\}$ we have $r \geq M(a)$, and as we noted at the outset, $M(a) \in h^{-1}\{a\}$. Therefore, $\ell_a = M(a)$ is the least element of $h^{-1}\{a\}$. Similarly, every $r \in h^{-1}\{a\}$ is below $J(a)$, so $g_a = J(a)$. The proofs of $\ell_b = M(b)$ and $g_b = J(b)$ are similar.

This proves Subclaim 3.

Subclaim 4. $h^{-1}\{0\}$ has least and greatest elements, namely, $\ell_0 = \bigwedge X$ and $g_0 = g_a \wedge g_b$.

Proof of Subclaim 4. $\ell_0 = \bigwedge X$ is obvious, so we need only verify that $g_0 = g_a \wedge g_b$. Observe that $h(g_a \wedge g_b) = h(g_a) \wedge h(g_b) = a \wedge b = 0$, so $g_a \wedge g_b \in h^{-1}\{0\}$. It remains to prove that $r \leq g_a \wedge g_b$ holds for all $r \in h^{-1}\{0\}$. Fix $r \in h^{-1}\{0\}$. Then $h(r \vee g_a) = h(r) \vee h(g_a) = 0 \vee a = a$, which places $r \vee g_a$ in $h^{-1}\{a\}$. Therefore, by maximality of g_a , we have $r \vee g_a \leq g_a$, whence $r \leq g_a$. Similarly, $r \leq g_b$. This proves Subclaim 4.

Subclaim 5. $h^{-1}\{1\}$ has least and greatest elements, namely $\ell_1 = \ell_a \vee \ell_b$ and $g_1 = \bigvee X$.

Proof of Subclaim 5. $g_1 = \bigvee X$ is obvious, so we need only verify that $\ell_1 = \ell_a \vee \ell_b$. Observe that $h(\ell_a \vee \ell_b) = h(\ell_a) \vee h(\ell_b) = a \vee b = 1$, so $\ell_a \vee \ell_b \in h^{-1}\{1\}$. It remains to prove that $r \geq \ell_a \vee \ell_b$ holds for all $r \in h^{-1}\{1\}$. Fix $r \in h^{-1}\{1\}$. Then $h(r \wedge \ell_a) = h(r) \wedge h(\ell_a) = 1 \wedge a = a$, which places $r \wedge \ell_a$ in $h^{-1}\{a\}$. Therefore, by minimality of ℓ_a , we have $r \wedge \ell_a \geq \ell_a$, whence $r \geq \ell_a$. Similarly, $r \geq \ell_b$. Now let $Y = \{(x, g_p), (g_p, x), (x, \ell_p), (\ell_p, x) : p \in \{0, a, b, 1\}, x \in X_p\}$. This proves Subclaim 5.

Subclaim 6. If $r \in F$ and $h(r) = p$, then $(r, \ell_p), (r, g_p) \in \langle Y \rangle$.

Proof of Subclaim 6. Either $r \in X_p$ or $r = s \wedge t$ or $r = s \vee t$. If $r \in X_p$, then the pair belongs to Y and the claim is trivial.

Suppose $r = s \wedge t$.

- If $h(r) = 1$, then $h(s) = h(t) = 1$. If we assume (the induction hypothesis) that $(s, \ell_1), (s, g_1), (t, \ell_1), (t, g_1)$ belong to $\langle Y \rangle$, then $(r, \ell_1) = (s \wedge t, \ell_1) = (s, \ell_1) \wedge (t, \ell_1) \in \langle Y \rangle$.
- If $h(r) = a$, then (wlog) $h(s) = a$ and $h(t) \in \{a, 1\}$. Assume (the induction hypothesis) that $(s, \ell_a), (s, g_a), (t, \ell_p), (t, g_p)$ belong to $\langle Y \rangle$. By Claim 1, $\ell_a \leq \ell_1$, so $\ell_a = \ell_a \wedge \ell_1$.
 - If $h(t) = 1$, then $(r, \ell_a) = (s \wedge t, \ell_a \wedge \ell_1) = (s, \ell_a) \wedge (t, \ell_1) \in \langle Y \rangle$.
 - If $h(t) = a$, then $(r, \ell_a) = (s \wedge t, \ell_a \wedge \ell_a) = (s, \ell_a) \wedge (t, \ell_a) \in \langle Y \rangle$.

- If $h(r) = 0$, then (wlog) that either (i) $h(s) = 0$, or (ii) $h(s) = a$, $h(t) = b$. If $h(s) = 0$, then $(s, \ell_0) \in \langle Y \rangle$ implies $(r, \ell_0) = (s \wedge t, \ell_0) = (s, \ell_0) \wedge (t, \ell_p) \in \langle Y \rangle$. If $h(s) = a$, $h(t) = b$, and $(s, \ell_a), (t, \ell_b) \in \langle Y \rangle$, then $(r, \ell_0) = (s \wedge t, \ell_0) = (s, \ell_a) \wedge (t, \ell_b) \in \langle Y \rangle$.

Similarly, in each of these three subcases we have $(r, g_p) \in \langle Y \rangle$.

Suppose $r = s \vee t$.

- If $h(r) = 0$, then $h(s) = h(t) = 0$. If we assume (the induction hypothesis) that $(s, \ell_p), (s, g_p), (t, \ell_p), (t, g_p)$ belong to $\langle Y \rangle$, then $(r, \ell_p) = (s \vee t, \ell_p) = (s, \ell_p) \vee (t, \ell_p) \in \langle Y \rangle$.
- If $h(r) = a$, then (wlog) $h(s) = a$ and $h(t) \in \{a, 0\}$. If we assume (the induction hypothesis) that $(s, \ell_p), (s, g_p), (t, \ell_p), (t, g_p)$ belong to $\langle Y \rangle$, then $(r, \ell_p) = (s \vee t, \ell_p) = (s, \ell_p) \vee (t, \ell_p) \in \langle Y \rangle$.
- If $h(r) = 1$, then (wlog) that either (i) $h(s) = 1$, or (ii) $h(s) = a$, $h(t) = b$.
 - If $h(s) = 1$, then $(s, \ell_1) \in \langle Y \rangle$ implies $(r, \ell_1) = (s \vee t, \ell_1) = (s, \ell_1) \vee (t, \ell_p) \in \langle Y \rangle$.
 - If $h(s) = a$, $h(t) = b$, and $(s, \ell_a), (t, \ell_b) \in \langle Y \rangle$, then $(r, \ell_1) = (s \vee t, \ell_1) = (s, \ell_a) \vee (t, \ell_b) \in \langle Y \rangle$.

Similarly, in each of these three subcases, we have $(r, g_p) \in \langle Y \rangle$. This proves Subclaim 6, and completes the proof of Prop 3.6. \square

4. MISCELLANEOUS NOTES

Let K be a finite subset of $\ker h$. Since K is finite, we can find an $N < \omega$ such that for all $\binom{p}{q} \in K$, the following implications are satisfied:

$$\begin{aligned} p \leq x &\implies q \leq x_N \\ p \leq y &\implies q \leq y_N \\ p \leq z &\implies q \leq z_N \end{aligned} \tag{4.1}$$

$$\begin{aligned} p \leq x \vee (y \wedge z) &\implies q \leq x_{N+1} \\ p \leq y \vee (x \wedge z) &\implies q \leq y_{N+1} \\ p \leq z \vee (x \wedge y) &\implies q \leq z_{N+1} \end{aligned} \tag{4.2}$$

Claim 4.3 If N is chosen as just described, and if $\binom{p}{q} \in \langle K \rangle$ then the implications 4.1 and 4.2 hold.

Proof. As usual, we proceed by induction on term complexity. If $\binom{p}{q} \in K$, then by choice of N , there is nothing to prove.

Case 1. Suppose $\binom{p}{q} = \binom{p_1}{q_1} \vee \binom{p_2}{q_2}$, where $\binom{p_1}{q_1}$ and $\binom{p_2}{q_2}$ satisfy (4.1) and (4.2). We show that $\binom{p}{q}$ satisfies these two implications as well. Recall, in the notation above, $x_1 := x \vee (y \wedge z)$.

Assume $p \leq x_1$. We show $q \leq x_{N+1}$. Since $p = p_1 \vee p_2 \leq x_1$, we have $p_1 \leq x_1$ and $p_2 \leq x_1$, so by the induction hypothesis, $q_1 \leq x_{N+1}$ and $q_2 \leq x_{N+1}$. Therefore, $q = q_1 \vee q_2 \leq x_{N+1}$, as desired.

Now assume $p \leq x$. We show $q \leq x_N$. Since $p = p_1 \vee p_2 \leq x$, we have $p_1 \leq x$ and $p_2 \leq x$, so by the induction hypothesis, $q_1 \leq x_N$ and $q_2 \leq x_N$. Therefore, $q = q_1 \vee q_2 \leq x_N$, as desired.

Case 2. Suppose $\binom{p}{q} = \binom{p_1}{q_1} \wedge \binom{p_2}{q_2}$, where $\binom{p_1}{q_1}$ and $\binom{p_2}{q_2}$ satisfy (4.1) and (4.2).

Assume $p \leq x_1 = x \vee (y \wedge z)$. We must show $q \leq x_{N+1}$. Since $p_1 \wedge p_2 \leq x_1$, then according to Theorem B.8, at least one of the following inequalities must hold:

- (1) $p_1 \leq x_1$;
- (2) $p_2 \leq x_1$;
- (3) $p_1 \wedge p_2 \leq x$;
- (4) $p_1 \wedge p_2 \leq y \wedge z$.

By the induction hypothesis, (1) implies $q_1 \leq x_{N+1}$ and (2) implies $q_2 \leq x_{N+1}$. In either case, $q = q_1 \wedge q_2 \leq x_{N+1}$, as desired. In case (3), Theorem B.8 implies that either $p_1 \leq x$ or $p_2 \leq x$, since x is a generator. Therefore, $q_1 \leq x_N$ or $q_2 \leq x_N$ and we conclude that $q \leq x_N \leq x_{N+1}$, as desired. It remains to prove $q \leq x_{N+1}$ for the final case in which $p_1 \wedge p_2 \leq y \wedge z$.

If $p_1 \wedge p_2 \leq y \wedge z$, then $p_1 \wedge p_2 \leq y$ and $p_1 \wedge p_2 \leq z$. Therefore, both of the following disjunctions hold:

- $p_1 \leq y$ or $p_2 \leq y$, and
- $p_1 \leq z$ or $p_2 \leq z$.

If $p_1 \leq y$ and $p_1 \leq z$, then $p_1 \leq x \vee (y \wedge z) = x_1$, so $q_1 \leq x_{N+1}$, so $q = q_1 \wedge q_2 \leq x_{N+1}$, as desired. Similarly, if $p_2 \leq y$ and $p_2 \leq z$, the desired conclusion holds. Finally, consider the case in which $p_1 \leq y$ and $p_2 \leq z$. In this case $q_1 \leq y_N$ and $q_2 \leq z_N$. Therefore, $q = q_1 \wedge q_2 \leq y_N \wedge z_N \leq x_N \vee (y_N \wedge z_N) = x_{N+1}$, as desired. \square

APPENDIX A. PROOF OF CONJECTURE UNDER SPECIAL ASSUMPTIONS

Prop. A.1. *Suppose $\mathbf{L} = \langle L, \wedge, \vee \rangle$ is a finite lattice and $h: \mathbf{F} \twoheadrightarrow \mathbf{L}$ a lattice epimorphism. Suppose also that whenever h is unbounded then there is a class of $\ker h$ containing both an infinite chain and a generator of \mathbf{F} . Then h is bounded whenever its kernel is a finitely generated sublattice of $\mathbf{F} \times \mathbf{F}$.*

Suppose h is not lower bounded. Then by Fact 1.1 there is an element $y_0 \in L$ such that $\beta_0(y_0) > \beta_1(y_0) > \dots$ is an infinite descending chain.

Let K be a finite subset of $\ker h$, say, $K = \{(p_1, q_1), \dots, (p_m, q_m)\} \subseteq \ker h$. We prove $\langle K \rangle \neq \ker h$. (Since K is an arbitrary finite subset of $\ker h$, this will prove $\ker h$ is not finitely generated.)

Let $x_0 \in X$ be a generator of \mathbf{F} that belongs to the class $h^{-1}\{y_0\}$ (so, $h(x_0) = y_0$).

Claim 1.1. There exists $N < \omega$ such that for all (p_i, q_i) in K , if $p_i \geq x_0$, then $q_i \geq \beta_N(y_0)$.

Proof. Fix i and $(p_i, q_i) \in K$ (so, $h(p_i) = h(q_i)$). Define N_i as follows:

Case 0. If $p_i \not\geq x_0$, let $N_i = 0$.

Case 1. If $p_i \geq x_0$, then $x_0 = x_0 \wedge p_i$, so $y_0 = h(x_0) = h(x_0) \wedge h(p_i) \leq h(p_i)$, so $y_0 \leq h(q_i)$. Also, $h(x_0 \wedge q_i) = h(x_0) \wedge h(q_i) = y_0$, so $x_0 \wedge q_i \in h^{-1}\{y_0\}$. Therefore (since $\{\beta_i(y_0)\}$ is an infinite descending chain in $h^{-1}\{y_0\}$) there exists $n_i > 0$ such that $x_0 \wedge q_i \geq \beta_{n_i}(y_0)$. Let $N_i = n_i$ in this case (so $q_i \geq \beta_{N_i}(y_0)$).

Since K is finite, we can find such N_i for each $(p_i, q_i) \in K$. Let $N = \max\{N_i : 1 \leq i \leq m\}$. Then for all $1 \leq i \leq m$ the following implication holds:

$$p_i \geq x_0 \implies q_i \geq \beta_N(y_0). \quad (\text{A.1})$$

Claim 1.2. There exists $N < \omega$ such that, for all $(p, q) \in \langle K \rangle$,

$$p \geq x_0 \implies q \geq \beta_N(y_0). \quad (\text{A.2})$$

Proof. Choose N as described in the proof of Claim 1.1 above so that for all $(p_i, q_i) \in K$ the implication (A.1) holds. Fix $(p, q) \in \langle K \rangle$. We prove (A.2) by induction on the complexity of (p, q) . If $(p, q) \in K$, then there's nothing to prove.

Case 1. Assume $(p, q) = (p_1, q_1) \wedge (p_2, q_2)$, where p_i, q_i ($i = 1, 2$) satisfy (A.2). Assume $p \geq x_0$. Then $p = p_1 \wedge p_2 \geq x_0$, so $p_1 \geq x_0$ and $p_2 \geq x_0$, so (by the induction hypothesis) $q_1 \geq \beta_N(y_0)$ and $q_2 \geq \beta_N(y_0)$. Therefore, $q = q_1 \wedge q_2 \geq \beta_N(y_0)$, as desired.

Case 2. Assume $(p, q) = (p_1, q_1) \vee (p_2, q_2)$, where p_i, q_i ($i = 1, 2$) satisfy (A.2). Assume $p \geq x_0$. Then $p = p_1 \vee p_2 \geq x$. Since x_0 is a generator, it is join prime in $\mathbf{F}(X)$, so either $p_1 \geq x_0$ or $p_2 \geq x_0$. Assume (wlog) $p_1 \geq x_0$. Then, (by induction hypothesis) $q_1 \geq \beta_N(y_0)$. Therefore, $q = q_1 \vee q_2 \geq q_1 \geq \beta_N(y_0)$, as desired.

Claim 1.3. K does not generate $\ker h$.

Proof. Let N be chosen as in the proof of Claim 1.2 above. Since $\beta_0(y_0) > \beta_1(y_0) > \dots$ is an infinite descending chain, $\beta_N(y_0) > \beta_{N+1}(y_0)$. The pair $(p, q) = (x_0, \beta_{N+1}(y_0))$ does not belong to $\langle K \rangle$, however it does belong to the kernel of h . This proves that the finite subset K does not generate $\ker h$. Since K was an arbitrary finite subset of $\ker h$, we have proved that $\ker h$ is not finitely generated.

APPENDIX B. BACKGROUND

Here are some useful definitions and results from the Free Lattices book by Freese, Jezek, and Nation [FJN95].

Definition B.1 (length of a term). Let X be a set. Each element of X is a term of length 1, also known as a *variable*. If t_1, \dots, t_n are terms of lengths k_1, \dots, k_n , then $t_1 \vee \dots \vee t_n$ and $t_1 \wedge \dots \wedge t_n$ are both terms of length $1 + k_1 + \dots + k_n$.

Examples. By the above definition, the terms

$$x \vee y \vee z \quad x \vee (y \vee z) \quad (x \vee y) \vee z$$

have lengths 4, 5, and 5, respectively. Reason: variables have length 1, so $x \vee y \vee z$ has length $1 + 1 + 1 + 1$. On the other hand, $x \vee y$ is a term of length 3, so $(x \vee y) \vee z$ has length $1 + 3 + 1$. Similarly, $x \vee (y \vee z)$ has length $1 + 1 + 3$.

Lemma B.2 ([FJN95, Lem. 1.2]). Let \mathcal{V} be a nontrivial variety of lattices and let $\mathbf{F}_{\mathcal{V}}(X)$ be the relatively free lattice in \mathcal{V} over X . Then,

$$\bigwedge S \leq \bigvee T \text{ implies } S \cap T \neq \emptyset \text{ for each pair of finite subsets } S, T \subseteq X. \quad (\text{B.1})$$

Lemma B.3 ([FJN95, Lem. 1.4]). Let \mathbf{L} be a lattice generated by a set X and let $a \in L$. Then

- (1) if a is join prime, then $a = \bigwedge S$ for some finite subset $S \subseteq X$.
- (2) if a is meet prime, then $a = \bigvee S$ for some finite subset $S \subseteq X$.

If X satisfies condition (B.1) above, then

- (3) for every finite, nonempty subset $S \subseteq X$, $\bigwedge S$ is join prime and $\bigvee S$ is meet prime.

Corollary B.4 ([FJN95, Cor. 1.5]). Let \mathcal{V} be a nontrivial variety of lattices and let $\mathbf{F}_{\mathcal{V}}(X)$ be the relatively free lattice in \mathcal{V} over X . For each finite nonempty subset $S \subseteq X$, $\bigwedge S$ is join prime and $\bigvee S$ is meet prime. In particular, every $x \in X$ is both join and meet prime. Moreover, if $x \leq y$ for $x, y \in X$, then $x = y$.

Theorem B.5 (Whitman's Condition, ver. 1). The free lattice $\mathbf{F}(X)$ satisfies the following condition:

(W) If $v = v_1 \wedge \dots \wedge v_r \leq u_1 \vee \dots \vee u_s = u$, then either $v_i \leq u$ for some i , or $v \leq u_j$ for some j .

Corollary B.6 ([FJN95, Cor. 1.9]). Every sublattice of a free lattice satisfies (W). Every element of a lattice satisfying (W) is either join or meet irreducible.

Theorem B.7 (Whitman's Condition, ver. 2). *The free lattice $\mathbf{F}(X)$ satisfies the following condition:*

(W+) *If $v = v_1 \wedge \cdots \wedge v_r \wedge x_1 \wedge \cdots \wedge x_n \leq u_1 \vee \cdots \vee u_s \vee y_1 \vee \cdots \vee y_m = u$, where $x_i, y_j \in X$, then either $x_i = y_j$ for some i and j , or $v_i \leq u$ for some i , or $v \leq u_j$ for some j .*

Theorem B.8 ([FJN95, Thm. 1.11]). *If $s = s(x_1, \dots, x_n)$ and $t = t(x_1, \dots, x_n)$ are terms and $x_1, \dots, x_n \in X$, then the truth of*

$$s^{\mathbf{F}(X)} \leq t^{\mathbf{F}(X)} \quad (\text{B.2})$$

can be determined by applying the following rules.

- (1) *If $s = x_i$ and $t = x_j$, then (B.2) holds iff $x_i = x_j$.*
- (2) *If $s = s_1 \vee \cdots \vee s_k$ is a formal join, then (B.2) holds iff $s_i^{\mathbf{F}(X)} \leq t^{\mathbf{F}(X)}$ for all i .*
- (3) *If $t = t_1 \wedge \cdots \wedge t_k$ is a formal meet, then (B.2) holds iff $s^{\mathbf{F}(X)} \leq t_i^{\mathbf{F}(X)}$ for all i .*
- (4) *If $s = x_i$ and $t = t_1 \vee \cdots \vee t_k$ is a formal join, then (B.2) holds iff $x_i \leq t_j^{\mathbf{F}(X)}$ for some j .*
- (5) *If $s = s_1 \wedge \cdots \wedge s_k$ is a formal meet and $t = x_i$, then (B.2) holds iff $s_j^{\mathbf{F}(X)} \leq x_i$ for some j .*
- (6) *If $s = s_1 \wedge \cdots \wedge s_k$ is a formal meet and $t = t_1 \vee \cdots \vee t_m$ is a formal join, then (B.2) holds iff $s_i^{\mathbf{F}(X)} \leq t^{\mathbf{F}(X)}$ for some i or $s^{\mathbf{F}(X)} \leq t_j^{\mathbf{F}(X)}$ for some j .*

Definition B.9 (up directed, continuous). A subset A of a lattice L is said to be *up directed* if every finite subset of A has an upper bound in A . It suffices to check this for pairs. A is up directed iff for all $a, b \in A$ there exists $c \in A$ such that $a \leq c$ and $b \leq c$. A lattice is *upper continuous* if whenever $A \subseteq L$ is an up directed set having a least upper bound $u = \bigvee A$, then for every b ,

$$\bigvee_{a \in A} (a \wedge b) = \bigvee_{a \in A} a \wedge b = u \wedge b.$$

Down directed and *down continuous* are defined dually. A lattice that is both up and down continuous is called *continuous*.

Theorem B.10 ([FJN95, Thm. 1.22]). *Free lattices are continuous.*

REFERENCES

- [FJN95] Ralph Freese, Jaroslav Ježek, and J. B. Nation. *Free lattices*, volume 42 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1995. URL: <http://dx.doi.org/10.1090/surv/042>, doi:10.1090/surv/042.

Email address: williamdemeo@gmail.com