

Representing Finite Lattices as Congruence Lattices

William DeMeo, Ralph Freese, Peter Jipsen

<http://math.hawaii.edu/~ralph/>

<http://uacalc.org/>

<https://github.com/UACalc/>

BLAST, Vanderbilt University, Aug 14–18, 2017

The Problem

Theorem (Grätzer-Schmidt)

*Every algebraic (so every finite) lattice is isomorphic to **Con** (**A**) for some (unary) algebra **A**.*

The Problem

Theorem (Grätzer-Schmidt)

Every algebraic (so every finite) lattice is isomorphic to $\mathbf{Con}(\mathbf{A})$ for some (unary) algebra \mathbf{A} .

Problem

*Is every finite \mathbf{L} isomorphic to $\mathbf{Con}(\mathbf{A})$ for some **finite** \mathbf{A} ?*

The Problem

Theorem (Grätzer-Schmidt)

Every algebraic (so every finite) lattice is isomorphic to $\mathbf{Con}(\mathbf{A})$ for some (unary) algebra \mathbf{A} .

Problem

*Is every finite \mathbf{L} isomorphic to $\mathbf{Con}(\mathbf{A})$ for some **finite** \mathbf{A} ?*

Since $\mathbf{Con}(\mathbf{A}) = \mathbf{Con}(\langle \mathbf{A}, \text{Pol}_1(\mathbf{A}) \rangle)$, **we assume all algebras are unary.**

Properties

Possible representation properties for a finite lattice \mathbf{L} :

(P1) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$.

Properties

Possible representation properties for a finite lattice \mathbf{L} :

- (P1) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$.
- (P2) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the all nonconstant operations are permutations.

Properties

Possible representation properties for a finite lattice \mathbf{L} :

- (P1) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$.
- (P2) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the all nonconstant operations are permutations.
- (P3) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the nonconstant operations generate a transitive permutation group.

Properties

Possible representation properties for a finite lattice \mathbf{L} :

- (P1) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$.
- (P2) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the all nonconstant operations are permutations.
- (P3) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the nonconstant operations generate a transitive permutation group.
- (P4) \mathbf{L} is isomorphic to an interval in the lattice of subgroups of a finite group.

Properties

Possible representation properties for a finite lattice \mathbf{L} :

- (P1) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$.
- (P2) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the all nonconstant operations are permutations.
- (P3) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the nonconstant operations generate a transitive permutation group.
- (P4) \mathbf{L} is isomorphic to an interval in the lattice of subgroups of a finite group.

$$(P3) \Rightarrow (P2) \Rightarrow (P1)$$

Properties

Possible representation properties for a finite lattice \mathbf{L} :

- (P1) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$.
- (P2) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the all nonconstant operations are permutations.
- (P3) \mathbf{L} is isomorphic to the congruence lattice of some finite algebra $\langle A, F \rangle$ where the nonconstant operations generate a transitive permutation group.
- (P4) \mathbf{L} is isomorphic to an interval in the lattice of subgroups of a finite group.

$$(P4) \Leftrightarrow (P3) \Rightarrow (P2) \Rightarrow (P1)$$

$(P3) \Leftrightarrow (P4)$

Let H be a subgroup of G . Let

$$A = \{aH : a \in G\} \quad (\text{left cosets})$$

$(P3) \Leftrightarrow (P4)$

Let H be a subgroup of G . Let

$$A = \{aH : a \in G\} \quad (\text{left cosets})$$

Make an algebra \mathbf{A} by adding operations

$$g : aH \mapsto gaH \quad (\text{left multiplication})$$

$(P3) \Leftrightarrow (P4)$

Let H be a subgroup of G . Let

$$A = \{aH : a \in G\} \quad (\text{left cosets})$$

Make an algebra \mathbf{A} by adding operations

$$g : aH \mapsto gaH \quad (\text{left multiplication})$$

Then **Con** $A \cong [H, G]$, the interval in the subgroup lattice.

Theorem (1980)

(P1) holds for *all* lattices iff (P4) holds for *all* lattices.

Theorem (1980)

(P1) holds for *all* lattices iff (P4) holds for *all* lattices.


The **size** of a representation $\mathbf{L} \cong \mathbf{Con}(\mathbf{A})$ is $|A|$. For $\mathbf{H} \leq \mathbf{G}$ the size in (P4) is $[G : H]$, the number of left H -cosets of \mathbf{G} .

Theorem (1980)

(P1) holds for *all* lattices iff (P4) holds for *all* lattices.

The **size** of a representation $\mathbf{L} \cong \mathbf{Con}(\mathbf{A})$ is $|A|$. For $\mathbf{H} \leq \mathbf{G}$ the size in (P4) is $[G : H]$, the number of left H -cosets of \mathbf{G} .

Example. The minimum size for \mathbf{L}_6 is 6:




Theorem (1980)

(P1) holds for *all* lattices iff (P4) holds for *all* lattices.

The **size** of a representation $\mathbf{L} \cong \mathbf{Con}(\mathbf{A})$ is $|A|$. For $\mathbf{H} \leq \mathbf{G}$ the size in (P4) is $[G : H]$, the number of left H -cosets of \mathbf{G} .

Example. The minimum size for \mathbf{L}_6 is 6:

\mathbf{L}_6




\mathbf{B}_6	0 1 2 3 4 5
$f(x)$	2 2 1 5 5 4
$g(x)$	3 4 4 0 1 1
$h(x)$	4 5 3 4 5 3

Theorem (1980)

(P1) holds for *all* lattices iff (P4) holds for *all* lattices.

The **size** of a representation $\mathbf{L} \cong \mathbf{Con}(\mathbf{A})$ is $|\mathbf{A}|$. For $\mathbf{H} \leq \mathbf{G}$ the size in (P4) is $[G : H]$, the number of left H -cosets of \mathbf{G} .

Example. The minimum size for \mathbf{L}_6 is 6:

\mathbf{L}_6		\mathbf{B}_6	0 1 2 3 4 5
		$f(x)$	2 2 1 5 5 4
		$g(x)$	3 4 4 0 1 1
		$h(x)$	4 5 3 4 5 3

Pálffy and Aschbacher have found groups $\mathbf{H} \leq \mathbf{G}$ representing this lattice. But Pálffy's example has $\mathbf{G} = \mathbf{A}_{11}$ and $|H| = 55$, so the size is $9! = 362880$.

Moral

Moral: Finding a representation with groups, (P4), may be much harder (and much bigger) than finding a (P1) representation.

New representable lattices from old

- all distributive lattices

New representable lattices from old

- all distributive lattices
- lattice duals (Hans Kurzweil, 1985, and R. Netter, 1986)

New representable lattices from old

- all distributive lattices
- lattice duals (Hans Kurzweil, 1985, and R. Netter, 1986)
- interval sublattices

New representable lattices from old

- all distributive lattices
- lattice duals (Hans Kurzweil, 1985, and R. Netter, 1986)
- interval sublattices
- direct products (Jiří Tůma, 1986)

New representable lattices from old

- all distributive lattices
- lattice duals (Hans Kurzweil, 1985, and R. Netter, 1986)
- interval sublattices
- direct products (Jiří Tůma, 1986)
- ordinal sums (Ralph McKenzie, 1984; John Snow, 2000)

New representable lattices from old

- all distributive lattices
- lattice duals (Hans Kurzweil, 1985, and R. Netter, 1986)
- interval sublattices
- direct products (Jiří Tůma, 1986)
- ordinal sums (Ralph McKenzie, 1984; John Snow, 2000)
- parallel sums (John Snow, 2000)

New representable lattices from old

- all distributive lattices
- lattice duals (Hans Kurzweil, 1985, and R. Netter, 1986)
- interval sublattices
- direct products (Jiří Tůma, 1986)
- ordinal sums (Ralph McKenzie, 1984; John Snow, 2000)
- parallel sums (John Snow, 2000)
- sublattices of representable lattices obtained as a union of a filter and an ideal (John Snow, 2000)

New representable lattices from old

- all distributive lattices
- lattice duals (Hans Kurzweil, 1985, and R. Netter, 1986)
- interval sublattices
- direct products (Jiří Tůma, 1986)
- ordinal sums (Ralph McKenzie, 1984; John Snow, 2000)
- parallel sums (John Snow, 2000)
- sublattices of representable lattices obtained as a union of a filter and an ideal (John Snow, 2000)
- overalgebras (DeMeo, 2013)

Pálffy-Pudlák Conditions

(A) \mathbf{L} is simple.

Pálffy-Pudlák Conditions

- (A) L is simple.
- (B) For each $x \neq 0$ in L , there are elements y and z such that $x \vee y = x \vee z = 1$ and $y \wedge z = 0$.

Pálffy-Pudlák Conditions

- (A) L is simple.
- (B) For each $x \neq 0$ in L , there are elements y and z such that $x \vee y = x \vee z = 1$ and $y \wedge z = 0$.
- (C) $|L| \neq 2$ and each element of L that is not an atom or 0 contains at least four atoms.

Pálffy-Pudlák Conditions

- (A) \mathbf{L} is simple.
- (B) For each $x \neq 0$ in L , there are elements y and z such that $x \vee y = x \vee z = 1$ and $y \wedge z = 0$.
- (C) $|L| \neq 2$ and each element of L that is not an atom or 0 contains at least four atoms.

Theorem

- If \mathbf{L} satisfies (A) and (B) then \mathbf{L} satisfies (P1) \Rightarrow (P2).

Pálffy-Pudlák Conditions

- (A) \mathbf{L} is simple.
- (B) For each $x \neq 0$ in L , there are elements y and z such that $x \vee y = x \vee z = 1$ and $y \wedge z = 0$.
- (C) $|L| \neq 2$ and each element of L that is not an atom or 0 contains at least four atoms.

Theorem

- If \mathbf{L} satisfies (A) and (B) then \mathbf{L} satisfies $(P1) \Rightarrow (P2)$.
- If \mathbf{L} satisfies (A), (B) and (C) then \mathbf{L} satisfies $(P1) \Rightarrow (P3)$.

McKenzie's variants

(B') If $\varphi : L \rightarrow L$ is any meet-preserving map such that $\varphi(x) > x$ for $x \neq 1$, then $\varphi(x) = 1$ for all x .

McKenzie's variants

- (B') If $\varphi : L \rightarrow L$ is any meet-preserving map such that $\varphi(x) > x$ for $x \neq 1$, then $\varphi(x) = 1$ for all x .
- (B'') The coatoms of L meet to 0.

McKenzie's variants

- (B') If $\varphi : L \rightarrow L$ is any meet-preserving map such that $\varphi(x) > x$ for $x \neq 1$, then $\varphi(x) = 1$ for all x .
- (B'') The coatoms of L meet to 0.

$$(B) \implies (B'') \implies (B').$$

McKenzie's variants

- (B') If $\varphi : L \rightarrow L$ is any meet-preserving map such that $\varphi(x) > x$ for $x \neq 1$, then $\varphi(x) = 1$ for all x .
- (B'') The coatoms of L meet to 0.

$$(B) \implies (B'') \implies (B').$$

Theorem

- *If \mathbf{L} satisfies (A) and (B') (or (B'')) then a minimal representation of \mathbf{L} witnesses that \mathbf{L} satisfies (P2). So,*

McKenzie's variants

- (B') If $\varphi : L \rightarrow L$ is any meet-preserving map such that $\varphi(x) > x$ for $x \neq 1$, then $\varphi(x) = 1$ for all x .
- (B'') The coatoms of L meet to 0.

$$(B) \implies (B'') \implies (B').$$

Theorem

- If \mathbf{L} satisfies (A) and (B') (or (B'')) then a minimal representation of \mathbf{L} witnesses that \mathbf{L} satisfies (P2). So,
- If (A) and (B') hold and $\mathbf{L} \cong \langle A, F \rangle$ is minimal, then F consists of permutations and constants.

Representations by intransitive groups

Suppose $\mathbf{A} = \langle A, G \rangle$ is a G -set and let $\mathbf{A}_i = \langle A_i, G \rangle$, $i < k$, be the minimal subalgebras of \mathbf{A} ; i.e. each set A_i is an orbit, or one-generated subuniverse, of \mathbf{A} .

Representations by intransitive groups

Suppose $\mathbf{A} = \langle A, G \rangle$ is a G -set and let $\mathbf{A}_i = \langle A_i, G \rangle$, $i < k$, be the minimal subalgebras of \mathbf{A} ; i.e. each set A_i is an orbit, or one-generated subuniverse, of \mathbf{A} .

Define congruences on A by the partitions

$$\tau = |A_0|A_1| \cdots |A_{k-1}| \quad (\text{the blocks are the orbits})$$

$$\tau_i = |A_i| \quad (\text{at most one nontrivial block})$$

$$\gamma_i = |A_i|A - A_i| \quad (\text{exactly two blocks unless } A_i = A)$$

Representations by intransitive groups

Suppose $\mathbf{A} = \langle A, G \rangle$ is a G -set and let $\mathbf{A}_i = \langle A_i, G \rangle$, $i < k$, be the minimal subalgebras of \mathbf{A} ; i.e. each set A_i is an orbit, or one-generated subuniverse, of \mathbf{A} .

Define congruences on A by the partitions

$$\tau = |A_0|A_1| \cdots |A_{k-1}| \quad (\text{the blocks are the orbits})$$

$$\tau_i = |A_i| \quad (\text{at most one nontrivial block})$$

$$\gamma_i = |A_i|A - A_i| \quad (\text{exactly two blocks unless } A_i = A)$$

We call τ the **intransitivity congruence**;

Representations by intransitive groups

Suppose $\mathbf{A} = \langle A, G \rangle$ is a G -set and let $\mathbf{A}_i = \langle A_i, G \rangle$, $i < k$, be the minimal subalgebras of \mathbf{A} ; i.e. each set A_i is an orbit, or one-generated subuniverse, of \mathbf{A} .

Define congruences on A by the partitions

$$\tau = |A_0|A_1| \cdots |A_{k-1}| \quad (\text{the blocks are the orbits})$$

$$\tau_i = |A_i| \quad (\text{at most one nontrivial block})$$

$$\gamma_i = |A_i|A - A_i| \quad (\text{exactly two blocks unless } A_i = A)$$

We call τ the **intransitivity congruence**;

Theorem

Let $\theta \in \mathbf{Con}(\mathbf{A})$, where $\mathbf{A} = \langle A, G \rangle$ and G is a group. Then

$$\tau = |A_0|A_1| \cdots |A_{k-1}| \quad (\text{the blocks are the orbits})$$

1 G acts transitively if and only if $\tau = 1_A$.

$$\tau = |A_0| |A_1| \cdots |A_{k-1}| \quad (\text{the blocks are the orbits})$$

- 1 G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- 2 The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- 1 G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- 2 The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- 3 The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(\mathbf{A}_i)$.

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(\mathbf{A}_i)$.
For $\theta_i \in \mathbf{Con}(\mathbf{A}_i)$, map $(\theta_0, \dots, \theta_{k-1}) \mapsto \theta_0 \cup \dots \cup \theta_{k-1}$.

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(\mathbf{A}_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
If $\theta \not\leq \gamma_i$, there are $a \in A_i$ and $b \notin A_i$ with $(a, b) \in \theta$. Since G acts transitively on each orbit, $\tau_i \leq \theta$. So $\tau \leq \theta$.

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(\mathbf{A}_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .

Follows from (3) and (4).

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .
- ⑥ If $k > 1$ and $|A_i| = 1$ for all i except 0 then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .

$\tau = |A_0| |A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .
- ⑥ If $k > 1$ and $|A_i| = 1$ for all i except 0 then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑦ If $k > 1$ and $[0_{\mathbf{A}}, \tau]$ is directly indecomposable then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .
- ⑥ If $k > 1$ and $|A_i| = 1$ for all i except 0 then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑦ If $k > 1$ and $[0_{\mathbf{A}}, \tau]$ is directly indecomposable then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑧ If $k = 2$ and $|A_1| = 1$ then τ is a coatom and everything is comparable with it.

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .
- ⑥ If $k > 1$ and $|A_i| = 1$ for all i except 0 then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑦ If $k > 1$ and $[0_{\mathbf{A}}, \tau]$ is directly indecomposable then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑧ If $k = 2$ and $|A_1| = 1$ then τ is a coatom and everything is comparable with it.

From (2) and (6).

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)


- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .
- ⑥ If $k > 1$ and $|A_i| = 1$ for all i except 0 then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑦ If $k > 1$ and $[0_{\mathbf{A}}, \tau]$ is directly indecomposable then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑧ If $k = 2$ and $|A_1| = 1$ then τ is a coatom and everything is comparable with it.
- ⑨ If τ is a coatom and $[0_{\mathbf{A}}, \tau]$ is directly indecomposable then everything is comparable with it.

$\tau = |A_0|A_1| \cdots |A_{k-1}|$ (the blocks are the orbits)

- ① G acts transitively if and only if $\tau = 1_{\mathbf{A}}$.
- ② The interval $[\tau, 1_{\mathbf{A}}]$ is isomorphic to $\text{Eq}(k)$.
- ③ The interval $[0_{\mathbf{A}}, \tau]$ is isomorphic to $\prod_{i=0}^{k-1} \mathbf{Con}(A_i)$.
- ④ If, for some i , $\theta \geq \bigvee_{j \neq i} \tau_j$ then $\theta \geq \tau$ or $\theta \leq \gamma_i$.
- ⑤ If $\theta \wedge \tau \prec \tau$ then $\theta \leq \gamma_i$ for some i .
- ⑥ If $k > 1$ and $|A_i| = 1$ for all i except 0 then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑦ If $k > 1$ and $[0_{\mathbf{A}}, \tau]$ is directly indecomposable then every coatom of $\mathbf{Con}(\mathbf{A})$ lies above τ .
- ⑧ If $k = 2$ and $|A_1| = 1$ then τ is a coatom and everything is comparable with it.
- ⑨ If τ is a coatom and $[0_{\mathbf{A}}, \tau]$ is directly indecomposable then everything is comparable with it.

From (8).


Examples: L_{14}



Example

- L_{14} satisfies (A) and (B'') so a minimal representation is permutational.


Examples: L_{14}



Example

- L_{14} satisfies (A) and (B'') so a minimal representation is permutational.


Examples: \mathbf{L}_{14}



Example

- \mathbf{L}_{14} satisfies (A) and (B'') so a minimal representation is permutational.
- $\mathbf{L}_{14} \cong \mathbf{Con} \langle A, G \rangle$ is not possible if G acts intransitively,


Examples: \mathbf{L}_{14}



Example

- \mathbf{L}_{14} satisfies (A) and (B'') so a minimal representation is permutational.
- $\mathbf{L}_{14} \cong \mathbf{Con} \langle A, G \rangle$ is not possible if G acts intransitively, so
- if $\mathbf{Con} \langle A, F \rangle$ is a minimal representation, then F generates a transitive group.


Examples: \mathbf{L}_{14}



Example

- \mathbf{L}_{14} satisfies (A) and (B'') so a minimal representation is permutational.
- $\mathbf{L}_{14} \cong \mathbf{Con} \langle A, G \rangle$ is not possible if G acts intransitively, so
- if $\mathbf{Con} \langle A, F \rangle$ is a minimal representation, then F generates a transitive group.
- Is \mathbf{L}_{14} representable? (Yes: as $[H, A_6]$ with $[A_6 : H] = 90$)


Examples: \mathbf{L}_{14}




Example

- \mathbf{L}_{14} satisfies (A) and (B'') so a minimal representation is permutational.
- $\mathbf{L}_{14} \cong \mathbf{Con} \langle A, G \rangle$ is not possible if G acts intransitively, so
- if $\mathbf{Con} \langle A, F \rangle$ is a minimal representation, then F generates a transitive group.
- Is \mathbf{L}_{14} representable? (Yes: as $[H, A_6]$ with $[A_6 : H] = 90$)
- Is this a minimum representation? (Don't know)


Examples: \mathbf{L}_{15} , (the dual of \mathbf{L}_{14})



Examples: \mathbf{L}_{15} , (the dual of \mathbf{L}_{14})




Examples: \mathbf{L}_{15} , (the dual of \mathbf{L}_{14})



Example

- $\mathbf{L}_{15} \cong \mathbf{Con} \langle \{0, 1, 2, 3\}, G \rangle$, G the group generated by the double transposition $0 \leftrightarrow 1, 2 \leftrightarrow 3$.


Examples: \mathbf{L}_{15} , (the dual of \mathbf{L}_{14})




Example

- $\mathbf{L}_{15} \cong \mathbf{Con} \langle \{0, 1, 2, 3\}, G \rangle$, G the group generated by the double transposition $0 \leftrightarrow 1, 2 \leftrightarrow 3$.
- $\mathbf{L}_{14} \cong \mathbf{L}_{15}^d$, which again proves \mathbf{L}_{14} is representable.

Examples: L_4 .




Examples: L_4 .



Example

- L_4 satisfies (B'') but not (A) so minimal representations need not be permutational.

Examples: \mathbf{L}_4 .



Example

- \mathbf{L}_4 satisfies (B'') but not (A) so minimal representations need not be permutational. In fact
- $\mathbf{L}_4 \cong \langle \{0, 1, 2, 3\}, f, g \rangle$, where

\mathbf{B}_4	0	1	2	3
$f(x)$	1	0	3	2
$g(x)$	0	0	2	2

Examples: L_4 .




Example


- L_4 satisfies (B'') but not (A) so minimal representations need not be permutational.
- But L_4 does have an intransitive representation on 6:

\mathbf{B}'_4	0	1	2	3	4	5
$f(x)$	1	2	0	4	5	3
$g(x)$	0	2	1	3	5	4

Examples: \mathbf{L}_{19} , a harder example:



Examples: \mathbf{L}_{19} , a harder example:



Lemma

Let $\mathbf{A} = \langle A, G \rangle$ be a finite algebra, where G is an intransitive group of permutations on A . Suppose the intransitivity congruence τ is a coatom. Then there do not exist congruences $0_{\mathbf{A}} < \psi < \theta$ in $\mathbf{Con}(\mathbf{A})$ with $\theta \wedge \tau = 0_{\mathbf{A}}$.

Proof

Lemma

Let $\mathbf{A} = \langle A, G \rangle$ be a finite algebra, where G is an intransitive group of permutations on A . Suppose the intransitivity congruence τ is a coatom. Then there do not exist congruences $0_{\mathbf{A}} < \psi < \theta$ in $\mathbf{Con}(\mathbf{A})$ with $\theta \wedge \tau = 0_{\mathbf{A}}$.


Proof.

Since τ is a coatom, there are exactly two orbits; call them B and C . Since $\theta \wedge \tau = 0_{\mathbf{A}}$, if $(x, y) \in \theta$ then $x = y$ or one is in B and the other is in C . So θ defines a bipartite graph between B and C . Since G acts transitively on both B and C , this graph corresponds to a bijection between B and C . The same applies to ψ . But equivalence relations corresponding to such graphs cannot be comparable. □

Small Lattices

Theorem


All lattices with at most 7 elements can be represented, with the one possible exception of \mathbf{L}_{10} :



Small Lattices

Theorem

All lattices with at most 7 elements can be represented, with the one possible exception of \mathbf{L}_{10} :




If $\mathbf{L}_{10} \cong \langle A, F \rangle$, then F generates a transitive group on A .

Small Lattices

Theorem

All lattices with at most 7 elements can be represented, with the one possible exception of \mathbf{L}_{10} :



If $\mathbf{L}_{10} \cong \langle A, F \rangle$, then F generates a transitive group on A .

Proof.

\mathbf{L}_{10} satisfies (A) and (B''). By part (5) of the intransitivity theorem, it cannot be represented with an intransitive group. \square

Finding Reps: Methods and Algorithms

Finding Reps: Methods and Algorithms

- Closure Method

Finding Reps: Methods and Algorithms

- Closure Method
- Overalgebras

Finding Reps: Methods and Algorithms

- Closure Method
- Overalgebras
- Ideal-Filter

Finding Reps: Methods and Algorithms

- Closure Method
- Overalgebras
- Ideal-Filter
- Duality

Finding Reps: Methods and Algorithms

- Closure Method
- Overalgebras
- Ideal-Filter
- Duality
- Group Methods (GAP)

Closure Method to find a Representation of \mathbf{L}

Closure Method to find a Representation of \mathbf{L}

- (1) Search through $\text{Eq}(X_k)$, $k = 2, 3, \dots$ finding sublattices isomorphic to \mathbf{L} .

Closure Method to find a Representation of \mathbf{L}

- (1) Search through $\text{Eq}(X_k)$, $k = 2, 3, \dots$ finding sublattices isomorphic to \mathbf{L} .
- (2) For each sublattice $\mathbf{L} \cong \mathbf{L}' \leq \text{Eq}(X_k)$ found, find the unary polymorphs of the members of L' ; that is, calculate the set F of all unary operations on X_k which respect all $\theta \in L'$.

Closure Method to find a Representation of \mathbf{L}

- (1) Search through $\text{Eq}(X_k)$, $k = 2, 3, \dots$ finding sublattices isomorphic to \mathbf{L} .
- (2) For each sublattice $\mathbf{L} \cong \mathbf{L}' \leq \text{Eq}(X_k)$ found, find the unary polymorphs of the members of L' ; that is, calculate the set F of all unary operations on X_k which respect all $\theta \in L'$.
- (3) For F found in the previous step, test if **Con** $(\langle X_k, F \rangle) = L'$. If so then $\mathbf{A} = \langle X_k, F \rangle$ is a minimal representation. Otherwise continue the search.

(a) **Find a small presentation of \mathbf{L} :**

The procedure can be sped up by first finding a presentation of \mathbf{L} with the minimal number of generators. Besides speeding up the search in $\text{Eq}(k)$, it is enough in calculating the unary polymorphs to respect the generators.

(b) **Subdirect Decompositions:**

Subdirect decompositions can be used to speed up finding unary polymorphs. For example, if $\theta_0, \theta_1 \in L' \leq \text{Eq}(X_k)$ with $\theta_0 \wedge \theta_1 = 0$, then X_k is naturally embedded into $X_k/\theta_0 \times X_k/\theta_1$. Since the operations in a direct product are component-wise, this cuts the search space of possible unary polymorphs from k^k down to $r^r s^s$, where r and s are the number of blocks in θ_0 and θ_1 .

(c) **Uniform Equivalence Relations:**


If it can be shown that the algebra of a minimal representation of \mathbf{L} has a transitive permutation group for its nonconstant unary polynomials, then we can restrict our search in $\text{Eq}(k)$ to uniform equivalence relations. Moreover the search for unary polymorphisms can be restricted to permutations.

(d) **Small generating set for the operations:**

Of course if $F' \subseteq F$ is a set of generators for the moniod F , we can take $\mathbf{A} = \langle X_k, F' \rangle$.


Nondist., linearly indec., small lattices

L_1




\mathbf{B}_1	0	1	2	3
$f(x)$	1	0	3	2
$g(x)$	1	0	1	0

L_2




\mathbf{B}_2	0	1	2
$f(x)$	0	1	2

L_3




\mathbf{B}_3	0	1	2	3	4	5	6
$f(x)$	0	1	2	1	2	1	0
$g(x)$	0	3	4	3	4	3	0
$h(x)$	6	5	2	5	2	5	6
$k(x)$	0	1	2	0	0	2	2


Method: overalgebras

L_4 


B_4	0	1	2	3
$f(x)$	1	0	3	2
$g(x)$	0	0	2	2

 L_5 


B₅	0	1	2	3	4	5	6	7	8	9	10	11	
<i>f</i> (<i>x</i>)	1	2	3	4	5	0	7	8	9	10	11	6	
<i>g</i> (<i>x</i>)	6	1	1	10	9	8	7	0	5	4	3	2	1
<i>h</i> (<i>x</i>)	0	0	0	6	0	0	0	6	0	0	0	0	

 L_6 


B_6	0	1	2	3	4	5
$f(x)$	2	2	1	5	5	4
$g(x)$	3	4	4	0	1	1
$h(x)$	4	5	3	4	5	3

L_7 

B_7	0	1	2	3	4	5
$f(x)$	1	0	0	4	3	3
$g(x)$	4	5	5	1	2	2
$h(x)$	3	3	4	3	3	4


 L_8 

B_8	0	1	2	3	4	5
$f(x)$	1	2	0	4	5	3
$g(x)$	3	5	4	0	2	1


 L_9 

B_9	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$f(x)$	0	0	0	0	0	2	1	2	1	3	4	5	3	4	5	
$g(x)$	0	0	0	0	0	6	7	6	7	10	11	12	10	11	12	
$h(x)$	13	14	15	1	9	8	15	14	13	15	1	9	8	8	1	9


Method: overalgebras

L_{10} 


**No finite algebra known with this
as its congruence lattice.**

 L_{11} 

A finite algebra with 108 elements known.

 L_{12} 

B_{12}	0	1	2	3	4	5	6	7	8
$f(x)$	0	0	3	3	3	6	6	6	0
$g(x)$	0	0	8	8	8	1	1	1	0
$h(x)$	0	5	5	4	0	0	5	4	4
$k(x)$	4	2	2	3	4	4	2	3	3
$l(x)$	5	5	7	7	7	6	6	6	5

L_{13} 

B_{13}	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
$f(x)$	0	1	2	1	2	1	0	0	1	2	2	1	0	0	1	2	1	2	0
$g(x)$	0	1	2	0	0	2	2	0	3	4	0	4	4	6	5	2	6	6	2
$h(x)$	0	1	2	3	4	5	6	0	1	2	4	5	6	0	1	2	3	4	6
$k(x)$	7	8	9	3	10	11	12	3	3	3	3	3	3	11	11	11	11	11	11
$l(x)$	13	14	15	16	17	5	18	13	16	17	17	16	13	5	5	5	5	5	5


Method: [overalgebras](#)

 L_{14} 


Upper interval in $\text{Sub}(A_6)$,
algebra of size 90

 L_{15} 

B_{15}	0	1	2	3
$f(x)$	1	0	3	2


 L_{16} 

Upper interval in $\text{Sub}(C_2.A_6)$
algebra of size 180


L_{17} 

B_{17}	0	1	2	3	4	5	6	7	8	9	10	11
$f(x)$	1	0	3	2	5	4	7	6	9	8	11	10
$g(x)$	4	7	5	6	8	11	9	10	0	3	1	2
$h(x)$	0	0	0	0	5	5	5	5	10	10	10	10

Method: filter-ideal in $\text{Sub}(A_4)$

 L_{18} 


Dual of 19, no explicit
small representation known

 L_{19} 

B_{19}	0	1	2	3	4	5	6	7
$f(x)$	0	1	1	0	4	5	5	4
$g(x)$	0	2	3	1	0	2	3	1
$h(x)$	7	6	6	7	3	2	2	3

 L_{20} 

Method: filter-ideal in $\text{SmallGroup}(216,153)$ in GAP

L_{21} 


B_{21}	0	1	2	3	4	5	6	7	8
$f(x)$	3	3	4	8	8	2	2	3	4
$g(x)$	0	0	6	1	1	0	0	5	6
$h(x)$	4	5	5	7	8	8	7	4	4

 L_{22} 


Dual of 23, no explicit small representation known

 L_{23} 


B₂₃	0	1	2	3	4	5
<i>f</i> (<i>x</i>)	0	1	0	1	4	4
<i>g</i> (<i>x</i>)	1	1	3	3	4	5
<i>h</i> (<i>x</i>)	3	2	3	2	5	5
<i>k</i> (<i>x</i>)	4	1	5	3	4	5

 L_{24} 


B_{24}	0	1	2	3
$f(x)$	1	1	2	2
$g(x)$	2	3	3	2

\mathbf{L}_{25} 

\mathbf{B}_{25}	0	1	2	3	4
$f(x)$	0	0	2	2	2
$g(x)$	0	1	0	1	1
$h(x)$	1	1	4	4	4
$k(x)$	2	3	2	3	3


 \mathbf{L}_{26} 

\mathbf{B}_{26}	0	1	2	3	4	5
$f(x)$	1	0	3	2	0	2
$g(x)$	4	4	5	5	1	3
$h(x)$	0	0	0	0	1	1
$k(x)$	3	5	3	5	3	3

L_{27} 


B_{27}	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$f(x)$	0	1	2	3	4	5	0	0	0	0	0	2	2	2	2	2
$g(x)$	4	5	3	4	5	3	5	3	4	5	3	4	5	4	5	3
$h(x)$	2	2	1	5	5	4	2	1	5	5	4	2	2	5	5	4
$k(x)$	3	4	4	0	1	1	4	4	0	1	1	3	4	0	1	1
$l(x)$	0	6	7	8	9	10	6	7	8	9	10	0	6	8	9	10
$m(x)$	11	12	2	13	14	15	12	2	13	14	15	11	12	13	14	15

Method: overalgebras


 L_{28} 

B_{28}	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$f(x)$	0	1	2	3	4	5	0	0	0	0	0	2	2	2	2	2
$g(x)$	3	3	4	3	3	4	3	4	3	3	4	3	3	3	3	4
$h(x)$	1	0	0	4	3	3	0	0	4	3	3	1	0	4	3	3
$k(x)$	4	5	5	1	2	2	5	5	1	2	2	4	5	1	2	2
$l(x)$	0	6	7	8	9	10	6	7	8	9	10	0	6	8	9	10
$m(x)$	11	12	2	13	14	15	12	2	13	14	15	11	12	13	14	15


Method: overalgebras

L_{29} 


B_{29}	0 1 2 3 4
$f(x)$	1 0 3 2 2
$g(x)$	2 4 2 4 3

L_{30} 


B_{30}	0 1 2 3 4
$f(x)$	0 3 4 3 4
$g(x)$	2 2 1 4 3

L_{31} 


B_{31}	0 1 2 3 4
$f(x)$	0 1 1 0 0
$g(x)$	1 1 2 2 2
$h(x)$	3 2 2 4 4

L_{32} 


B_{32}	0 1 2 3 4
$f(x)$	0 1 1 3 3
$g(x)$	1 2 2 4 4
$h(x)$	3 3 4 3 4

L_{33} 

B_{33}	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$f(x)$	1	3	2	0	9	11	10	8	13	15	14	12	5	7	6	4
$g(x)$	11	8	10	9	7	4	6	5	15	12	14	13	3	0	2	1
$h(x)$	14	15	12	13	10	11	8	9	6	7	4	5	2	3	0	1

 L_{34} 

B₃₄	0	1	2	3
f(x)	0	1	3	2

 L_{35} 

B_{35}	0	1	2	3
$f(x)$	1	1	2	3
$g(x)$	2	3	3	3

Problem

- What about nonunary algebras?

Problem

- What about nonunary algebras?

Problem

*Which (finite) lattices can be represented as **Con A**, where **A** has a Taylor term?*

Resources

- [SmallLatticeReps.ua](http://math.hawaii.edu/~ralph/SmallLatticeReps.ua), a UACalc file with most of the \mathbf{B}_i 's:

`http://math.hawaii.edu/~ralph/`

- The slides of this talk are there too.
- [SmallLatticeReps.ua](https://github.com/UACalc/SmallAlgebras/) and other algebra files:

`https://github.com/UACalc/SmallAlgebras/`

- Get UACalc at

`http://uacalc.org/`

- Source at

`https://github.com/UACalc/uacalcsrc/`