

Problems for Discussion 7, 11/06/13

Compiled by Mai Le

1 Questions about overlap add or overlap-save?


2 DSP tricks

Byron is faced with computing **lots** of discrete Fourier transforms. He will, of course, use the FFT algorithm, but he is behind schedule and needs to get his results as quickly as possible. He gets the idea of computing **two** transforms at one time by computing the transform of $s[n] = s_1[n] + js_2[n]$, where $s_1[n]$ and $s_2[n]$ are two real-valued signals of which he needs to compute the spectra. The issue is whether he can retrieve the individual DFTs from the results or not.

- What will the DFT $S[k]$ of this complex-valued signal in terms of $S_1[k]$ and $S_2[k]$, the DFTs of the original signals?
- Byron's friend, an Aggie who knows some signal processing, says that retrieving the wanted DFTs is easy: "Just find the real and imaginary parts of $S[k]$." Show that this approach is too simplistic.
- While his friend's idea is not correct, it does not give him an idea. What approach will work? **Hint:** Use the symmetry properties of the DFT.
- How does the number of computations change with this approach? Will Byron's idea ultimately lead to a faster computation of the required DFTs?

3 Upsampling and the DFT


Let $x[n]$ be a finite-duration sequence of length 8 with the 8-point DFT $X[k]$ shown below.




A new sequence $y[n]$ of length 16 is defined by

$$y[n] = \begin{cases} x\left[\frac{n}{2}\right], & n \text{ even} \\ 0, & n \text{ odd} \end{cases}$$


Which plot corresponds to $Y[k]$, the 16-point DFT of $y[n]$?




(a)



(b)




(c)



(d)



(e)



(f)

4 DFT and the DTFT

Let $x[n]$ be real-valued and non-negative over $0 \leq n \leq N-1$ and zero otherwise. Let $X[k]$ be the N -point DFT of $x[n]$ and $X(e^{j\omega})$ be the DTFT of $x[n]$.

True or false?

If $X(e^{j\omega})$ can be written as

$$X(e^{j\omega}) = B(\omega)e^{j\alpha\omega}$$

where $B(\omega)$ is real and α is a real constant, then $X[k]$ can be expressed as

$$X[k] = A[k]e^{j\gamma k}$$

where $A[k]$ is real and γ is a real constant.

True or false?

If $X[k]$ can be expressed as

$$X[k] = A[k]e^{j\gamma k}$$

where $A[k]$ is real and γ is a real constant, then $X(e^{j\omega})$ can be written as

$$X(e^{j\omega}) = B(\omega)e^{j\alpha\omega}$$

where $B(\omega)$ is real and α is a real constant.

5 Removing Distortion with the DFT

Let a distorted signal $y[n]$ be the output of an LTI system (defined by $h[n]$) on the true signal $x[n]$. Our goal is to recover $x[n]$ from $y[n]$ with knowledge of the filter $h[n]$. In theory, $x[n]$ can be recovered from $y[n]$ by passing $y[n]$ through an inverse filter having a system function equal to the reciprocal of the system function of the distorting filter.

Suppose that the distortion is caused by an FIR filter with impulse response

$$h[n] = \delta[n] - \frac{1}{2}\delta[n - n_0]$$

where n_0 is a positive integer, i.e. the distortion of $x[n]$ take sthe form of an echo at delay n_0 .

- (a) Determine the z-transform $H(z)$ and the N -point DFT $H[k]$ of the impulse response $h[n]$. Assume that $N = 4n_0$.
- (b) Let $H_i(z)$ denote the system inverse function of the inverse filter, and let $h_i[n]$ be the corresponding impulse response. Determine $h_i[n]$. Is this an FIR or an IIR filter? What is the duration of $h_i[n]$?
- (c) Suppose that we use an FIR filter of length N in an attempt to implement the inverse filter, and let the N -point DFT of the FIR filter be

$$G[k] = \frac{1}{H[k]}, \quad k = 0, 1, \dots, N-1$$

What is the impulse response $g[n]$ of the filter?

- (d) It might appear that the FIR filter with DFT $G[k] = \frac{1}{H[k]}$ implements the inverse filter perfectly. After all, one might argue that the FIR distorting filter has an N -point DFT $H[k]$ and the FIR filter in cascade has an N -point DFT $G[k] = \frac{1}{H[k]}$, and since $G[k]H[k] = 1$ for all k , we have implemented an all-pass, nondistorting filter. What is the fallacy in this argument?
- (e) Perform the convolution of $g[n]$ with $h[n]$, and thus determine how well the FIR filter with N -point DFT $G[k] = \frac{1}{H[k]}$ implements the inverse filter.